A Conserved Histone H3-H4 Interface Regulates DNA Damage Tolerance and Homologous Recombination during the Recovery from Replication Stress

Masafumi Hayashi,^{a,b} Kenji Keyamura,^a Asami Yoshida,^a Mariko Ariyoshi,^c Genki Akanuma,^a Takashi Hishida^a

^aDepartment of Molecular Biology, Graduate School of Science, Gakushuin University, Tokyo, Japan

^bDepartment of Life Science, College of Science, Rikkyo University, Tokyo, Japan

^cGraduate School of Frontier Biosciences, Osaka University, Osaka, Japan

ABSTRACT In eukaryotes, genomic DNA is packaged into nucleosomes, which are the basal components coordinating both the structures and functions of chromatin. In this study, we screened a collection of mutations for histone H3/H4 mutants in *Saccharomyces cerevisiae* that affect the DNA damage sensitivity of DNA damage tolerance (DDT)-deficient cells. We identified a class of histone H3/H4 mutations that suppress methyl methanesulfonate (MMS) sensitivity of DDT-deficient cells (referred to here as the histone SDD mutations), which likely cluster on a specific H3-H4 interface of the nucleosomes. The histone SDD mutations did not suppress the MMS sensitivity of DDT-deficient cells in the absence of Rad51, indicating that homologous recombination (HR) is responsible for DNA damage resistance. Furthermore, the histone SDD mutants showed reduced levels of PCNA ubiquitination after exposure to MMS or UV irradiation, consistent with decreased MMS-induced mutagenesis relative to that of wild-type cells. We also found that histone SDD mutants lacking the INO80 chromatin remodeler impair HR-dependent recovery from MMS-induced replication arrest, resulting in defective S-phase progression and increased Rad52 foci. Taken together, our data provide novel insights into nucleosome functions, which link INO80-dependent chromatin remodeling to the regulation of DDT and HR during the recovery from replication blockage.

KEYWORDS DNA damage tolerance, INO80, budding yeast, histones, homologous recombination

Accurate and complete DNA replication is crucial for the maintenance of genome integrity. However, replication fork progression is often impeded by many types of DNA damage arising from endogenous and exogenous sources because replicative DNA polymerases are unable to bypass them (1). Such replication stress can lead to mutagenesis and chromosomal rearrangements, which are hallmarks of cancer (2, 3). Therefore, the ability to overcome replication problems is essential for the faithful transmission of genetic information upon cell division. In eukaryotes, the DNA damage tolerance (DDT) pathway facilitates the bypass of fork-blocking lesions without removing them; this allows the restarting of stalled replication forks and/or repair of single-stranded DNA (ssDNA) gaps generated behind replication forks, thus ensuring the completion of replication (4–8).

The DDT pathway can be subdivided into two parallel branch pathways that can bypass DNA lesions through either translesion DNA synthesis (TLS) or an error-free damage bypass mechanism, both of which are controlled by covalent ubiquitin modification of proliferating cell nuclear antigen (PCNA) (9). In *Saccharomyces cerevisiae*, the Rad6-Rad18, E2-E3 complex plays a crucial role in the DDT pathway via

Citation Hayashi M, Keyamura K, Yoshida A, Ariyoshi M, Akanuma G, Hishida T. 2021. A conserved histone H3-H4 interface regulates DNA damage tolerance and homologous recombination during the recovery from replication stress. *Mol Cell Biol* 41:e00044-20. <https://doi.org/10.1128/MCB.00044-20>.

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Takashi Hishida, takashi.hishida@gakushuin.ac.jp.

Received 9 February 2020

Returned for modification 23 March 2020

Accepted 24 January 2021

Accepted manuscript posted online 1 February 2021

Published 24 March 2021

monoubiquitylation of PCNA at a conserved K164 residue. Monoubiquitylated PCNA activates TLS, during which either the correct or an incorrect nucleotide is incorporated opposite the base that remains at the lesion site (9–12). Indeed, DNA polymerase ζ , involved in the error-prone TLS, is responsible for most spontaneous and induced mutagenesis (13, 14). PCNA can be further modified by the Mms2-Ubc13 E2-conjugating enzyme and the Rad5 E3 ligase, which form a K63-linked polyubiquitin chain (9, 15). This modification of PCNA promotes the error-free bypass of DNA lesions, in which fork-blocking lesions are bypassed by template switching using the undamaged sister chromatid as a template (16–18). Two-dimensional (2D) gel analysis revealed that Rad51-dependent sister chromatid junctions were formed in a PCNA polyubiquitination-dependent manner (19), demonstrating that homologous recombination (HR) contributes to Rad5-Mms2-Ubc13 template switch events during recovery from replication stress. Independent of its role in the DDT pathway, HR is required for the restart of collapsed forks and the repair of postreplicative ssDNA gaps (19–23). In this context, PCNA is modified at K164 and (to a lesser extent) at Lys127 by SUMO during S phase (9). This leads to recruitment of the Srs2 helicase to the stalled replication fork and subsequent inhibition of unscheduled HR events (24–26), which is referred to as the salvage HR pathway. In genetic backgrounds defective in both PCNA ubiquitination and SUMOylation, the salvage HR pathway predominantly facilitates the repair of the ssDNA gap left behind the replication fork, thereby enabling DNA replication to complete during replication stress. However, recent evidence suggests that such HR-dependent recovery from replication stress frequently results in genomic instability (20, 27, 28). Therefore, replication-associated HR function has to be tightly controlled to prevent genomic instability, which could otherwise interfere with replication completion and lead to genomic instability.

In eukaryotes, genomic DNA is packaged into chromatin, which potentially interferes with the accessibility of factors involved in transcription, replication, and repair. Cells have evolved highly coordinated mechanisms to cope with chromatin compaction and accessibility, including covalent modification of histones and ATP hydrolysis-dependent nucleosome remodeling (29–31). ATP-dependent chromatin remodeling complexes can alter the physical state of nucleosomes through their activities, including histone exchange, eviction of nucleosomes, and nucleosome remodeling/sliding (32, 33). The budding yeast INO80 complex is the one of the four classes of nucleosome remodelers and consists of 14 different subunits in addition to the Ino80 Snf2-like ATPase (33, 34). These subunits include three actin-related proteins (Arp4, Arp5, and Arp8) and two RuvB-related helicases (Rvb1 and Rvb2) (35, 36). Accumulating evidence has implicated the INO80 complex in DNA damage and replication stress responses (37, 38). Yeast cells lacking Ino80 or Arp8 are viable but show increased sensitivity to a broad range of genotoxic agents, such as UV light, methyl methanesulfonate (MMS), and IR (ionizing irradiation) (39–42). In budding yeast and mammals, the INO80 complex is recruited to a double-strand-break (DSB) site in a H2A phosphorylation-related and/or cell cycle-regulated manner and promotes both nucleosome eviction around DSBs and DSB-induced chromatin mobility, underscoring its role in DSB repair (43–47). Independent of this role in DSB repair, yeast cells lacking the INO80 complex exhibit slow progression during unperturbed S phase and are unable to resume replication after treatment with hydroxyurea (HU) or MMS (39, 41, 42). In mammalian cells, fork progression is also significantly delayed (42). These observations demonstrate that the INO80 remodeler is involved in normal DNA replication as well as in the recovery from replication stress and plays a direct role in the restart of stalled replication forks.

In this study, we explored the role of nucleosomes in the replication stress response by screening for histone H3/H4 mutants that affect the DNA damage sensitivity of *rad18 Δ* cells, and we identified many histone alleles that have not previously been linked with the DNA damage response. Among these, six H3/H4 alleles that cluster on the lateral surface of the nucleosome suppressed the sensitivity of *rad18 Δ* strains to MMS and/or HU. In particular, histone H3 R69 and H3 E73 residues are part of the α 1

TABLE 1 Histone H3 and H4 mutations that affect the MMS/HU sensitivity of *rad18Δ* cells^a

H3 allele	MMS	HU	H4 allele	MMS	HU
K4A	+	—	R17K	—	++
K4Q	—	+	N25D	s	s
Q5E	+	++	R36A	—	+
A7S	++	++	R36K	++	—
K14A	—	+	R39K	+	+
K14R	—	+	K44A	+	—
K14Q	—	+	K44R	+	—
A15S	—	+	E83A	++	+
K18Q	++	++	V57A	+	—
H39A	++	+	L62A	+	—
R40A	++	++	E63Q	—	+
R40K	+	++	S64D	—	+
K42Q	+	+	T71A	—	+
V46A	+	—	T71D	—	+
R49A	++	++	E74A	—	+
K56A	—	+	H75A	+	—
K56Q	++	++	R78K	—	+
S57D	—	+	R92A	+	+
R69A	—	s	Q93A	+	+
R69K	++	++	R95A	—	+
E73A	s	s	T96A	—	+
E73Q	s	s	L97A	+	+
Q76E	—	s			
R83A	s	—			
I89A	+	+			
S95A	—	++			
Y99F	+	—			
L100A	—	+			
F104A	+	+			
L109A	++	++			
K115R	+	—			
K121A	—	+			
R134A	—	+			

^a “+” and “++” indicate increased MMS and/or HU sensitivity relative to that of *rad18Δ* cells, “s” indicates moderate suppression of MMS and/or HU sensitivity relative to that of *rad18Δ* cells, and “—” indicates MMS and/or HU sensitivity similar to that of *rad18Δ* cells.

helix, which can interact directly with H4 N25; here we refer to this patch as the histone SDD (suppressor of DDT defect). We also show that the INO80 complex and HR pathway are required for the suppression of DDT deficiency by histone SDD mutations. Thus, our data highlight the role of chromatin structure as an important modulator of DDT and HR functions.

RESULTS

Identification of histone H3/H4 mutations that affect the DNA replication stress response. To identify histone residues required for the replicative stress response, we screened a collection of mutations for histone H3 and H4 mutants that affect the DNA damage sensitivity of DDT-deficient *rad18Δ* cells. To this end, we utilized synthetic genetic array (SGA) analysis to construct double mutants between *rad18Δ* and a library of systematic substitutions of histone H3/H4 (48, 49) and successfully constructed 333 *rad18Δ* histone *H3^m/H4^m* double mutants (see Materials and Methods).

The sensitivities of all double mutants to MMS or HU using the spot assay were compared to those of the respective single mutants. Histone mutants that conferred increased sensitivities to either MMS or HU or both in a *rad18Δ* background were retested to confirm their sensitivities, and the results are summarized in Table 1. In total, we identified 49 mutants (28 in H3 and 21 in H4) in this screen. Among these 49 alleles, 18 have been identified as conferring sensitivity to DNA damage in previous large-scale screens (50), which validates the accuracy of our screen. On the other hand, we failed to recover 28 histone mutations that were reported to confer MMS and/or

HU sensitivity in previous studies. This may be due to our use of a different screening strategy and/or our stringent criteria for evaluating MMS and HU sensitivity. Indeed, 26 histone mutants, except for *H4^{I46A}* and *H4^{T80A}* mutants, were less sensitive to MMS than were mutants in *rad18Δ* cells (data not shown), implying that these histone mutants do not cause additive effects in *rad18Δ* cells at the low MMS concentrations used in this screening. In contrast, while *H4^{I46A}* and *H4^{T80A}* mutants showed enhanced sensitivity to MMS and HU, their mutations did not exacerbate the MMS and HU sensitivity of *rad18Δ* cells (Fig. 1A), suggesting that these are epistatic to *rad18Δ* cells. Therefore, it is likely that these are screened out, because the sensitivity of the double mutant in these cases is the same as that of *rad18Δ* cells.

In addition to these MMS- and/or HU-sensitive histone alleles, we found six histone H3/H4 alleles (affecting five amino acid residues) that partially suppressed the sensitivity of *rad18Δ* mutants to MMS and/or HU (*H3^{R69A}*, *H3^{E73A}*, *H3^{E73Q}*, *H3^{Q76E}*, *H3^{R83A}*, and *H4^{N25D}*) (Fig. 1B; Table 1). All five residues are identical to the corresponding residues of human histones H3 and H4, including variants. With the exception of *H3^{R83A}*, these histone single mutants exhibited MMS and HU sensitivities similar to those of wild-type cells (Fig. 1C). In contrast, *H3^{R83A}* cells were slightly more sensitive to MMS than wild-type cells (Fig. 1C), as reported previously (51). We visualized the spatial positions of histone suppressor residues using the yeast nucleosome core structure (Fig. 1D). All residues are clustered in a specific region comprising the lateral surface of the nucleosome. H3 R69 and H3 R83 are located in close proximity to make contacts with the phosphate backbone of nucleosomal DNA (Fig. 1D). Importantly, histone H3 R69 and H3 E73 residues in the $\alpha 1$ helix take part in the H3-H4 interface via interacting directly with H4 N25, thus representing a particular nucleosome patch (Fig. 1D). We refer to this patch as the SDD (suppressor of DDT defect) region. Subsequently, we subjected the H3 R69, H3 E73, and H4 N25 variants to further analysis, with a particular focus on *H3^{R69A}* as a representative histone suppressor mutant.

Histone SDD mutations rescue the DNA damage sensitivity of *pol30^{K164R}* cells.

Rad18 is a central E3 ubiquitin ligase in DDT that is required for ubiquitin modification of PCNA at K164 (9). To further examine the effects of these histone SDD mutants on the DDT pathway, a *pol30^{K164R}* allele was integrated at the *POL30* genomic locus in these histone mutants. The MMS sensitivity of *pol30^{K164R}* cells was partially suppressed by the *H3^{R69A}*, *H3^{E73Q}*, and *H4^{N25D}* mutations (Fig. 2A). We also assessed these histone mutants for their sensitivity to chronic low-dose UV (CLUV) irradiation, because DDT-deficient cells are highly sensitive to this genotoxic insult (52). All histone SDD mutations suppressed the CLUV sensitivity of *pol30^{K164R}* cells (Fig. 2A). In addition, the other histone SDD mutations, *H3^{E73A}*, *H3^{Q76E}* and *H3^{R83A}*, identified in this study also suppressed the MMS sensitivity of *pol30^{K164R}* cells (data not shown).

When cells are exposed to genotoxic agents, the DNA damage checkpoint plays a crucial role in maintaining genome stability by arresting cell cycle progression and promoting DNA damage repair. To test whether histone SDD mutation affects the DNA damage checkpoint, we analyzed damage-induced phosphorylation of Rad53 and H2A (γ -H2A). Similar levels of phosphorylated Rad53 and γ -H2A were detected in wild-type and *H3^{R69A}* cells following MMS treatment (Fig. 2B), indicating that *H3^{R69A}* does not perturb the normal DNA damage checkpoint.

Suppression of MMS and CLUV sensitivity in DDT-deficient cells requires a functional HR repair pathway. Both *SRS2* deletion and lack of PCNA SUMOylation (*pol30^{K127R/K164R}*) suppress the MMS and CLUV sensitivity of DDT-deficient cells by a mechanism that requires a functional HR pathway (24, 25, 52). We therefore hypothesized that *H3^{R69A}* might facilitate HR-dependent recovery from replication stress in DDT-deficient cells. Consistent with this hypothesis, the *H3^{R69A}* mutation did not rescue the MMS and CLUV sensitivity of *pol30^{K164R} rad51Δ* cells (Fig. 2A). Similar behavior was also observed in the *H3^{E73Q}* and *H4^{N25D}* strains (Fig. 2A). These results indicate that the histone SDD mutations can suppress the MMS and CLUV sensitivity of DDT-deficient cells in an HR-dependent manner. Thus, it is likely that the SDD region specifically influences the role of histones in the DDT pathway. We note that the other three alleles,

FIG 1 Identification of histone SDD (suppressor of DDT-deficient cells) mutations. (A) MMS and HU sensitivities of histone *H4^{I46A}* and *H4^{T80A}* mutants. Serial dilutions (10-fold) of equal numbers of cells were spotted onto YPDA plates containing of MMS or HU. The plates were incubated at 30°C for 3 days. (B and C) MMS and HU sensitivities of histone H3/H4 SDD mutants. The relevant strains were spotted and incubated as for panel A. (D) Location of histone SDD residues on the nucleosome. The DNA helix is shown in purple. (Left) Front view of the nucleosome. (Right) The enlarged detail of the H3-H4 interface that makes contact with DNA is presented in the same orientation. Side chains of H3 R69, H3 E73, H3 Q76, H3 R83, and H4 N25 are shown in stick models. Dashed lines indicate hydrogen bonds. The drawings were produced using PyMol (<https://www.pymol.org/>) based on the atomic model of yeast nucleosome core particle (Protein Data Bank code 1ID3).

H3^{E73A}, *H3^{Q76E}* and *H3^{R83A}*, suppress the MMS sensitivity of *pol30^{K164R}* cells in an HR-dependent manner (data not shown). These results imply that the histone suppressor alleles exert their suppressive effects via a similar mechanism of action.

We next examined whether PCNA SUMOylation affects the suppression of MMS sensitivity of *pol30^{K164R}* cells by *H3^{R69A}*. The DDT-deficient *pol30^{K127R/K164R}* mutant was sensitive to MMS but to a lesser extent than *pol30^{K164R}* cells (Fig. 2C), because it lacks both SUMOylation sites. Interestingly, the *H3^{R69A}* mutation still suppressed the MMS

FIG 2 Histone SDD mutations suppress the DNA damage sensitivity of DDT-deficient cells. (A) Serial dilutions (10-fold) of equal numbers of cells were spotted onto YPDA plates with or without the indicated concentrations of MMS. The plates were incubated at 30°C for 3 days. For CLUV, the plates were incubated in the presence or absence of CLUV for 3 days at 30°C. (B) MMS-induced Rad53 and histone H2A phosphorylation. Cells were treated with 0.2% MMS for 2 h. Protein extracts were prepared and separated by SDS-PAGE. Western blotting was done with antibodies against Rad53, H2A-S129 phosphorylation, H3, and tubulin. (C to E) Histone SDD and PCNA-SUMO-Srs2 have different regulatory functions for the DDT pathway. The spot assays were done as described for panel A.

sensitivity of *pol30^{K127R/K164R}* cells (Fig. 2C). In addition, *H3^{R69A} srs2Δ* cells were more sensitive than *srs2Δ* cells (Fig. 2D), suggesting that the histone SDD has Srs2-independent functions. We note that the histone SDD mutations suppress the MMS sensitivity of *rad18Δ* cells less efficiently than they suppress that of *pol30^{K164R}* cells. This may be due to the various degrees to which the salvage HR events are inhibited by the PCNA-SUMO-Srs2, because the SUMO modification of PCNA at K164 still occurs in *rad18Δ* cells. This hypothesis was further supported by the observation that the *H3^{R69A}* mutation substantially suppressed the MMS sensitivity of *rad18Δ siz1Δ* cells that were defective for both ubiquitin and SUMO modifications of PCNA (Fig. 2E). Collectively, these results suggest that *H3^{R69A}*-dependent suppression of DDT deficiency is, at least partly, distinct from that caused by defects in the PCNA-SUMO-Srs2.

FIG 3 DNA damage-induced recombination and mutation frequencies in the wild-type and *H3^{R69A}* strains. (A) Recombination frequency in the wild-type and *H3^{R69A}* strains. Cells grown to early log phase at 30°C were treated with 0.01% MMS for 5 h. Intrachromosomal recombination frequencies between direct repeat sequences in *lys2BA-URA3-lys2BG* of the haploid strains were determined (see Materials and Methods). Error bars indicate the standard errors of the three independent experiments. (B) Mutation frequency in the wild-type and *H3^{R69A}* strains. Cells grown to early log phase at 30°C were treated with 0.01% MMS for 5 h. Canavanine-resistant mutants were selected on synthetic complete medium lacking arginine and containing canavanine. Error bars indicate the standard errors of the three independent experiments. The significance of the differences between the wild-type and *H3^{R69A}* strains was evaluated with an unpaired *t* test (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Histone SDD mutations affect recombination and mutation frequencies. Our results demonstrated that although histone SDD mutations do not result in an MMS-sensitive phenotype, combining a histone SDD mutation with a defect in DDT and HR pathways results in a synergistically enhanced phenotype, suggesting that the histone SDD patch is involved in the regulation of the DDT/HR pathways under MMS-induced replication stress. We therefore assessed whether histone SDD single mutation affects recombination and mutation frequencies because the DDT/HR pathway choice is tightly controlled to prevent genomic instability. We first measured the frequency of intrachromosomal mitotic recombination at the *lys2BA::URA3::lys2BG* allele on chromosome II in the wild-type and *H3^{R69A}* strains. In this assay, gene conversion events between differently mutated *lys2* alleles generate Lys⁺ Ura⁺ cells, whereas most of Lys⁺ Ura⁻ cells can arise from pop-out recombination by single-strand annealing events. We found no substantial effect of *H3^{R69A}* on the spontaneous frequency of gene conversion and pop-out recombination. In contrast, when cells were treated with MMS, *H3^{R69A}* cells had increased frequencies of both recombination events compared with the wild-type strains (Fig. 3A).

We next examined the frequency of forward mutations of the *CAN1* gene, which result in resistance to canavanine (Can^r). In the absence of MMS, Can^r mutation frequencies were similar in both the wild-type and *H3^{R69A}* strains (Fig. 3B). Notably, when cells were treated with MMS, Can^r mutation frequency was increased in both strains, but its level was lower in the *H3^{R69A}* mutant than in the wild-type strain (Fig. 3B). Taken together, these data suggest that histone SDD is involved in both DDT-dependent mutagenesis and suppression of HR in response to DNA damage and thus support the idea that histone SDD contributes to the regulation of DDT and HR during DNA damage-induced replication stress.

The histone SDD promotes DNA damage-induced ubiquitination of PCNA. Our data imply that the histone SDD mutations might affect posttranslational modifications of PCNA. To evaluate the presence of modified PCNA species, we used a strain expressing a c-Myc-tagged *POL30* (PCNA) gene that was integrated at the *POL30* genomic locus. Western blot analysis confirmed previously reported ubiquitin and SUMO modifications of PCNA at K164 following UV irradiation (9); slow-migrating bands, representing ubiquitination and SUMOylation of PCNA at K164, were abolished in the *pol30^{K164R}* mutant (Fig. 4A). PCNA ubiquitination and SUMOylation at K164 were also absent in

FIG 4 Histone SDD promotes DNA damage-induced PCNA ubiquitination. (A) UV-induced modifications of PCNA. Asynchronous cell cultures were exposed to UV light at 100 J/m² and incubated at 30°C for 2 h. Protein extracts were prepared, and PCNA-Myc was detected by Western blotting with anti-c-Myc. Ubiquitylated and SUMOylated species are indicated. (B and C) MMS-induced modifications of PCNA. Cultures of wild-type or *H3^{R69A}* *POL30*-Myc cells grown to early log phase at 30°C were arrested in G₁ with α -factor for 2 h. MMS (0.03%) was added to cell cultures for 20 min before release from G₁ arrest. Cell cultures were then washed and released into S phase in the presence of MMS (0.03%). Samples from synchronized cell cultures were taken at the indicated time points after release from G₁ block. Cell cycle progression was determined by flow cytometry (B), and PCNA-Myc was detected by Western blotting with anti-c-Myc antibody (C). The slower-migrating bands are indicated by arrowheads. An image obtained at low exposure is shown at the bottom. The intensities of PCNA-Ub relative to those of total PCNA were quantified, and the results were normalized to the signal at time zero (right). Error bars indicate standard deviations from three independent experiments. The significance of the differences between the wild-type and *H3^{R69A}* strains was evaluated with an unpaired *t* test (*, *P* < 0.05). (D) UV-induced modifications of PCNA. Log-phase cultures of wild-type or *H3^{R69A}* *POL30*-Myc cells were exposed to UV light at 100 J/m² and incubated at 30°C. Samples were taken at the indicated times and analyzed by Western blotting with anti-c-Myc antibody. An image obtained at low exposure is shown at the bottom. Quantification of band intensities was done as for panel C. Values represent the means from two independent experiments (right).

FIG 5 INO80 chromatin remodeler is necessary for replication fork recovery in *H3^{R69A}* cells. (A) Genetic interactions between *H3^{R69A}* and chromatin remodelers. Equal numbers of cells were serially diluted (10-fold per dilution) and spotted onto YPDA plates containing the indicated concentrations of MMS. The plates were incubated at 30°C for 3 days. (B) Delayed S-phase progression in *H3^{R69A} arp8Δ* cells. Cells were arrested in G₁ with α -factor for 2 h and were then released into YPDA medium. Samples were taken at the indicated time points after release from G₁ block, and cell cycle progression was determined by flow cytometry. (C) G₁-synchronized cells were released into YPDA medium containing 0.03% MMS. Samples were analyzed by flow cytometry. (D) G₁-synchronized cells were released into YPDA medium containing 0.1% MMS for 1 h. After MMS treatment, cells were washed with sodium thiosulfate and released into YPDA medium. Samples were taken at the indicated time points after release from MMS and were analyzed by flow cytometry. (E) Deletion of *SRS2*, but not *H3^{R69A}*, suppresses the MMS sensitivity of *pol30^{K164R} arp8Δ* cells. The spot assays were done as described for panel A. The plates were incubated at 30°C for 3 days.

the *rad18Δ* and the *siz1Δ* strains, respectively (Fig. 4A). To monitor the status of PCNA ubiquitination, synchronized cultures were treated with MMS for 20 min before release from G₁ arrest, and were then released into S phase in the presence of MMS. Fluorescence-activated cell sorting (FACS) analysis revealed that *H3^{R69A}* cells slowed their progression through S phase, similarly to the wild-type cells (Fig. 4B). While treatment of cells with MMS induces ubiquitin modifications of PCNA in both wild-type and *H3^{R69A}* cells, the level of PCNA ubiquitination was reduced in *H3^{R69A}* cells in comparison with that in wild-type cells (Fig. 4C). A similar reduction in PCNA ubiquitination was also found post-UV irradiation (100 J/m²) (Fig. 4D). We note that *H3^{R69A}* reduced overall levels of ubiquitinated PCNA upon MMS or UV exposure, suggesting that histone SDD affects not only the regulation of the TLS pathway but also that of Rad5-Mms2-Ubc13 error-free DDT. This is consistent with the observation that *H3^{R69A}* fully suppressed the CLUV sensitivity of *pol30^{K164R}* cells (Fig. 2A), because the TLS-deficient strain (*rad30Δ rev3Δ rev1Δ* mutant) did not show a detectable effect on the CLUV sensitivity, whereas the *rad5Δ* mutant had CLUV hypersensitivity equivalent to that of the *pol30^{K164R}* mutant (52). Taken together, these results suggest that the histone SDD region is involved in the regulation of DDT pathway at the level of PCNA ubiquitin modifications.

INO80 facilitates resumption of the cell cycle in *H3^{R69A}* cells after release from MMS-induced replication arrest. Chromatin modifications caused by histone modifying or ATP-dependent nucleosome remodeling factors play important roles in the cellular response to replication stress (31, 38, 53). Since modifications of histone SDD residues have not been reported to date, our investigation focused primarily on a subset of ATP-dependent nucleosome remodelers. INO80, RSC, and Fun30 have previously been implicated in the repair of DNA double-strand breaks (DSBs) and/or DNA damage tolerance (31, 32, 39, 41, 43, 44, 54, 55). Arp8, an actin-related protein, is a member of the Ino80 complex and is required for Ino80 ATPase and chromatin remodeling activities (56). Two closely related RSC complexes that function in the DNA damage response contain either Rsc1 or Rsc2 in a mutually exclusive manner (57). To test whether these nucleosome remodelers contribute to misregulation of DDT function in *H3^{R69A}* cells, we constructed *H3^{R69A} arp8Δ*, *H3^{R69A} rsc1Δ*, *H3^{R69A} rsc2Δ*, and *H3^{R69A} fun30Δ* mutants and tested their growth and MMS sensitivities. We found that *H3^{R69A} arp8Δ* cells grew very poorly even in the absence of exogenous DNA damage (Fig. 5A), whereas none of the other mutants exhibited a delay in growth rate. FACS analysis of yeast synchronized in the cell cycle revealed that the majority of wild-type, *H3^{R69A}*, and *arp8Δ* cells achieved 2C DNA content within 60 to 75 min after release from G₁ arrest (Fig. 5B). In contrast, *H3^{R69A} arp8Δ* cells were unable to complete DNA replication even after 120 min (Fig. 5B), indicating that S phase progression was compromised. In addition, the *H3^{R69A} arp8Δ* mutants were more sensitive to MMS than each single mutant, whereas *H3^{R69A}* did not affect the MMS sensitivity of other nucleosome remodeler-deficient mutants (Fig. 5A). These results demonstrate that INO80 plays a critical role in DNA replication in *H3^{R69A}* cells.

We next tested whether the INO80 complex affects S phase progression in *H3^{R69A}* cells treated with MMS. Cells were arrested in G₁ by α -factor and then released into medium containing a low concentration of MMS (0.03%). *H3^{R69A}* and *arp8Δ* cells showed a slight delay in completing S phase compared with the wild type (Fig. 5C). In contrast, the majority of *H3^{R69A} arp8Δ* cells remained in G₁/S phase throughout the experiments (Fig. 5C), indicating a severe defect in S phase progression. One possibility is that *H3^{R69A} arp8Δ* cells have a defect in the replication fork recovery process. To test this, G₁ synchronized cells were treated with MMS (0.1%) for 1 h and then released into MMS-free medium for recovery. Wild-type and *H3^{R69A}* cells progressed through S phase with roughly equal kinetics, as they resumed the cell cycle and completed S phase 3 h after MMS removal (Fig. 5D). As reported previously (39), *arp8Δ* cells did not complete S phase at the same time point (3 h) during replication recovery (Fig. 5D). Strikingly, the majority of *H3^{R69A} arp8Δ* cells arrested at G₁/S phase even 3 h after MMS removal (Fig. 5D). These results further emphasize the important role of INO80 complex in replication fork recovery in *H3^{R69A}* cells.

***H3^{R69A}* suppresses the MMS sensitivity of *pol30^{K164R}* cells in an Ino80-dependent manner.** To investigate whether INO80 function is linked to the suppression of DDT deficiency in the histone SDD mutant background, we examined DNA damage sensitivity in the *arp8Δ* strain background. We found that *pol30^{K164R} arp8Δ* cells are more sensitive to MMS than either of the single mutants, indicating that Arp8 has functions independent of the DDT pathway in the DNA damage response (Fig. 5E). Remarkably, the *H3^{R69A} pol30^{K164R} arp8Δ* triple mutants are as sensitive as *pol30^{K164R} arp8Δ* cells to MMS (Fig. 5E, top), suggesting that *H3^{R69A}*-dependent suppression of the DDT deficiency requires INO80 function. In addition, the suppression effect by *H3^{R69A}* is likely not due to an inactivation of SUMO-PCNA-associated Srs2 functions, as deletion of *SRS2* still suppressed the *pol30^{K164R} arp8Δ* sensitivity to MMS (Fig. 5E, bottom).

***H3^{R69A} arp8Δ* cells have a defect in HR-dependent recovery from MMS-induced replication arrest.** Our data imply that histone SDD has a regulatory role in the INO80-dependent processing of replication-associated lesions. To test this possibility, we constructed *H3^{R69A} rad51Δ*, *arp8Δ rad51Δ*, and *H3^{R69A} arp8Δ rad51Δ* cells and examined their growth and MMS sensitivity. We found that the *rad51Δ* mutation partially suppressed the growth defect of *H3^{R69A} arp8Δ* cells under normal growth conditions (Fig. 6A, left). Indeed, when G₁-synchronized *H3^{R69A} arp8Δ* and *H3^{R69A} arp8Δ rad51Δ* cells were released into fresh media to stimulate S phase progression, 80% of *H3^{R69A} arp8Δ rad51Δ* cells completed DNA replication in 90 min, whereas only 50% of *H3^{R69A} arp8Δ* cells did even at 120 min (Fig. 6A, right). These findings indicate that the lack of HR function in *H3^{R69A} arp8Δ* cells alleviates their growth defect. Thus, it is likely that the inhibition of unperturbed S phase progression of *H3^{R69A} arp8Δ* cells is caused in part by toxic recombination events, which is consistent with the hypothesis that the histone SDD patch regulates HR through INO80 functions.

As for MMS treatment, we found that the *H3^{R69A}* mutation rendered *rad51Δ* cells more sensitive to MMS (Fig. 6B), which is in agreement with the role of the histone SDD in the DDT pathway. Moreover, while the *rad51Δ* mutation partially suppressed the growth defect of *H3^{R69A} arp8Δ* cells, *H3^{R69A} arp8Δ rad51Δ* triple mutants were more sensitive to MMS than *arp8Δ rad51Δ* cells (Fig. 6B). Since the growth rates of both strains are somewhat different in the absence of exogenous DNA damage, we quantified the ability of these strains to form colonies in the presence of 0.05% MMS. Under these conditions, viability of *H3^{R69A} arp8Δ rad51Δ* cells (0.29%) was at least 20-fold lower than that of *arp8Δ rad51Δ* cells (6.9%). Taken together, these results suggest that the increased MMS sensitivity of *arp8Δ rad51Δ* cells conferred by *H3^{R69A}* is likely due to the channeling of replication-blocking lesions to the HR pathway.

To further examine whether HR is impeded in *H3^{R69A} arp8Δ* cells, we observed Rad52-yellow fluorescent protein (Rad52 YFP) focus formation after transient exposure to MMS, because Rad52 nuclear focus formation is a cellular indicator of HR (58). Wild-type and *H3^{R69A}* cells exhibited increased Rad52-YFP foci 2 h after release from MMS exposure, and the number of cells with foci dropped to about 10% within 6 h (Fig. 6C and D). Consistent with a previous study (39), *arp8Δ* cells showed a delayed disappearance of MMS-induced Rad52-YFP foci (Fig. 6D). Remarkably, *H3^{R69A} arp8Δ* cells had an increased number of Rad52-YFP foci during recovery, and Rad52-YFP foci persisted at a later time point (10 h) (Fig. 6C and D), indicating an impaired HR function during recovery in *H3^{R69A} arp8Δ* cells. Taken together, these results suggest that INO80-dependent HR might serve as an alternative mechanism that compensates for the impaired DDT pathway in *H3^{R69A}* cells.

DISCUSSION

Using a comprehensive histone H3 and H4 mutant library, we have identified many histone alleles that enhance the MMS/HU sensitivity of *rad18Δ* cells. Among 49 H3/H4 mutations identified in this screen, 31 alleles have not previously been linked with the DNA replication stress response. Thus, our strategy proved to be effective for the identification of histone mutations that affect the DNA damage and replication stress

FIG 6 Genetic and functional interactions between $H3^{R69A}$, $arp8\Delta$, and $rad51\Delta$ mutants. (A) The relevant strains were streaked onto YPDA plates and incubated at 30°C for 3 days (left). Cells were arrested in G₁ with α -factor for 2 h and were then released into YPDA medium. Samples were taken at the indicated time points, and DNA content was determined by flow cytometry (right). (B) MMS sensitivity of the $H3^{R69A} arp8\Delta rad51\Delta$ triple mutant. Serial dilutions (10-fold) of equal numbers of cells were spotted onto YPDA plates with or without MMS. The plates were incubated at 30°C for 3 days. (C) MMS-induced Rad52-YFP foci in wild-type, $H3^{R69A}$, $arp8\Delta$, and $H3^{R69A} arp8\Delta$ (Continued on next page)

FIG 7 A model for the functional interaction of the histone SDD patch and INO80 in the control of DDT and HR pathways. The histone SDD patch might affect the chromatin remodeling activity of INO80 following replication stress, which is responsible for DDT/HR output when replication is impeded. (Left) In the wild-type background, the association of INO80 with the histone SDD patch and/or the histone SDD patch itself would promote replicative bypass of lesions by the DDT pathway. (Right) Histone SDD mutations would influence the crucial interactions between INO80 and the nucleosome in order to regulate the DDT, salvage HR, or both pathways, thereby channeling the cellular mechanism dealing with replication-blocking lesions into the salvage HR pathway (see the text for more details).

response. Further studies will be needed to determine how these residues contribute to these processes.

In addition to MMS/HU sensitive mutants, we identified six H3/H4 alleles (histone SDD mutations; *H3^{R69A}*, *H3^{E73A}*, *H3^{E73Q}*, *H3^{Q76E}*, *H3^{R83A}*, and *H4^{N25D}*) that suppress the MMS, UV, and/or HU sensitivity of DDT-deficient cells. To our knowledge, despite the large number of studies conducted on histone mutations that reported DNA damage responses, limited information is currently available on the functions of histone SDD residues, except for H3 R83; *H3^{R83A}* mutation causes defects in ribosomal DNA (rDNA) and telomere silencing (59). However, in contrast to other histone SDD mutations, *H3^{R83A}* cells are slightly more sensitive to MMS (Fig. 1C), implying that it is involved in DNA damage responses that are distinct from those of the SDD region. Thus, the histone SDD region has not previously been linked with the DNA damage response. One explanation for this is that histone SDD mutations confer resistance to genotoxic drugs only in DDT-deficient cells and therefore are difficult to identify via screening for the histone single mutants. Furthermore, our study demonstrates that Rad51 is critical for the suppression of DDT defects engendered by histone SDD mutations; this is consistent with the observation that DNA damage-induced recombination frequency increased in *H3^{R69A}* cells. These results suggest that the suppression of DDT deficiency in histone SDD mutants might be due to an exacerbated HR function under replication stress conditions. Notably, this effect appears to be unrelated to an impaired PCNA-SUMO-Srs2 function but instead requires INO80 to mediate HR-dependent recovery from replication stress in *H3^{R69A}* cells (Fig. 7). Thus, histone SDD and Srs2 have distinct regulatory functions for the DDT pathway and likely cooperate to determine DDT output when DNA replication is impeded. In this context, the levels of PCNA ubiquitination upon DNA damage are substantially reduced in histone SDD mutants compared with those in wild-type cells. This is consistent with a decreased mutation frequency in *H3^{R69A}* cells. We speculate that the histone SDD patch positively regulates the DDT pathway, although the detailed mechanism remains unclear (Fig. 7). In addition to its role in DDT regulation, an alternative, but not mutually exclusive, possibility is that the

FIG 6 Legend (Continued)

arp8Δ cells. Cells grown to early log phase at 30°C were treated with 0.1% MMS for 20 min and released into YPDA medium. At the indicated time points, samples were taken for fluorescence microscopy. Representative images of Rad52-YFP foci are shown. (D) Quantitation of cells with Rad52-YFP foci. Error bars indicate the standard errors of the three independent experiments.

enhanced salvage HR pathway in histone SDD mutants indirectly affects the DDT function (Fig. 7), as shown previously (24). Taken together, our findings demonstrate novel regulatory mechanisms within the replication stress response and highlight the role of chromatin structure as an important modulator of DDT and HR functions.

INO80 is highly conserved in eukaryotes, including yeast and mammals, and is implicated in diverse DNA transactions such as transcription, DNA replication, homologous recombination, and DNA repair (31, 38, 53, 60, 61). The INO80 complex is recruited to sites of stalled replication forks, stabilizes stalled forks, and also regulates DDT and HR during the recovery from MMS- and HU-induced replication stress (39, 41). These results suggest that the ability of INO80 to remove/translocate nucleosomes at the stalled fork facilitates the recruitment of essential factors for repair and recovery. In the present study, we showed that introduction of *H3^{R69A}* onto the *arp8Δ* background causes a modest growth defect under unperturbed conditions and results in poor viability, defects in replication completion, and accumulation of Rad52 foci after treatment with MMS. We also demonstrated that *H3^{R69A}* suppresses the MMS sensitivity of DDT-deficient cells in an Ino80-dependent manner. These results suggest that the histone SDD mutations enhance the salvage HR pathway in recovery from DNA damage-induced replication arrest, likely through modulating the activity and/or chromatin accessibility of INO80 chromatin remodeler, which fits well with the known roles of yeast and mammalian INO80 in promoting HR (43, 44, 46, 60, 61). Although precisely how histone SDD affects INO80 activity is unclear, one possibility is that the histone SDD mutations affect the histone-DNA contact required for the accessibility of enzymes that mediate the DDT pathway, thereby creating a dependence on INO80 chromatin remodeling activity in order to overcome such nucleosome impediments at stalled forks. However, we could not demonstrate that histone SDD mutation promotes the enrichment of INO80 complex upon chromatin DNA following DNA damage (our unpublished data). Alternatively, the histone SDD region might directly affect the chromatin remodeling activity of INO80 following replication stress. Previous structural and functional studies in yeast have reported that INO80 Arp module initially binds to the extranucleosomal DNA, and prominent conformational changes in the INO80 complex promote nucleosome contacts, which drive remodeling reactions by INO80 via coupling ATPase activity to nucleosome repositioning (62, 63). Thus, the role of the Arp8 module binding to DNA is not simply to recruit INO80 to nucleosomes but rather to promote allosteric changes in INO80 required to efficiently remodel nucleosomes. In this context, the INO80 complex subunit Arp8 directly binds to nucleosomes (56, 64), and especially, its C-terminal domain interacts with the histone H3-H4 interface containing H3 Leu71 and H4 Leu23 residues (65), which is very close to or overlaps with the SDD patch. Therefore, our current hypothesis is that SDD mutations may influence the balance of the regulation of DDT and HR in a manner that modulates crucial interactions between INO80 and nucleosomal substrates in order to regulate nucleosome translocation, thus channeling the cellular mechanism dealing with replication-blocking lesions into HR (Fig. 7). Further biochemical studies will be needed to elucidate how the histone SDD region affects INO80 chromatin remodeling activity during DNA damage-induced replication stress. The results of such studies should advance our understanding of the molecular mechanisms by which chromatin remodeling enzymes and chromatin topology regulate DDT and HR pathways during the replication stress response.

MATERIALS AND METHODS

Yeast strains and media. All yeast strains used in this study are derivatives of BY4741 (Open Biosystems) and are listed in Table S1 in the supplemental material. The yeast synthetic histone H3 and H4 mutant collection was purchased from Dharmacon (now Horizon). It consists of 345 histone H3 and H4 alleles, where every H3 and H4 amino acid was systematically changed to alanine or serine, and the modifiable residues were mutated to mimic constitutively posttranslational modification states (for example, lysine residues were mutated to glutamine to mimic the acetylation state) (49). Standard genetic procedures were used for strain construction and medium preparation (66). Strains carrying a C-terminal c-Myc epitope tag or YFP were made by transforming the cells with PCR-amplified cassettes

that carry homologies to the C terminus of *POL30* as explained previously (67, 68). Yeast cells were routinely grown in YPD medium (1% yeast extract, 2% peptone, 2% dextrose) containing 0.003% adenine sulfate (YPDA). Synthetic complete (SC) medium was used in strain selection and recombination and mutation frequency analysis. For the G₁ synchronization of cells, the cultures were treated for 2 h with 10 μ g/ml of α -factor (Sigma-Aldrich).

Construction of a histone H3/H4 mutation library in DDT-deficient *rad18* Δ cells. The histone and *rad18* Δ double mutant collection was constructed by systematic genetic array (SGA) analysis (48). Briefly, the MH101 haploid strain (BY4741 *rad18* Δ ::*LEU2 MAT α*) was mated to each *MAT α* haploid library of histone H3 or H4 single mutants and mated cells were incubated on SC-Ura-Leu plates at 30°C, producing a collection of doubly heterozygous diploids. Following sporulation, the set of desired haploid double mutants was subsequently selected according to selection markers (HIS, URA, LEU, and NAT). Of 345 histone mutants, 328 *rad18* Δ H3/H4^m double mutants were successfully constructed using this approach. For reasons unknown, we were unable to obtain five *rad18* Δ H3/H4^m double mutants (H3^m [R49A, R63A, and Q93E]/H4^m [S69A and Y98A]), and therefore, they were constructed by introducing a *rad18* Δ ::*LEU2* construct to each histone mutants. The remaining 12 mutants (H3^m [R17K, K42A, G44A, R53K, K56A, T58D, L61A, and Q93E]/H4^m [G28A, Y51A, K91A, and Y98A]) were not analyzed further in this study because they also contained additional unexpected mutations and/or did not contain the desired mutations following sequencing during screening. We note that all double mutants are viable, but *rad18* Δ cells with one of eight histone mutations (H3^m [R49A, R63A, S95A, and L109A]/H4^m [R36A, R36K, I46A, and E53A]) show a slow growth phenotype compared with *rad18* Δ single mutants. For DNA damage sensitivity screening, 10-fold serial dilutions of strains were spotted onto YPDA plates with or without MMS (0.0003%) or HU (25 mM). The plates were incubated for 3 days at 30°C. We then evaluated relative colony-forming ability of these strains by comparison to those of the *rad18* Δ strains expressing one copy of wild-type histones H3 and H4, according to the previous study (49, 50) (<http://www.histonehits.org>).

Sensitivity to MMS, HU, and CLUV. Asynchronous cell cultures grown in liquid YPDA medium were normalized to 2×10^8 cells/ml at appropriate dilutions, and 10-fold serial dilutions starting with equal numbers of cells were performed in 96-well plates. Aliquots (3 μ l) of each dilution were spotted onto YPDA plates containing MMS (Sigma-Aldrich) or HU (Sigma-Aldrich). The plates were then incubated for 3 days at 30°C. For CLUV sensitivity assays, the plates were incubated for 3 days at 30°C under continuous exposure to UV irradiation at a dose of ≈ 0.1 J/m²/min as described in previous studies (52, 69).

Recombination and mutation frequency. Intrachromosomal mitotic recombination was assessed by using heteroalleles of *lys2* in a tandem duplication, *lys2BA::URA3::lys2BG* on chromosome II (70). Lys⁺ Ura⁺ recombinants can occur by gene conversion without crossing over, whereas Lys⁺ Ura⁻ recombinants can occur by single-strand annealing or by gene conversion with crossing over, but the latter event is rare. Recombination frequencies were determined as described previously (70). Briefly, strains were streaked out on YPDA plates and grown at 30°C. Five colonies for each strain were suspended in YPDA medium and grown to early logarithmic phase. The cell cultures were then treated or not treated with 0.01% MMS for 5 h (MMS+ or MMS-). Subsequently, cells were washed with 5% sodium thiosulfate to inactivate the drug and plated onto plates at an appropriate dilution to determine the total cell number on YPDA plates and the number of recombinant mutants on SC plates without Lys and/or Ura. Plates were incubated at 30°C for 3 days. The recombination frequency was determined by dividing the total number of recombinants on SC plates without Lys and/or Ura by the total number of viable cells. The Lys⁺ Ura⁻ frequency was calculated by subtracting the Lys⁺ Ura⁺ frequency from the Lys⁺ frequency. The average was determined for at least three independent sets of experiments.

Mutation frequencies at *CAN1* locus were determined as described previously (10, 69). Five independent colonies of a strain were used to measure mutation frequencies. Briefly, colonies of cells were grown to early logarithmic phase in YPDA medium. The cell cultures were then treated or not with 0.01% MMS for 5 h. Aliquots were removed from the MMS- and MMS+ cultures, washed with 5% sodium thiosulfate, and plated onto YPDA medium at an appropriate dilution and SC-arginine plates containing 60 μ g/ml of canavanine medium. Plates were incubated at 30°C for 3 days. The mutation frequency was determined by dividing the number of Can^r colonies by the total number of viable cells, and the average was determined for at least three independent sets of experiments. A *t* test was used to determine the significance of differences between wild-type and H3^{R69A} strains in recombination and mutation frequencies.

Yeast cell extract and Western blotting. Yeast cell extracts were prepared from yeast cultures using the trichloroacetic acid method, as described previously (26). For Western blotting of PCNA, yeast cell extracts were prepared from yeast cultures using Na₂S₂O₃ methods, as described previously (66). Protein samples were separated using SDS-polyacrylamide gels, and proteins were transferred to polyvinylidene difluoride (PVDF) membranes. PCNA-Myc, Rad53, H2A, H3, and tubulin were detected with anti-c-Myc (MBL), anti-Rad53 (Santa Cruz), anti- γ -H2A (phospho S129) (Millipore), anti-H3 (Abcam), and anti- α -tubulin (Sigma) antibodies, respectively. The protein bands were visualized by enhanced chemiluminescence detection (Western blot ultrasensitive horseradish peroxidase [HRP] substrate; TaKaRa). The intensity of each protein band was quantified using ImageJ software.

Fluorescence microscopy and analysis of Rad52-YFP localization. Cells grown to early log phase at 30°C were treated with 0.1% MMS for 20 min. Subsequently, cells were washed with 5% sodium thiosulfate and released into YPDA. Samples were collected at various time points and resuspended in phosphate-buffered saline (PBS). Fluorescence microscopy was performed on a Zeiss Axioplan2. Images were visualized using the Lumina Vision imaging software program (Mitani Corporation). More than 100 individual cells were scored for each strain. Data were obtained for at least three independent sets of experiments.

Cell cycle arrest and flow cytometry. Cells were grown to early log phase at 30°C in YPDA medium and treated for 2 h with 10 μ g/ml of α -factor (Sigma). To release cells from α -factor arrest, cells were washed twice with water. Samples were collected at various times for fluorescence-activated cell sorting (FACS) analysis and protein extraction. FACS analysis was performed as described previously (71). Briefly, cells were fixed in 70% ethanol, washed, and resuspended in 50 mM sodium citrate (pH 7.5). Samples were treated with RNase A (0.25 mg/ml) for 1 h at 50°C and then with proteinase K (1 mg/ml) for 1 h at 50°C. Cells were stained with propidium iodide (16 μ g/ml) for 1 h at 4°C. The DNA contents of cells were analyzed with a Becton, Dickinson, FACSCanto II flow cytometer.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 0.1 MB.

ACKNOWLEDGMENTS

We thank Tomomi Ohtawara for technical assistance and members of the Hishida laboratory for helpful discussions.

This work was supported by JSPS KAKENHI grant numbers JP17K07290 and JP23114007 (to T.H.) and JP20K05790 (to G.A.).

REFERENCES

- Zeman MK, Cimprich KA. 2014. Causes and consequences of replication stress. *Nat Cell Biol* 16:2–9. <https://doi.org/10.1038/ncb2897>.
- Halazonetis TD, Gorgoulis VG, Bartek J. 2008. An oncogene-induced DNA damage model for cancer development. *Science* 319:1352–1355. <https://doi.org/10.1126/science.1140735>.
- Negrini S, Gorgoulis VG, Halazonetis TD. 2010. Genomic instability—an evolving hallmark of cancer. *Nat Rev Mol Cell Biol* 11:220–228. <https://doi.org/10.1038/nrm2858>.
- Branzei D, Foiani M. 2010. Maintaining genome stability at the replication fork. *Nat Rev Mol Cell Biol* 11:208–219. <https://doi.org/10.1038/nrm2852>.
- Ulrich HD. 2011. Timing and spacing of ubiquitin-dependent DNA damage bypass. *FEBS Lett* 585:2861–2867. <https://doi.org/10.1016/j.febslet.2011.05.028>.
- Saugar I, Ortiz-Bazan MA, Tercero JA. 2014. Tolerating DNA damage during eukaryotic chromosome replication. *Exp Cell Res* 329:170–177. <https://doi.org/10.1016/j.yexcr.2014.07.009>.
- Bonner JN, Zhao X. 2016. Replication-associated recombinational repair: lessons from budding yeast. *Genes (Basel)* 7:48. <https://doi.org/10.3390/genes7080048>.
- Wong RP, Garcia-Rodriguez N, Zilio N, Hanulova M, Ulrich HD. 2020. Processing of DNA polymerase-blocking lesions during genome replication is spatially and temporally segregated from replication forks. *Mol Cell* 77:3–16.e14. <https://doi.org/10.1016/j.molcel.2019.09.015>.
- Hoege C, Pfander B, Moldovan GL, Pyrowolakis G, Jentsch S. 2002. RAD6-dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature* 419:135–141. <https://doi.org/10.1038/nature00991>.
- Stelter P, Ulrich HD. 2003. Control of spontaneous and damage-induced mutagenesis by SUMO and ubiquitin conjugation. *Nature* 425:188–191. <https://doi.org/10.1038/nature01965>.
- Andersen PL, Xu F, Xiao W. 2008. Eukaryotic DNA damage tolerance and translesion synthesis through covalent modifications of PCNA. *Cell Res* 18:162–173. <https://doi.org/10.1038/cr.2007.114>.
- Waters LS, Minesinger BK, Wiltout ME, D'Souza S, Woodruff RV, Walker GC. 2009. Eukaryotic translesion polymerases and their roles and regulation in DNA damage tolerance. *Microbiol Mol Biol Rev* 73:134–154. <https://doi.org/10.1128/MMBR.00034-08>.
- Lawrence CW, Christensen RB. 1979. Ultraviolet-induced reversion of *cyc1* alleles in radiation-sensitive strains of yeast. III. *rev3* mutant strains. *Genetics* 92:397–408.
- Kozmin SG, Pavlov YI, Kunkel TA, Sage E. 2003. Roles of *Saccharomyces cerevisiae* DNA polymerases Poleta and Polzeta in response to irradiation by simulated sunlight. *Nucleic Acids Res* 31:4541–4552. <https://doi.org/10.1093/nar/gkg489>.
- Parker JL, Ulrich HD. 2009. Mechanistic analysis of PCNA poly-ubiquitylation by the ubiquitin protein ligases Rad18 and Rad5. *EMBO J* 28:3657–3666. <https://doi.org/10.1038/emboj.2009.303>.
- Unk I, Hajdu I, Blastyak A, Haracska L. 2010. Role of yeast Rad5 and its human orthologs, HLTf and SHPRH in DNA damage tolerance. *DNA Repair (Amst)* 9:257–267. <https://doi.org/10.1016/j.dnarep.2009.12.013>.
- Giannattasio M, Zwicky K, Follonier C, Foiani M, Lopes M, Branzei D. 2014. Visualization of recombination-mediated damage bypass by template switching. *Nat Struct Mol Biol* 21:884–892. <https://doi.org/10.1038/nsmb.2888>.
- Prado F. 2014. Homologous recombination maintenance of genome integrity during DNA damage tolerance. *Mol Cell Oncol* 1:e957039. <https://doi.org/10.4161/23723548.2014.957039>.
- Branzei D, Foiani M. 2008. Regulation of DNA repair throughout the cell cycle. *Nat Rev Mol Cell Biol* 9:297–308. <https://doi.org/10.1038/nrm2351>.
- Carr AM, Lambert S. 2013. Replication stress-induced genome instability: the dark side of replication maintenance by homologous recombination. *J Mol Biol* 425:4733–4744. <https://doi.org/10.1016/j.jmb.2013.04.023>.
- Heyer WD, Ehmsen KT, Liu J. 2010. Regulation of homologous recombination in eukaryotes. *Annu Rev Genet* 44:113–139. <https://doi.org/10.1146/annurev-genet-051710-150955>.
- Gonzalez-Prieto R, Munoz-Cabello AM, Cabello-Lobato MJ, Prado F. 2013. Rad51 replication fork recruitment is required for DNA damage tolerance. *EMBO J* 32:1307–1321. <https://doi.org/10.1038/emboj.2013.73>.
- Branzei D, Szakal B. 2016. DNA damage tolerance by recombination: molecular pathways and DNA structures. *DNA Repair (Amst)* 44:68–75. <https://doi.org/10.1016/j.dnarep.2016.05.008>.
- Pfander B, Moldovan GL, Sacher M, Hoege C, Jentsch S. 2005. SUMO-modified PCNA recruits Srs2 to prevent recombination during S phase. *Nature* 436:428–433. <https://doi.org/10.1038/nature03665>.
- Papouli E, Chen S, Davies AA, Huttner D, Krejci L, Sung P, Ulrich HD. 2005. Crosstalk between SUMO and ubiquitin on PCNA is mediated by recruitment of the helicase Srs2p. *Mol Cell* 19:123–133. <https://doi.org/10.1016/j.molcel.2005.06.001>.
- Hishida T, Ohya T, Kubota Y, Kamada Y, Shinagawa H. 2006. Functional and physical interaction of yeast Mgs1 with PCNA: impact on RAD6-dependent DNA damage tolerance. *Mol Cell Biol* 26:5509–5517. <https://doi.org/10.1128/MCB.00307-06>.
- Iraqi I, Chekkal Y, Jmari N, Pietrobon V, Freon K, Costes A, Lambert SA. 2012. Recovery of arrested replication forks by homologous recombination is error-prone. *PLoS Genet* 8:e1002976. <https://doi.org/10.1371/journal.pgen.1002976>.
- Keyamura K, Arai K, Hishida T. 2016. Srs2 and Mus81-Mms4 prevent accumulation of toxic inter-homolog recombination intermediates. *PLoS Genet* 12:e1006136. <https://doi.org/10.1371/journal.pgen.1006136>.
- Sinha M, Watanabe S, Johnson A, Moazed D, Peterson CL. 2009. Recombinational repair within heterochromatin requires ATP-dependent chromatin remodeling. *Cell* 138:1109–1121. <https://doi.org/10.1016/j.cell.2009.07.013>.
- House NC, Koch MR, Freudenreich CH. 2014. Chromatin modifications and DNA repair: beyond double-strand breaks. *Front Genet* 5:296. <https://doi.org/10.3389/fgene.2014.00296>.

31. Poli J, Gasser SM, Papamichos-Chronakis M. 2017. The INO80 remodeller in transcription, replication and repair. *Philos Trans R Soc Lond B Biol Sci* 372:20160290. <https://doi.org/10.1098/rstb.2016.0290>.
32. Narlikar GJ, Sundaramoorthy R, Owen-Hughes T. 2013. Mechanisms and functions of ATP-dependent chromatin-remodeling enzymes. *Cell* 154:490–503. <https://doi.org/10.1016/j.cell.2013.07.011>.
33. Gerhold CB, Gasser SM. 2014. INO80 and SWR complexes: relating structure to function in chromatin remodeling. *Trends Cell Biol* 24:619–631. <https://doi.org/10.1016/j.tcb.2014.06.004>.
34. Shen X, Mizuguchi G, Hamiche A, Wu C. 2000. A chromatin remodelling complex involved in transcription and DNA processing. *Nature* 406:541–544. <https://doi.org/10.1038/35020123>.
35. Jonsson ZO, Jha S, Wohlschlegel JA, Dutta A. 2004. Rvb1p/Rvb2p recruit Arp5p and assemble a functional Ino80 chromatin remodeling complex. *Mol Cell* 16:465–477. <https://doi.org/10.1016/j.molcel.2004.09.033>.
36. Tosi A, Haas C, Herzog F, Gilmozzi A, Berninghausen O, Ungewickell C, Gerhold CB, Lakomek K, Aebersold R, Beckmann R, Hopfner KP. 2013. Structure and subunit topology of the INO80 chromatin remodeler and its nucleosome complex. *Cell* 154:1207–1219. <https://doi.org/10.1016/j.cell.2013.08.016>.
37. Falbo KB, Shen X. 2012. Function of the INO80 chromatin remodeling complex in DNA replication. *Front Biosci (Landmark Ed)* 17:970–975. <https://doi.org/10.2741/3967>.
38. Morrison AJ. 2017. Genome maintenance functions of the INO80 chromatin remodeller. *Philos Trans R Soc Lond B Biol Sci* 372:20160289. <https://doi.org/10.1098/rstb.2016.0289>.
39. Shimada K, Oma Y, Schleker T, Kugou K, Ohta K, Harata M, Gasser SM. 2008. Ino80 chromatin remodeling complex promotes recovery of stalled replication forks. *Curr Biol* 18:566–575. <https://doi.org/10.1016/j.cub.2008.03.049>.
40. Papamichos-Chronakis M, Peterson CL. 2008. The Ino80 chromatin-remodeling enzyme regulates replisome function and stability. *Nat Struct Mol Biol* 15:338–345. <https://doi.org/10.1038/nsmb.1413>.
41. Falbo KB, Alabert C, Katou Y, Wu S, Han J, Wehr T, Xiao J, He X, Zhang Z, Shi Y, Shirahige K, Pasero P, Shen X. 2009. Involvement of a chromatin remodeling complex in damage tolerance during DNA replication. *Nat Struct Mol Biol* 16:1167–1172. <https://doi.org/10.1038/nsmb.1686>.
42. Vassileva I, Yanakieva I, Peycheva M, Gospodinov A, Anachkova B. 2014. The mammalian INO80 chromatin remodeling complex is required for replication stress recovery. *Nucleic Acids Res* 42:9074–9086. <https://doi.org/10.1093/nar/gku605>.
43. Morrison AJ, Highland J, Krogan NJ, Arbel-Eden A, Greenblatt JF, Haber JE, Shen X. 2004. INO80 and gamma-H2AX interaction links ATP-dependent chromatin remodeling to DNA damage repair. *Cell* 119:767–775. <https://doi.org/10.1016/j.cell.2004.11.037>.
44. van Attikum H, Fritsch O, Hohn B, Gasser SM. 2004. Recruitment of the INO80 complex by H2A phosphorylation links ATP-dependent chromatin remodeling with DNA double-strand break repair. *Cell* 119:777–788. <https://doi.org/10.1016/j.cell.2004.11.033>.
45. Tsukuda T, Fleming AJ, Nickoloff JA, Osley MA. 2005. Chromatin remodeling at a DNA double-strand break site in *Saccharomyces cerevisiae*. *Nature* 438:379–383. <https://doi.org/10.1038/nature04148>.
46. Gospodinov A, Vaissiere T, Krastev DB, Legube G, Anachkova B, Herceg Z. 2011. Mammalian Ino80 mediates double-strand break repair through its role in DNA end strand resection. *Mol Cell Biol* 31:4735–4745. <https://doi.org/10.1128/MCB.06182-11>.
47. Horigome C, Oma Y, Konishi T, Schmid R, Marcomini I, Hauer MH, Dion V, Harata M, Gasser SM. 2014. SWR1 and INO80 chromatin remodelers contribute to DNA double-strand break perinuclear anchorage site choice. *Mol Cell* 55:626–639. <https://doi.org/10.1016/j.molcel.2014.06.027>.
48. Tong AH, Evangelista M, Parsons AB, Xu H, Bader GD, Page N, Robinson M, Raghibizadeh S, Hogue CW, Bussey H, Andrews B, Tiers M, Boone C. 2001. Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science* 294:2364–2368. <https://doi.org/10.1126/science.1065810>.
49. Dai J, Hyland EM, Yuan DS, Huang H, Bader JS, Boeke JD. 2008. Probing nucleosome function: a highly versatile library of synthetic histone H3 and H4 mutants. *Cell* 134:1066–1078. <https://doi.org/10.1016/j.cell.2008.07.019>.
50. Huang H, Maertens AM, Hyland EM, Dai J, Norris A, Boeke JD, Bader JS. 2009. HistoneHits: a database for histone mutations and their phenotypes. *Genome Res* 19:674–681. <https://doi.org/10.1101/gr.083402.108>.
51. Hodges AJ, Gallegos IJ, Laughery MF, Meas R, Tran L, Wyrick JJ. 2015. Histone sprocket arginine residues are important for gene expression, DNA repair, and cell viability in *Saccharomyces cerevisiae*. *Genetics* 200:795–806. <https://doi.org/10.1534/genetics.115.175885>.
52. Hishida T, Kubota Y, Carr AM, Iwasaki H. 2009. RAD6-RAD18-RAD5-pathway-dependent tolerance to chronic low-dose ultraviolet light. *Nature* 457:612–615. <https://doi.org/10.1038/nature07580>.
53. Bartholomew B. 2014. Regulating the chromatin landscape: structural and mechanistic perspectives. *Annu Rev Biochem* 83:671–696. <https://doi.org/10.1146/annurev-biochem-051810-093157>.
54. Niimi A, Chambers AL, Downs JA, Lehmann AR. 2012. A role for chromatin remodellers in replication of damaged DNA. *Nucleic Acids Res* 40:7393–7403. <https://doi.org/10.1093/nar/gks453>.
55. Chen X, Cui D, Papusha A, Zhang X, Chu CD, Tang J, Chen K, Pan X, Ira G. 2012. The Fun30 nucleosome remodeller promotes resection of DNA double-strand break ends. *Nature* 489:576–580. <https://doi.org/10.1038/nature11355>.
56. Shen X, Ranallo R, Choi E, Wu C. 2003. Involvement of actin-related proteins in ATP-dependent chromatin remodeling. *Mol Cell* 12:147–155. [https://doi.org/10.1016/s1097-2765\(03\)00264-8](https://doi.org/10.1016/s1097-2765(03)00264-8).
57. Chambers AL, Brownlee PM, Durley SC, Beacham T, Kent NA, Downs JA. 2012. The two different isoforms of the RSC chromatin remodeling complex play distinct roles in DNA damage responses. *PLoS One* 7:e32016. <https://doi.org/10.1371/journal.pone.0032016>.
58. Alvaro D, Lisby M, Rothstein R. 2007. Genome-wide analysis of Rad52 foci reveals diverse mechanisms impacting recombination. *PLoS Genet* 3:e228. <https://doi.org/10.1371/journal.pgen.0030228>.
59. Park JH, Cosgrove MS, Youngman E, Wolberger C, Boeke JD. 2002. A core nucleosome surface crucial for transcriptional silencing. *Nat Genet* 32:273–279. <https://doi.org/10.1038/ng982>.
60. van Attikum H, Fritsch O, Gasser SM. 2007. Distinct roles for SWR1 and INO80 chromatin remodeling complexes at chromosomal double-strand breaks. *EMBO J* 26:4113–4125. <https://doi.org/10.1038/sj.emboj.7601835>.
61. Lademann CA, Renkawitz J, Pfander B, Jentsch S. 2017. The INO80 complex removes H2A.Z to promote presynaptic filament formation during homologous recombination. *Cell Rep* 19:1294–1303. <https://doi.org/10.1016/j.celrep.2017.04.051>.
62. Brahma S, Ngubo M, Paul S, Udagama M, Bartholomew B. 2018. The Arp8 and Arp4 module acts as a DNA sensor controlling INO80 chromatin remodeling. *Nat Commun* 9:3309. <https://doi.org/10.1038/s41467-018-05710-7>.
63. Knoll KR, Eustermann S, Niebauer V, Oberbeckmann E, Stoehr G, Schall K, Tosi A, Schwarz M, Buchfellner A, Korber P, Hopfner KP. 2018. The nuclear actin-containing Arp8 module is a linker DNA sensor driving INO80 chromatin remodeling. *Nat Struct Mol Biol* 25:823–832. <https://doi.org/10.1038/s41594-018-0115-8>.
64. Gerhold CB, Winkler DD, Lakomek K, Seifert FU, Fenn S, Kessler B, Witte G, Luger K, Hopfner KP. 2012. Structure of actin-related protein 8 and its contribution to nucleosome binding. *Nucleic Acids Res* 40:11036–11046. <https://doi.org/10.1093/nar/gks842>.
65. Saravanan M, Wuerges J, Bose D, McCormack EA, Cook NJ, Zhang X, Wigley DB. 2012. Interactions between the nucleosome histone core and Arp8 in the INO80 chromatin remodeling complex. *Proc Natl Acad Sci U S A* 109:20883–20888. <https://doi.org/10.1073/pnas.1214735109>.
66. Amberg DC, Burke DJ, Strathern JN. 2005. Methods in yeast genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
67. Longtine MS, McKenzie III A, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR. 1998. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14:953–961. [https://doi.org/10.1002/\(SICI\)1097-0061\(199807\)14:10<953::AID-YEA293>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U).
68. Reid RJ, Lisby M, Rothstein R. 2002. Cloning-free genome alterations in *Saccharomyces cerevisiae* using adaptamer-mediated PCR. *Methods Enzymol* 350:258–277. [https://doi.org/10.1016/s0076-6879\(02\)50968-x](https://doi.org/10.1016/s0076-6879(02)50968-x).
69. Haruta N, Kubota Y, Hishida T. 2012. Chronic low-dose ultraviolet-induced mutagenesis in nucleotide excision repair-deficient cells. *Nucleic Acids Res* 40:8406–8415. <https://doi.org/10.1093/nar/gks580>.
70. Hishida T, Ohno T, Iwasaki H, Shinagawa H. 2002. *Saccharomyces cerevisiae* MGS1 is essential in strains deficient in the RAD6-dependent DNA damage tolerance pathway. *EMBO J* 21:2019–2029. <https://doi.org/10.1093/emboj/21.8.2019>.
71. Hayashi M, Keyamura K, Hishida T. 2018. Cyclin-dependent kinase modulates budding yeast Rad5 stability during cell cycle. *PLoS One* 13:e0204680. <https://doi.org/10.1371/journal.pone.0204680>.