Structure and activity of SLAC1 channels for stomatal signaling in leaves

Ya-nan Deng^{a,b,c}, Hamdy Kashtoh^{a,b,1}, Quan Wang^{d,1}, Guang-xiao Zhen^{a,b,c}, Qi-yu Li^{a,b,c}, Ling-hui Tang^{a,b,c}, Hai-long Gao^d, Chun-rui Zhang^{a,b,c}, Li Qin^{a,b,c}, Min Su^{a,b}, Fei Li^e, Xia-he Huang^{a,b}, Ying-chun Wang^{a,b,c}, Qi Xie^{a,b,c}, Oliver B. Clarke^{f,g,2}, Wayne A. Hendrickson^{f,h,2}, and Yu-hang Chen^{a,b,c,i,2}

^aState Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, 100101 Beijing, China; ^bInnovative Academy of Seed Design, Chinese Academy of Sciences, 100101 Beijing, China; ^cCollege of Advanced Agricultural Sciences, University of Chinese Academy of Sciences, 100049 Beijing, China; ^dNational Laboratory of Biomacromolecules, CAS Center for Excellence in Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, 100101 Beijing, China; ^eDepartment of Biology, New York University, New York, NY 10003; ^fDepartment of Physiology and Cellular Biophysics, Columbia University, New York, NY 10032; ^gDepartment of Anesthesiology, Columbia University, New York, NY 10032; ^hDepartment of Biochemistry and Molecular Biophysics, Columbia University, New York, NY 10032; and ⁱCAS Center for Excellence in Biomacromolecules, Chinese Academy of Sciences, 100101 Beijing, China

Contributed by Wayne A. Hendrickson, March 24, 2021 (sent for review July 20, 2020; reviewed by Julian I. Schroeder and Robert M. Stroud)

Stomata in leaves regulate gas exchange between the plant and its atmosphere. Various environmental stimuli elicit abscisic acid (ABA); ABA leads to phosphoactivation of slow anion channel 1 (SLAC1); SLAC1 activity reduces turgor pressure in aperture-defining guard cells; and stomatal closure ensues. We used electrophysiology for functional characterizations of *Arabidopsis thaliana* SLAC1 (AtSLAC1) and cryoelectron microscopy (cryo-EM) for structural analysis of *Brachypodium distachyon* SLAC1 (BdSLAC1), at 2.97-Å resolution. We identified 14 phosphorylation sites in AtSLAC1 and showed nearly 330-fold channel-activity enhancement with 4 to 6 of these phosphorylated. Seven SLAC1-conserved arginines are poised in BdSLAC1 for regulatory interaction with the N-terminal extension. This BdSLAC1 structure has its pores closed, in a basal state, spring loaded by phenylalanyl residues in high-energy conformations. SLAC1 phosphorylation fine-tunes an equilibrium between basal and activated SLAC1 trimers, thereby controlling the degree of stomatal opening.

abscisic acid signaling | phosphorylation | channel activation | cryo-EM | electrophysiology

The stomatal pore, formed by a pair of specialized guard cells in plant leaves, serves as the gateway for water transpiration and atmospheric CO₂ influx for photosynthesis (1, 2). These pores need to be tightly controlled, as inadequate CO₂ intake is harmful and excessive water loss is devastating for plants (3). The guard cells respond to a wide range of environmental stimuli, including high CO₂ levels, O₃, low air humidity, and drought, and transduce these signals into appropriate turgor pressure changes that can adjust stomatal pore aperture (4, 5). It is well established that turgor pressure of guard cells is regulated by ion transport across the membrane, notably anions and potassium ions (6–8).

The phytohormone abscisic acid (ABA) plays a central role in controlling stomatal closure via activation of a complex signaling pathway that is mediated by receptors, kinase/phosphatases, and ion channels (5, 9). It has been known that the activation of a slow anion current in guard cells is a key event leading to stomatal closure (10–13). The corresponding gene, slow anion channel 1 (SLAC1), was discovered in genetic screens for O₃ or CO₂ sensitivity in *Arabidopsis*; when this particular channel was mutated out, the mutants showed impaired response to stimuli (14–16).

Arabidopsis thaliana SLAC1 (AtSLAC1) contains 556 amino acid (aa) residues, with a conserved transmembrane domain (TMD) between hydrophilic tails at both the N and C termini (15). Within the ABA signaling pathway, SLAC1 channel activity is controlled by a protein kinase-phosphatase pair (OST1/ABI1) (17–19). In the absence of ABA, OST1 kinase is bound and inhibited by ABI1 phosphatase (20–22). Under drought condition, ABA is accumulated into guard cells and perceived by the PYR1 receptor (23, 24). This resulting hormone–receptor complex then binds to phosphatase ABI1 to form a ternary hormone–receptor–phosphatase

complex (25–28), which activates OST1 kinase, and leads to phosphorylation of SLAC1 (17, 18). Subsequently, SLAC1 releases anions (Cl[−] and NO₃[−]) from guard cells, leading to membrane depolarization. Depolarization in turn activates potassium channels to release K⁺ via K⁺ channels and drives water out of the cell, decreasing guard cell turgor to close down the stomatal pore (8).

It has been shown that SLAC1 activation is due to phosphorylation at its N terminus (17, 18, 29–31). The phosphorylation of S120 by OST1 is essential but not sufficient for SLAC1 activation monitored in *Xenopus* oocytes (18). S59 has been shown to be the target for both OST1 and other kinases (29, 31). In another phosphoproteomic analysis, using the fragment SLAC1^{1–186} as substrate, S86 and S113 were also shown to be phosphorylated by OST1 (32). OST1 also phosphorylates SLAC1 at the C terminus (17), with T513 identified as a potential phosphorylation site for SLAC1 activation (31). Although multiple phosphorylation sites have been proposed, the underlying mechanism for SLAC1 activation remains elusive.

Significance

Stomatal openings in plant leaves admit CO₂ and release water vapor between plant and atmosphere. Environmental stimuli such as darkness or dryness lead to closure, whereas light or low CO₂ stimulate opening. Each stoma is defined by a pair of guard cells, and its pore aperture is controlled by ion channels that regulate the turgor pressure in these cells. SLAC1 anion channels are of central importance here, and environmental factors affect channel activity by controlling SLAC1 phosphorylation. Our studies define multiple sites of regulatory phosphorylation in the context of an atomic structure of the SLAC1 channel, providing a mechanistic understanding of the fine-tuning of channel activity and thereby of stomatal apertures in response to the environment.

Author contributions: Y.-h.C. designed research; Y.-n.D., Q.W., G.-x.Z., Q.-y.L., L.-h.T., H.-l.G., C.-r.Z., L.Q., M.S., X.-h.H., O.B.C., and Y.-h.C. performed research; Y.-n.D., H.K., F.L., Y.-c.W., Q.X., O.B.C., W.A.H., and Y.-h.C. analyzed data; and H.K., W.A.H., and Y.-h.C. wrote the paper.

Reviewers: J.I.S., University of California San Diego; and R.M.S., University of California, San Francisco.

The authors declare no competing interest.

Published under the PNAS license.

¹H.K. and Q.W. contributed equally to this work.

²To whom correspondence may be addressed. Email: oc2188@cumc.columbia.edu, wah2@cumc.columbia.edu, or yuhang.chen@genetics.ac.cn.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2015151118/-DCSupplemental>.

Published April 29, 2021.

Our previous study on a SLAC1 bacterial homolog from *Haemophilus influenzae* (*HiTehA*) revealed a superfamily of trimeric anion channels (33) and found each protomer gated by a highly conserved phenylalanine. We observed an unusual high-energy rotameric conformation for this gating phenylalanine (F262 in *HiTehA*), and further mutational study on the corresponding site in *AtSLAC1* (F450) suggested a unique mechanism for channel activation (33). Nevertheless, since *HiTehA* shares a mere 19% in sequence identity with *AtSLAC1* and lacks the substantial N and C termini that are essential for channel activation, questions remain about SLAC1 structure and its regulation by kinase phosphorylation (7).

Here we describe the cryoelectron microscopy (cryo-EM) structure of SLAC1 from *Brachypodium distachyon* (*BdSLAC1*) at 2.97-Å resolution, and we further characterize the channel activity on *Arabidopsis* SLAC1. *BdSLAC1* shares 63% overall sequence identity with *AtSLAC1*, with 73% for the pore-forming TMD and 48% for the regulatory N and C termini. As for the bacterial homolog, each *BdSLAC1* TMD comprises five tandemly repeated helical hairpins that surround a transmembrane pore in each protomer of the trimeric assembly. The SLAC1 pores are electropositive, consistent with its function as an anion channel. The regulatory N-terminal domain (190 aa) and C-terminal domain (65 aa) are flexibly associated and not discernible in the structure.

We employed a SLAC1::OST1 fusion design for examining SLAC1 phosphorylation, whereby we systematically characterized critical sites responsible for channel activation. Inspired by the structure, we also conducted mutagenesis on the highly conserved phenylalanine residues related to channel gating. Altogether, our structural and functional analyses provide insights into SLAC1 gating and activity modulation. These findings allow us to propose a mechanism for finely tuned SLAC1 control of the stomatal pore in response to environmental stimuli.

Results

Biochemical Production and Structure Determination of Plant SLAC1.

Fifteen plant SLAC1s were selected for gene synthesis and cloned into a modified pREP1 vector with capture tags for expression in the fission yeast *Schizosaccharomyces pombe* (34). Among those 15 SLAC1s, *BdSLAC1* had the best expression and purification yield; however, it is not stable in most detergents, such as the commonly used dodecylmaltoside (DDM). Improved stability and gel-filtration profiles were achieved by using a styrene-maleic acid (SMA) polymer to extract the SLAC1 protein into SMA lipid particles (SMALPs) by using a detergent-free procedure for membrane protein solubilization (35) (*SI Appendix, Fig. S1A*). When solubilized in 2% SMA polymer, *BdSLAC1* was a suitable candidate for structural and functional characterization. No additional SMA polymer or detergents were needed to maintain the solubilized proteins in solution. We determined the structure of *BdSLAC1* by cryo-EM at 2.97-Å resolution (Fig. 1A and *SI Appendix, Fig. S1 B–H* and Table S1).

The cryo-EM map for the *BdSLAC1* TMD is of high quality (Fig. 2A), allowing us to build an accurate model containing residues 191 to 385 and 395 to 512. Owing to flexibility, neither N- nor C-terminal segments are resolved in the current structure. Intriguingly, perhaps as a benefit of using SMA rather than detergent for membrane extraction (36), we found densities for two lipids per protomer. Both are positioned as from the outer leaflet. Three single-chain molecules (perhaps sphingolipids from yeast) are bundled at the threefold axis of the trimer, and what seems to be a typical diglyceride binds at peripheral sites between protomers (Fig. 2A and B).

Structure of Plant SLAC1. The *BdSLAC1* is a symmetric trimer, and subunits are tightly associated, burying a total surface area of ~6,600 Å² within interfaces. Remarkably, *BdSLAC1* has positive electrostatic potential surface on both the extracellular and cytoplasmic

sides (Fig. 1B), whereas bacterial *HiTehA* has positive charges only on the cytoplasmic surface. Each SLAC1 protomer has 10 TM helices, adopting similar topology as *HiTehA* (33). They are arranged as five pairs of helical hairpins that are assembled into a unique protein fold with quasi-fivefold symmetry. TM_{odd} helices create a channel pore across membrane, which is blocked by a highly conserved phenylalanine residue (F460 in *BdSLAC1*). TM_{even} helices are longer and straighter, but more inclined, surrounding the inner pore and forming an outer layer (Fig. 1C). Short juxtamembrane amphipathic helices, H_{0,1} before TM₁ and H_{2,3} between TM₂ and TM₃, are located near the N terminus on the cytoplasmic side.

Similar to TRIC (34), another trimeric channel, SLAC1 also has three independent pores, one through each protomer. Interestingly, highly conserved Pro or Gly residues are located in the middle of the pore-forming helices (Gly in TM₁/TM₅, Pro in TM₃/TM₇/TM₉), which generate kinks and contribute to formation of a relatively uniform pore across membrane (Fig. 2C). The pore has a diameter of ~6 Å, except for blockage by the phenyl ring of highly conserved F460 near the middle of the membrane. Highly conserved positive charged residues, including Arg or Lys, are found near the channel pore at both cytoplasmic and extracellular sides, rendering the pore surface electropositive, consistent with SLAC1 function as an anion-conducting channel.

Despite limited sequence similarity, plant SLAC1 shares similar overall topology and trimeric assembly architecture with bacterial *HiTehA* (33). Nevertheless, there are distinctive differences between plant SLAC1 and bacterial TehA. When *BdSLAC1* is compared to *HiTehA*, the rmsd is 1.90 Å for 269 superimposed C_α of the protomer, and is 2.25 Å for 793 superimposed C_α of the trimer. The most divergent regions are located near loops connecting the TM helices. A noticeable distinction is near the middle of TM₅, where *HiTehA* has a π -helical turn caused by the insertion of A146 beside the P145-mediated kink (*SI Appendix, Fig. S2*). Another difference is at the middle of TM₉ where a proline-mediated kink (P461 in *BdSLAC1*) replaces a water-mediated kink at a glycine residue (G263 in *HiTehA*), but nevertheless maintains a similar conformation. Additionally, a short juxtamembrane helix (H_{0,1}) connects the flexible regulatory N-terminal domain to TM₁ in SLAC1 (packing against H_{2,3}).

A TehA-based homology model inspired our previous SLAC1 mutagenesis for functional analysis on gating and selectivity; however, inadequate similarity between TehA and SLAC1/SLAHs also caused errors in this and other homology modeling. The alanine insertion in TM₅ of TehA led to misplaced predictions of pore-lining residues in that helix (33, 37), and a two-residue misalignment in TM₇ led to the prediction of a tryptophan residue (W404 in *AtSLAC1*, and W356 in *AtSLAH2*) as pore lining (37), whereas actually it is located opposite to the pore in *BdSLAC1*. Obviously having an actual SLAC1 structure provides a much improved basis for studying plant SLAC1 or SLAH channels.

SLAC1 Activation by OST1 Kinase. There were puzzles remaining for SLAC1 activation via OST1 phosphorylation. Coexpression of *Arabidopsis* SLAC1 with OST1 in *Xenopus* oocytes only produced weak currents, whereas their coexpression with split-YFP association (SLAC1::YFP^C and OST1::YFP^N) resulted in moderate stimulation (17, 18, 33). The poor activation of coexpression of SLAC1 and OST1 individually may be due to the lack of sufficient interaction between channel and kinase, while enhanced currents were achieved with the physical linkage from split-YFP constructs. These difficulties complicated the identification residues responsible for activating phosphorylation. To reconstitute and optimize channel activation via OST1 phosphorylation, we created a construct of SLAC1 fused directly with OST1, (SLAC1::OST1, named SOF) and performed electrophysiological measurement in oocytes after cRNA injections (Fig. 3A). We observed remarkably enhanced macroscopic current from the SLAC1::OST1 construct,

Fig. 1. Cryo-EM structure of *BdSLAC1*. (A) *BdSLAC1* trimer embedded in SMALP, top view (Above) and side view (Below). The protomers are colored in blue, green, and red. The transparent surface indicates the boundary of the SMALP nanoparticle. (B) Electrostatic potential at the extracellular (Left) and intracellular (Right) surfaces. Electronegative and electropositive potential are colored in degrees of red and blue saturation, respectively. One protomer is drawn as a ribbon diagram and colored spectrally from blue at its N terminus to red at its C terminus. Stars indicate the position of highly conserved arginine residues (R196, R265, R272, R331, R332, R385, and R442). (C) Ribbon diagram of a *BdSLAC1* protomer is colored spectrally as in B and viewed from outside (Left) and within (Right) the membrane. The side chains of F460 and F285 are shown as stick bonds within the channel pore.

much greater than in the split-YFP assay (18). To validate SLAC1 activation by OST1 phosphorylation, we disrupted the OST1 kinase activity by introducing a function impaired D160A mutation (38), and observed no apparent current for the SLAC1::OST1(D160A) construct. We further confirmed ABI1 inhibition to OST1 activity via coexpression of SLAC1::OST1 with ABI1 (*SI Appendix, Fig. S3*).

As our previous work suggested that conserved F450 plays an important role in channel gating, we then analyzed channel conductance properties for SLAC1 variants (wild type [WT], F450L, and F450A), in the presence and absence of OST1 and in the SLAC1::OST1 fusion construct. Compared to the coexpression of SLAC1/OST1 separately, the exceptionally large currents in SLAC1::OST1 construct suggests highly efficient channel activation. Removal of the phenyl ring in F450A SLAC1 led to increased currents both in absence and presence of OST1; however, the F450A SLAC1::OST1 current is much reduced from that with wild-type SLAC1::OST1 (Fig. 3 *B* and *C*). Consistent with our previous study, the F450L mutant shows negligible currents in all three cases (Fig. 3*D*). Puzzles from these findings are examined further below, but it seems clear that conserved gating phenylalanine F450 is crucial for SLAC1 activation.

Characterization of Phosphorylation Sites. We generated the same SLAC1::OST1 construct for expression in *S. pombe*. We expressed, purified, and performed mass spectrometry analyses on the fusion

protein (*SI Appendix, Fig. S4A*). Unexpectedly, we found 14 sites of phosphorylation, including 13 in the N-terminal domain and 1 in the C-terminal segment (Fig. 4*A* and *SI Appendix, Fig. S4 B–F* and Table S2). Two of these sites (S59 and S120) have been characterized previously (18, 29–32), whereas the other 12 sites are newly identified in the current study. Whether and how these potential sites are involved in SLAC1 activation becomes an intriguing question.

To understand the role of these sites, we generated single mutants by substituting the identified candidates, serine or threonine to alanine in the SLAC1::OST1 construct, and performed conductance measurement in oocytes. We observed notable current reduction for four sites (S59, S86, S113, and S120), suggesting critical roles for these in channel activation (Fig. 4*B* and *SI Appendix, Fig. S5A*). Currents were further reduced in pairwise combinations, and more still in the quadruple substitution. Moreover, the addition of T114A or/and S116A at patch 1 (S113, T114, S116, and S120) virtually abolished currents, and the patch1 quadruple mutation itself also nearly eliminated current. By contrast, the quintuple mutation at patch 2 (S134, T137, T142, S146, and S152) only slightly reduced the current (Fig. 4*C*). We conclude from these observations that six key sites, including S59, S86, S113, T114, S116, and S120, play important roles for SLAC1 activation by OST1 phosphorylation.

To confirm the above observation, we further designed a SLAC1::OST1 mutant construct (named SLAC1::OST1-13A or SOF-13A), which replaced all 13 phosphorylation sites in the

Fig. 2. The cryo-EM map for *BdSLAC1*. (A) Density distribution from *BdSLAC1* cryo-EM at 2.97 Å and superimposed model (protein, yellow; lipids, orange). The central lipid bundle and the peripheral lipid are shown. (B) Representative cryo-EM density map for the TM_{odd} segments. The helices are drawn from the unrolled pore, as oriented in Fig. 1C. The maps are contoured at a level of 0.67, and density is shown within a 2.5-Å zone around the atoms belonging to each segment, using the “sop zone” command in UCSF Chimera. (C) Kinks in the pore-forming TM_{odd} helices, orientated as in B. The kink-mediated Pro or Gly residues (G207 in TM_1 , P279 in TM_3 , G347 in TM_5 , P407 in TM_7 , and P461 in TM_9) are shown and colored magenta.

N-terminal domain to alanine residues. The SOF-13A shows little current activity; however, substantial current was restored upon removal of the phenyl ring of F450 in SOF-13A_F450A, compared to the SLAC1 F450A mutant. This demonstrates that the lack of current in SOF-13A is due to loss of phosphorylation, rather than failure of surface expression in oocytes (SI Appendix, Fig. S5B). This observation also suggests that the reduced currents of other such mutants (singly and in combination) most likely reflect impaired channel activity due to the reduced phosphorylation. Next, we examined functional significance of the six key residues by restoring them (singly or in combination) back into the SOF-13A construct (Fig. 4D). Among these, only A113S shows large current, which is enhanced appreciably for the A86S/A113S double restoration. SLAC1 activity was completely restored when four to six key phosphorylation sites were restored back into SOF-13A (Fig. 4D). These results confirm the

critical roles for these six key sites in SLAC1 activation, and combined phosphorylation of multiple sites potentiates SLAC1 to full activity.

Additionally, we created phosphomimetic mutants for the six key sites by substitution to aspartate residues. Single phosphomimetics did not produce noticeable currents, whereas combinations of two to six sites gradually stimulated channel activities to a substantial level (Fig. 4E and SI Appendix, Fig. S5 C and D). Overall, these results suggest synergetic phosphorylation of multiple sites can provide a fine-tuning mechanism for modulating various SLAC1 activities.

Some puzzles remain regarding C-terminal phosphorylation sites. Although S543 was phosphorylated in the mass spectrometry analyses, we observed no current reduction upon substitution to alanine in the SLAC1::OST1 construct, and found no stimulation for S543D SLAC1. On the other hand, another study showed that phosphomimetic mutant T513D stimulated currents (31), even though we detected no phosphorylation in our experiment. We reproduce the finding on T513D and also fail to observe phosphorylation on T111, a previously unreported site. T111A substitution showed no current reduction in SLAC1::OST1, but the phosphomimetic mutant T111D did stimulate currents in SLAC1 (SI Appendix, Fig. S5 E and F). These observations indicate that both T111 and T513 may play important roles in channel activation; yet further investigation is needed.

SLAC1 Pore and Gating Mechanism. Little is known about how SLAC1 phosphorylation is coupled to channel gating, but new clues come from structural features of pore-lining residues in our current structure. The SLAC1 pore is lined with highly conserved and generally hydrophobic residues (Fig. 5 A and B). The SLAC superfamily shares an extremely conserved gating phenylalanine (F450 in *AtSLAC1*, F460 in *BdSLAC1*, and F262 in *HtTehA*). In *HtTehA*, the phenyl ring (F262) is restricted in a high-energy conformation ($-160^\circ/-4^\circ$ for χ_1/χ_2), by contacts with neighboring L18 (TM_1) and V210 (TM_7) (33). The corresponding residues in *BdSLAC1* are L206 and A409, respectively; and both are conserved in plant SLAC1s. The replacement of Val by Ala would partially release the constraint on the gating phenylalanine. In *BdSLAC1*, the corresponding gating phenylalanine F460 also adopts a strained conformation ($159^\circ/47^\circ$ for χ_1/χ_2). We found that another pore-lining phenylalanine (F285 in *BdSLAC1*, TM_3) lays in a close van der Waals contact with the gating phenylalanine F460, having a χ_1/χ_2 angle of $-94^\circ/-31^\circ$. Both phenylalanine residues are highly conserved in plant SLAC1s (F267 and F450 in *AtSLAC1*). Different from bacterial *HtTehA*, these dual phenyl rings interact and render both in energy unfavorable conformations, suggesting a critical role in channel gating (Fig. 5C). In addition, the kink at the middle of TM_9 in *HtTehA* (mediated by G263 and a water molecule) is replaced by a proline-mediated kink in SLAC1 (P451 in *AtSLAC1* and P461 in *BdSLAC1*).

Taken together, distinguishing structural characteristics of the *BdSLAC1* pore indicate that the mechanism for channel activation in plant SLAC1 is distinctive from that in bacterial homologs. The conformation of proline-kinked TM_9 in SLAC1 is similar to that with the water-mediated kink in *TehA* (SI Appendix, Fig. S6A), but the distinctions are such that they impact pore geometry and likely affect conductance. In examining the gating mechanism in *Arabidopsis* SLAC1, we first substituted P451 with glycine (as in bacterial *TehA*) and also alanine for testing in SLAC1::OST. These mutations reduced channel currents markedly, respectively, to 36% and 24% of wild type (SI Appendix, Fig. S6 B and C), likely due to impaired channel activities or possibly from altered trafficking to the plasma membrane.

We also mutated the pore phenylalanine residues, F276 and F450, to alanine in both SLAC1 and SLAC1::OST, and measured resulting conductance in oocytes (SI Appendix, Fig. S7). Compared to the small but significant current in the wild type,

Fig. 3. Electrophysiological characterization of AtSLAC1::OST1. (A, Above) Construct design for SLAC1::OST1 (SOF); AtSLAC1 is fused directly with AtOST1. (Below) Typical *Xenopus* oocyte microelectrode voltage-clamp current traces (−110 mV to 70 mV in 20 mV steps). (B and C) Current-voltage (I-V) relations for *Arabidopsis* SLAC1 wild type, F450A mutant. Note different y axis scales. (D) Effects of gating residue mutations. Mean chloride currents, measured at −90 mV, are shown comparing WT AtSLAC1 and its variants (F450A and F450L), alone, coexpressed with OST1, and SLAC1::OST1.

the F276A mutant showed enhanced leaky conductance with altered properties in the absence of OST1, suggesting a role for F276 in channel gating (Fig. 5D). The roles of F276 and F450 are complicated, however. The F276A, F450A, and F276A/F450A mutants all increased currents relative to wild type when in unphosphorylated SLAC1, by 154%, 2100%, and 668%, respectively, whereas they decreased currents to 62%, 21%, and 58% of wild type, respectively, when in the SLAC1::OST1-activated channel (Fig. 5E).

There is no simple explanation for the results from pore-phenylalanine mutations; effects of the phenyl ring deletions are opposite in unphosphorylated and phosphorylated channels. The pore as restructured after activation must differ substantially, and at present indecipherably, from the *BdSLAC1* basal-state pore. It makes sense from the basal-state structure that the current should be greatly enhanced when the phenyl ring is removed in F450A, but the substantial reduction of that current in the F276A/F450A double mutant implies added conformational change. The relatively modest increase of current upon phosphorylation of F450A (3.3-fold) compared to the ~330-fold increase for the WT channel (Fig. 3D) suggests radical change. Furthering the complications, another study has found that F450A leads to large currents in oocytes, but not in the absence of OST1 (39); this latter observation conflicts with our findings here and those reported previously (33). Activation might plausibly entail pore-opening quaternary rearrangements in the WT trimer, and the propensity for such activation might differ for different mutants. Indeed, the virtually complete loss of phosphoactivation in the F450L channel may simply mean that this mutant cannot undergo conformational activation.

Discussion

Plants must maintain a delicate balance between gas exchange necessary for photosynthesis and water conservation in the surrounding environment, which they do by regulating stomatal pores in the leaf epidermis. Environmental factors, such as elevated CO₂ levels, darkness, and drought, activate SLAC1 channels to induce stomatal closure through reduction of turgor pressure in guard

cells. The SLAC1 activation is controlled largely by OST1 phosphorylation (6–8), and SLAC1 activity adjusts the turgor pressure in guard cells and thereby controls stomatal apertures; however, the underlying mechanism for SLAC1 phosphoactivation is largely unknown.

Following the leads of others, we have studied SLAC1 activity by expression in *Xenopus* oocytes to gain insight into OST1 phosphorylations of SLAC1, recognizing that SLAC1 regulation in plant cells can differ from that obtained in oocytes. The complement of protein kinases and other relevant factors in plant guard cells likely differs from those present in oocytes, and differing SLAC1 channels may be presented differently onto oocyte plasma membranes. Nevertheless, despite the reservations, we have gained useful insights from electrophysiology in oocytes that complement our structural analysis. Here too, we face limitations. We were able to perform the functional studies with *Arabidopsis* SLAC1 and the structural analysis on *Brachypodium* SLAC1; and, whereas ideally we would have both functional and structural data from the very same SLAC1 molecules, we must rely on the high sequence similarity (63% aa identity overall, 73% for the transmembrane domain) in drawing conclusions.

Previous studies in *Xenopus* oocytes showed that OST1 alone barely activates SLAC1, usually at the level significantly less than those observed by CDPKs or CIPKs. A possible explanation could be weak or transient interaction between SLAC1 and OST1, perhaps due to the lack of an unknown mediating factor in the heterologous system. Indeed, when OST1 and SLAC1 were drawn together through split YFP association (SLAC1::YFP^C and OST1::YFP^N), SLAC1 activation via OST1 phosphorylation in oocytes was enhanced substantially (18). We suspected that the split YFP encounters may not have been complete, and, so we took away the problem by using the direct fusion SLAC1::OST1 to assure avidity. Thereby, OST1 phosphorylation enhanced *AtSLAC1* channel activity ~330-fold (Fig. 3D). Another explanation for low SLAC1 activation by simple coexpression of SLAC1 and OST1 has recently come to light from work of Takahashi et al. (40), suggesting that MAPKKK kinase can activate OST1 by phosphorylating OST1

Fig. 4. Characterization of phosphorylation sites in AtSLAC1. (A) Phosphorylation sites identified in mass spectrometry by using SLAC1::OST1. All phosphorylated residues are indicated, 13 at the N terminus and 1 at the C terminus. (B) Effects of single alanine mutations in SLAC1::OST1. Mutations S59A, S86A, S113A, and S120A resulted in substantially reduced currents. The asterisks indicate significance of one-way ANOVA analyses and Tukey's test: *** $P < 0.01$ and **** $P < 0.0001$. (C) Effects of mutant combinations in SLAC1::OST1. Double mutants (S59A/S120A, S86A/S113A, and S113A/S120A) and quadruple mutant (S59A/S86A/S113A/S120A) significantly reduced currents, and additions of such S/T-A mutations to that quadruple mutant essentially eliminated currents. The patch-1 quadruple mutant (S113A/T114A/S116A/S120A) almost abolished channel current, whereas the patch-2 quadruple mutant (S134A/T137A/T142A/S146A/S152A) showed no effect. (D) Effects of restoring key sites (S59, S86, S113, T114, S116, and S120) back into SLAC1-13A::OST1. The SLAC1-13A::OST1 construct contains 13 alanine mutations for the identified phosphorylation N-terminal-domain residues, and serves as a "blank" control construct. (E) Electrophysiological analyses of SLAC1 phosphomimetics. Aspartate substitutions of two to six sites gradually restore channel activities to a substantial level (in the absence of OST1) (mean $\pm$ SEM, $n > 8$).

Fig. 5. The SLAC1 channel gating. (A) Cross-section through the model of *Bd*SLAC1 F460A. The model is oriented as in Fig. 1C, with the electrostatic potential (Left) and surface conservation (Right) shown on the external surface of the molecular envelope. (B) Pore-lining residues are indicated in TM_{odd} helices of the *Bd*SLAC1 rolled-open model. (C) Molecular basis for conformational strain between the highly conserved pore lining phenyl residues F460 and F285 (respectively, corresponding to F450 and F276 in *At*SLAC1). The pore-forming TM_{odd} helices are shown and presented as ribbon diagrams with selected pore-lining side chains drawn in stick representation, top view (Left) and side view (Right, with TM₅ removed). The dual phenyl rings interact and render both in energetically unfavorable conformations, with $\chi_1/2$ at $-94^\circ/-31^\circ$ (F285) and $159^\circ/47^\circ$ (F460), suggesting critical roles in channel gating. (D) Representative current traces of SLAC1 and F276 SLAC1. (E) Mean chloride currents, measured at -90 mV, are shown comparing WT, F276A, F450A, and F276A/F450A mutant, in the absence of OST1 and in SLAC1::OST1.

S171; however, native background MAPKKK activity seems to be missing in *Xenopus* oocytes. Coincidentally, we found that OST1 S171 was phosphorylated in the SLAC1::OST1 fusion when expressed in *S. pombe* (SI Appendix, Table S2).

The present study appreciably advances understanding of the mechanism for phosphoactivation of SLAC1. Besides obtaining an atomic-level picture of *Bd*SLAC1, we identified 14 sites of phosphorylation by using an *At*SLAC1::OST1 fusion construct to enhance avidity of the kinase–substrate interaction. Among them, six residues located in the flexible N-terminal domain (S59, S86, S113, T114, S116, and S120) all play key roles in activating the channel. Whether and which of these sites are phosphorylated *in vivo* in plants, by OST1 and other kinases, needs further investigation. It is also unclear whether any phosphorylation occurs in the SLAC1 transmembrane domain, from which we have not observed any phosphorylated peptides in the mass spectrum of the purified SLAC1::OST1 protein.

Phosphorylation is one of the most significant of posttranslational modifications, used to modulate channels and transporters in many signaling pathways and by diverse mechanisms. Some transport proteins are inactive when their cytoplasmic loops/tails block the channel pore at resting state (41, 42). Phosphorylation usually induces conformational changes to release inhibiting blockage in these proteins, and in turn renders them active. Phosphorylation of plant aquaporin (*SoPIP2*) rearranges its cytoplasmic gate-blocking loop to open the channel (41). Another example is the CFTR chloride channel, for which phosphorylation disengages its regulatory domain from an inhibitory position and leads to channel opening (42). In yet another case,

phosphorylation switches the dual-affinity nitrate transporter NRT1.1 from the low-to-high-affinity mode by decoupling the dimer and increasing structure flexibility (43, 44). What then is the basis for SLAC1 channel activation after phosphorylation? As we have seen, efficacious SLAC1 phosphorylation sites are at *At*SLAC1 positions 59, 86, and a patch of four residues among 113 to 120, each imbedded in an island of conservation; yet, despite especially strong juxta-TMD conservation (*Bd*SLAC1 residues 154 to 188), the entire N-terminal domain (1 to 188) is invisible in the current *Bd*SLAC1 structure and presumed to be flexibly associated. Thus, the actual mechanism of SLAC1 phosphoregulation remains elusive.

In principle, we contemplate two alternative mechanisms: one a binding-activation model in which phosphorylated domains interact with the TMD to exert pore-opening conformational changes, and the second an inhibition-release model in which interactions that hold the channel in a closed state would be released upon phosphorylation. We initially disfavored the latter since baseline channel activity and stomatal openings are the same for $\Delta N/\Delta C$ SLAC1 as for WT SLAC1 (45), even though ABA responsiveness was abolished, and because the N-terminal domain is only flexibly associated and not TMD bound in the cryo-EM structure. Moreover, as for the binding-activation model, we could imagine phosphorylated N-terminal domain residues being attracted to the positively charged cytoplasmic TMD surface of SLAC1, which features seven arginine residues that are identical in all SLAC1-identified sequences (R187, R256, R263, R321, R322, R375, and R432 in *AtSLAC1*) and five sites having either lysine or arginine residues at >94% incidence (K255, K320, K325, K384, and K440 in

Puzzles that remain for these and other future investigations include the conformational changes that SLAC1 undergoes upon phosphorylation, the mechanism by which the phosphorylated N-terminal domain of SLAC1 promotes the activated conformation, possible allosteric modulation by bicarbonate (39) and other ligands, whether calcium-dependent kinases and other kinases play complementary as well as overlapping roles, the role of lipids in channel stability and/or activity, and the structural basis for anion selectivity. Deep insight into the mechanism of SLAC1 control on stomatal pores will be valuable for designing drought-tolerant or water-efficient plant cultivation.

Protein Production and Structure Determination. A structural genomics approach, screening 15 plant species, was used to identify *BdSLAC1* as being suitable for expression in *S. pombe* for use in structure determination. Purified *BdSLAC1* was applied to gold grids, frozen in a TFS Vitrobot, and micrographs were collected in a Titan Krios equipped with a K2 detector. Images were motion corrected, particles were picked and classified, three-

ACKNOWLEDGMENTS. We thank Dr. Guang-zhe Yang for suggestions on electrophysiological experiments and De-lin Li, Ling-min Yuan, and other colleagues for expression screening in the early stages of the project. We are also grateful to the staff at the Center for Biological Imaging, Institute of Biophysics, Chinese Academy of Sciences. This project was financially supported by the National Key Research and Development Program of China (Grants 2016YFA0500503 and 2020YFA0509903), the Strategic Priority Research Program of the Chinese Academy of Sciences (Precision Seed Design and Breeding, Grant XDA24020305), and the National Natural Science Foundation of China (Grants 31470728, 31322005, and 31872721) to Y.-h.C., by NIH Grant R35GM134920-01 and NSF Grant MCB-1934628 to F.L., and by NIH Grants R01GM107462 and P41GM116799 to W.A.H.