Identification and Characterization of *Mycobacterium smegmatis* and *Mycobacterium avium* subsp. *paratuberculosis* Zinc Transporters

Elke Goethe,^a Ayla Gieseke,^a Kristin Laarmann,^a Janita Lührs,^a Ralph Goethe^a

^aInstitute for Microbiology, Department of Infectious Diseases, University of Veterinary Medicine Hannover, Hannover, Germany

ABSTRACT Zinc uptake in bacteria is essential to maintain cellular homeostasis and survival. ZnuABC is an important zinc importer of numerous bacterial genera which is expressed to restore zinc homeostasis when the cytosolic concentration decreases beyond a critical threshold. Upon zinc limitation, the fast-growing nonpathogenic organism *Mycobacterium smegmatis* (MSMEG) as well as the ruminant pathogen *M. avium* subsp. *paratuberculosis* (MAP) increases expression of genes encoding ZnuABC homologues but also of genes encoding other transporters. This suggests an involvement of these transporters in zinc homeostasis. Here, we characterized the putative zinc transporters of MSMEG (ZnuABC and ZnuABC2) and MAP (ZnuABC, MptABC, and MAP3774-76). Deletion of either ZnuABC or ZnuABC2 in MSMEG did not lead to growth defects but did lead to an increased expression of zinc marker genes in MSMEGΔ*znuABC*, indicating cytosolic zinc limitation. However, chromatin immunoprecipitation proved direct binding of the global zinc regulator Zur to promoter regions of both *znuABC* and *znuABC2*. Simultaneous deletion of both transporters caused severe growth defects, which could be restored either by homologous complementation with single ZnuABC transporters or supplementation of growth media with zinc but not iron, manganese, cobalt, or magnesium. Heterologous complementation of the double mutant with MAP transporters also resulted in reconstitution of growth. Nonradioactive FluoZin-3AM (FluoZin-3 acetoxymethyl derivative) zinc uptake assays directly revealed the competence of all transporters to import zinc. Finally, structural and phylogenetic analyses provided evidence of a novel class of ZnuABC transporters, represented by the ZnuABC2 of MSMEG, which is present only in actinobacteria, mainly in the genera *Nocardia* and *Streptomyces* and in fast-growing mycobacteria.

IMPORTANCE Zinc is necessary for bacterial growth but is toxic when present in excess. Hence, bacterial cells have developed systems to alter intracellular concentration. Regulation of these systems is primarily executed at the transcriptional level by regulator proteins which sense femtomolar changes in the zinc level. In environmental and pathogenic mycobacteria, zinc starvation induces expression of common zinc import systems, such as the ZnuABC transporter, but also of other additional not-yet-characterized transport systems. In this study, we characterized the role of such systems in zinc transport. We showed that transport systems of both species whose transcription is induced upon zinc starvation can exchangeably restore cellular zinc homeostasis in transporter-deficient mutants by transporting zinc into the cell.

KEYWORDS metal transport, mycobacteria, zinc transporter, zinc uptake, *znuABC*

Zinc is an essential micronutrient for all living cells. It is of major importance in numerous cellular processes and in host-pathogen interactions (1). Its availability is crucial for bacterial survival and growth, both in the environment and in the host.

Citation Goethe E, Gieseke A, Laarmann K, Lührs J, Goethe R. 2021. Identification and characterization of *Mycobacterium smegmatis* and *Mycobacterium avium* subsp. *paratuberculosis* zinc transporters. *J Bacteriol* 203:e00049-21. <https://doi.org/10.1128/JB.00049-21>.

Editor Tina M. Henkin, Ohio State University

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Elke Goethe, elke.goethe@tiho-hannover.de, or Ralph Goethe, ralph.goethe@tiho-hannover.de.

Received 29 January 2021

Accepted 4 March 2021

Accepted manuscript posted online 15 March 2021

Published 7 May 2021

Despite being highly abundant in the intra- and extracellular environments, bioavailability of free zinc is in the picomolar range, since nearly all zinc is protein bound (2–4). This is essential, since excess free zinc is toxic for cells (5, 6). During bacterial infection, host cells can actively alter zinc concentration (e.g., by the production of metal-binding proteins and zinc transporters) to either starvation or excess (7, 8). This process, which belongs to the antimicrobial host defense (so-called nutritional immunity), is directed to perturb bacterial protein function, due to missing or false incorporation of metal ions, and consequently loss of cell vitality. On the other hand, bacteria maintain their own zinc homeostasis by the expression of zinc storage, export, and uptake systems in response to high or low availability of the metal (9). This allows fast adaptation to ecological niches with different zinc availability encountered by the bacterium.

The high-affinity zinc importer ZnuABC is a crucial factor in the sensitive equilibrium of zinc homeostasis in many bacterial species. This ATP-binding cassette (ABC) transporter is constituted by the proteins ZnuA, a soluble periplasmic metallochaperone, ZnuB, which constitutes a membrane permease, and ZnuC, an ATPase (10). The importance of this transporter has been demonstrated, e.g., for *Escherichia coli*, *Vibrio cholerae*, *Pseudomonas aeruginosa*, *Salmonella enterica*, and *Yersinia pestis* (11–18). In these species, ZnuABC deletion resulted in growth defects and/or impairment of pathogenicity.

The genes encoding ZnuABC homologues are also present in many mycobacterial species (19–21). Hints of an involvement of ZnuABC transporters in zinc metabolism in mycobacteria were provided by studies addressing zinc starvation-induced gene expression. Such studies showed enhanced expression of *znuA*, *znuB*, and partially *znuC* upon zinc starvation in *Mycobacterium tuberculosis* (MTB), the environmental species *Mycobacterium smegmatis* (MSMEG), and the ruminant pathogen and causative agent of paratuberculosis *Mycobacterium avium* subsp. *paratuberculosis* (MAP). In addition, it was demonstrated that *znu* genes were repressed by the Fur-like family regulator Zur in MTB and MSMEG (19, 20), which was also shown for other bacterial species, such as *E. coli* and *Bacillus subtilis* (12, 22). Interestingly, our recent studies with MSMEG (20) and MAP (21) revealed the presence of additional zinc starvation-regulated operons predicted to encode ABC-type transporters. Beside the classical ZnuABC transporter homologue (msmeg_6045-47), a second ABC-type importer (msmeg_6049-51), which was suggested to resemble the ZnuABC2 transporter of *Mycobacterium vanbaalenii* (MVAN) (<https://regprecise.lbl.gov>), was also highly induced upon Zur deletion and zinc starvation in MSMEG (20). The presence of two ZnuABC transporters was also reported for *Streptomyces coelicolor* (SCO) (23, 24) but, to our knowledge, not for other bacteria. Upon zinc starvation of MAP, expression of the operons encoding the classical ZnuABC (*map_0487-89*), the *Mycobacterium paratuberculosis* transporter MptABC (*map_3734-36*) and a predicted Mn^{2+}/Zn^{2+} transporter, MAP3774-76 (*map_3774-76*), were induced. Even though MptABC and MAP3774-76 have also been proposed to be related to iron transport (25–27), *mptA* has clearly been shown to be regulated in a zinc-dependent manner by Zur (21). The presence of additional zinc transporters in MAP is in contrast to MTB (28), and no homologues of MptABC and MAP3774-76 could be identified in the closely related subspecies *Mycobacterium avium* subsp. *avium* and *Mycobacterium avium* subsp. *hominissuis* (21). This might be explained by the location of MptABC and MAP3774-76 on the MAP-specific genomic insertions LSP14 and LSP15, respectively (29), which are part of a MAP-specific zinc-responsive genomic island (ZnGI). The common ZnuABC, also present in other mycobacteria, is encoded in a different region (21). This specific location might indicate a particular relevance of the operons for the zinc metabolism of MAP.

Aside from their regulation by zinc, the structure and direct role of the predicted classical and additional zinc transporters in MSMEG and MAP zinc uptake have not yet been addressed. Hence, we were interested in characterizing the structure of these transporters and clarifying the role of the predicted MSMEG and MAP zinc transporters in zinc homeostasis. For this, we constructed single and double transporter mutations

FIG 1 Growth of MSMEG *znuABC* single mutants in different media. MSwt, mutant strains MSΔ1 and MSΔ2, and the complemented strains MSΔ1::A and MSΔ2::B were grown in MB to an OD₆₀₀ of approximately 2.0, washed twice with PBS, and inoculated at an OD₆₀₀ of 0.05 in MB (A) or 0.1 in SM (B). Growth was monitored daily by measurement of OD₆₀₀ for 4 (A) or 8 (B) days. Shown are the results of three independent replicates in duplicate. (C and D) Substitution of *znuABC* transporters at the transcriptional level. Expression of MSMEG_RS29170 (*znuA2*, primers 27 and 28) or MSMEG_RS29150 (*znuC*, primers 23 and 24) and MSMEG_RS29260 (*rpmG*, primers 37 and 38) in MSΔ1 (C) and MSΔ2 (D) and the complemented strains MSΔ1::A or MSΔ2::B, respectively, was analyzed by qRT-PCR. Shown are the results of at least three independent replicates in duplicate. Statistical analysis was performed using Student's *t* test. *, *P* < 0.05; ****, *P* < 0.001.

in MSMEG, which were complemented either homologously with corresponding MSMEG zinc transporters or heterologously with predicted MAP zinc transporters. Mutants and complemented strains were subjected to growth experiments, zinc uptake assays, and chromatin immunoprecipitation (ChIP) analyses to clarify the regulation and impact of these transporters in zinc homeostasis.

RESULTS

ZnuABC and ZnuABC2 of MSMEG are regulated by Zur. We previously showed that upon zinc limitation, *M. smegmatis* (MSMEG) and the pathogen *M. avium* subsp. *paratuberculosis* (MAP) increase expression of not only the genes encoding ZnuABC homologues but also genes encoding other transporters, suggesting that they are additional zinc importers (20, 21). In MSMEG, we found two predicted ZnuABC transporter homologues encoded by *msmeg_6045-47* (reannotated MSMEG_RS29145-55, ZnuABC^{MSMEG}) and *msmeg_6049-51* (MSMEG_RS29165-75, ZnuABC2^{MSMEG}), and in MAP, we found the predicted ZnuABC transporter homologue genes *map_0487-89* (ZnuABC^{MAP}) as well as two other predicted transporters, MptABC (*map_3734-36*) and MAP3774-76 (20, 21). However, besides the zinc-dependent regulated gene expression, the role of these transporters in zinc homeostasis is still unclear.

Since ZnuABC transporters have been shown to be essential for bacterial growth in other bacterial species, we were first interested in analyzing the impact of single transporters on MSMEG zinc homeostasis by deleting ZnuABC^{MSMEG} and ZnuABC2^{MSMEG}. However, individual deletion of *znuABC* (MSΔ1) or *znuABC2* (MSΔ2) in MSMEG wild type (MSwt) led to slightly reduced growth but no significant growth defects in different media (Fig. 1A and B). Thus, we hypothesized a substitution of the lost transporter by induced expression and activity of the remaining zinc transporter. Therefore, we analyzed the expression of MSMEG_RS29170 (representing *znuA2*) and the zinc marker gene MSMEG_RS29260 (*rpmG*), encoding an alternative ribosomal protein

FIG 2 Impact of Zur on MSMEG *znuABC* and *znuABC2* expression. MSΔ6 complemented with HA-tagged Zur (MSΔ6::L) was grown in standard MB (control) or treated with 10 μ M TPEN or 500 μ M ZnSO₄. ChIP from cross-linked lysates was performed using anti-HA and anti-IgG1 κ (isotype control). DNA from ChIP and input control was subjected to qRT-PCR with primers targeting the promoter of *znuABC*, *znuABC2*, or an intragenic region (*znuint*). Shown are the results of at least three independent experiments in duplicate; statistical analysis of relative fold enrichment from promoter regions compared to the intragenic control was performed using Student's *t* test (unpaired). ***, *P* < 0.0005.

(ARP), which has been shown to be strongly induced upon zinc starvation (20), in MSΔ1. Interestingly, *znuA2* and *rpmG* expression was significantly induced compared to that of the wild type, while induction was abolished in the complemented strain MSΔ1::A. Induced expression of *znuA2* and *rpmG* indicates a lower concentration of free zinc in MSΔ1 (Fig. 1C) and hence suggests a role of ZnuABC in zinc import of MSMEG. However, this concentration change did not influence growth. In contrast, expression of *rpmG* and *znuC* (MSMEG_RS29150) was not increased in MSΔ2 (Fig. 1D), which indicated that cytosolic zinc concentration in MSΔ2 is maintained at wild-type levels. Together, the data suggested that ZnuABC^{MSMEG} is more important for zinc homeostasis than ZnuABC2^{MSMEG} or that ZnuABC2^{MSMEG} lacks a zinc import function.

This was surprising, since we showed previously that *znuABC* and *znuABC2* expression in MSMEG is zinc-dependently regulated by Zur (20). Therefore, we investigated the binding of Zur to the promoters of these operons. ChIP analyses with a MSΔ*zur* mutant strain (MSΔ6) which was complemented with hemagglutinin (HA)-tagged Zur (MSΔ6::L) revealed a significant enrichment of the promoter DNA of MSMEG *znuABC* and *znuABC2* in anti-HA pulldown samples but not of the intragenic control *znuint* (Fig. 2). In addition, treatment with the zinc-chelating agent TPEN [*N,N,N',N'*-tetrakis(2-pyridylmethyl)ethylenediamine] reduced and increase of zinc in the medium enhanced Zur binding, which is consistent with the general function of Zur as a repressor in zinc-replete conditions. These data clearly demonstrated a direct and zinc-dependent binding of Zur to the promoters of both ZnuABC transporters in MSMEG.

Both MSMEG ZnuABC transporters are exchangeable. The above data showed that despite its zinc-dependent regulation by Zur, ZnuABC2^{MSMEG} deletion presumably did not change the cytosolic zinc concentration, leaving its role in zinc transport open. To better understand the role of ZnuABC2^{MSMEG} in zinc uptake, we deleted *znuABC2* in MSΔ1, resulting in MSΔ*znuABC*Δ*znuABC2* (MSΔΔ4) and complemented this mutant either with only one transporter (MSΔΔ4::A or MSΔΔ4::B) or with both transporters (MSΔΔ4::AB). Growth of MSΔΔ4 was significantly impaired in standard liquid MB medium (see Materials and Methods) (Fig. 3A) and heavily impaired on standard MB agar. No growth was visible after 4 days; however, after 10 days, small colonies appeared (Fig. 3B, control). Interestingly, the growth defect on standard MB agar was nullified upon complementation with one of the ZnuABC transporters, ZnuABC^{MSMEG} or ZnuABC2^{MSMEG}, in a Spot-On assay, suggesting that each transporter can substitute for the other. Without complementation, the growth defect of the double mutant was specifically restored when the MB agar was supplemented with zinc in a concentration-dependent manner, whereas no growth was detectable with manganese, iron,

FIG 3 Growth experiments with MSΔΔ4 and complemented strains. (A) Strains MSwt and MSΔΔ4 as well as homologously complemented strain MSΔΔ4::A::B (*znuABC* and *znuABC2*) were grown in MB. Growth was monitored daily by OD₆₀₀ determination for 4 days. Shown are the results of three independent replicates in duplicate. Statistical analysis was performed using Student's *t* test. *, *P* < 0.05. (B) MSwt, MSΔΔ4, and homologously complemented strains MSΔΔ4::A (*znuABC*), MSΔΔ4::B (*znuABC2*), and MSΔΔ4::A::B were grown in MB, diluted in PBS to an OD₆₀₀ of 0.1, diluted 10-fold to 10⁻⁴, and plated on MB standard agar (control) and MB supplemented with 1 or 5 μM TPEN or 25 or 100 μM ZnSO₄, MnSO₄, CoCl₂, MgSO₄, or FeNH₃-citrate. Agar plates were incubated for 4 days at 37°C. The control was incubated for up to 10 days. Assays were repeated three times in duplicate. Shown are the results of the 10⁻¹ dilution of a representative replicate.

cobalt, or magnesium (Fig. 3B). These results clearly indicate the importance and specificity of both transporters for zinc uptake.

Predicted MAP zinc importers restore MSΔΔ4 growth-defective phenotype.

Having established MSΔΔ4, we used this readout system to investigate the MAP transporters MptABC, MAP3774-76, and ZnuABC^{MAP} with regard to their impact in zinc homeostasis.

For this, MSΔΔ4 was complemented with MptABC, MAP3774-76, and ZnuABC^{MAP}, resulting in strains MSΔΔ4::C, MSΔΔ4::D, and MSΔΔ4::E, respectively. To show whether the putative MAP zinc transporters were also able to restore the zinc-dependent growth defect of MSΔΔ4, we repeated the above-mentioned Spot-On assays on supplemented MB agars with these strains. As shown in Fig. 4A, only ZnuABC^{MAP} was able to fully restore the phenotype on standard MB agar after 4 days of incubation. However, complementation of MSΔΔ4 with MptABC and MAP3774-76 led to better growth when the MB agar was supplemented with zinc than was seen in MSΔΔ4, and this enhanced growth compared to the mutant was also visible after 10 days without supplement. Thus, all MAP transporters seem to contribute to zinc uptake in the double mutant, with ZnuABC^{MAP} being the most efficient.

To further emphasize the role of MSMEG and MAP transporters in zinc uptake, we performed growth experiments in metal-depleted minimal Sauton's medium (SM). In SM, the double mutant MSΔΔ4 was not able to grow (Fig. 4B). Complementation with MSMEG ZnuABC transporters and also with each of the MAP transporters completely

FIG 4 Growth of MSΔΔ4 with different putative MAP zinc transporters. (A) MSwt, MSΔΔ4, and heterologously complemented strains MSΔΔ4::C (MptABC), MSΔΔ4::D (MAP3774-76), and MSΔΔ4::E (ZnuABC^{MAP}) were grown in MB, diluted in PBS to an OD₆₀₀ of 0.1, diluted 10-fold up to 10⁻⁴, and plated on MB standard agar (control) and MB supplemented with 1 or 5 μM TPEN or 25 or 100 μM ZnSO₄, MnSO₄, CoCl₂, MgSO₄, or FeNH₃-citrate. Agar plates were incubated for 4 days at 37°C. The control was incubated for up to 10 days. Assays were repeated three times in duplicate. Shown are the results of the 10⁻¹ dilution of a representative replicate. (B to E) Growth of MSΔΔ4 in Sauton's medium (SM). Strains were grown in MB, washed and inoculated to an initial OD₆₀₀ of 0.1 into fresh SM. SM standard (B) was supplemented with (C) 10 μM TPEN or (D) 1 μM or (E) 10 μM ZnSO₄. Growth was monitored daily for 8 days by OD₆₀₀ measurement. Shown are the results of at least three independent replicates in duplicate.

restored growth to wild-type level (Fig. 4B). The restoration of MSΔΔ4 growth defect was also possible upon supplementation of SM with ZnSO₄ (Fig. 4D and E), which corresponds with our results from the Spot-On assay. Conversely, TPEN treatment leads to reduced growth of all strains (Fig. 4C).

FIG 5 FluoZin-3AM zinc uptake assay. MSMEG strains were grown in SM for 3 days, washed with PBS, adjusted to an OD_{600} of 2.0, and treated with 1 μ M FluoZin-3AM or left untreated (MSwt w/o FluoZin [control]) for 30 min. Samples of each strain were adjusted to an OD_{600} of 0.1 (approximately 1×10^7 bacteria) and subjected to a zinc uptake assay. $ZnSO_4$ (25 μ M) was added after 1 h of incubation (blue arrow) for 3 h; subsequently, 1 μ M TPEN was added for 1 h. Fluorescence (relative light units [RLU]) was measured at extinction/excitation wavelengths of 495/516 nm every 10 min. RLU was normalized to OD_{600} (input). Shown are the results of three experiments in triplicate (with standard errors of the means [SEM]). Statistical analysis was performed using one-way analysis of variance (ANOVA). *, $P < 0.05$.

MSMEG and MAP zinc importers contribute to zinc uptake. Overall, all growth experiments indicated an involvement of ZnuABC^{MSMEG}, ZnuABC2^{MSMEG}, MptABC, MAP3774-76, and ZnuABC^{MAP} in zinc transport. We subsequently performed a nonradioactive zinc uptake assay by use of the fluorochrome FluoZin-3AM (FluoZin-3 acetoxymethyl derivative) to directly illustrate zinc uptake by the different transporters according to a protocol by Bayle et al. (30) and Chimalapati et al. (31). In this assay, zinc uptake was indicated by an increasing fluorescence, as only binding of zinc to intracellular FluoZin-3 leads to fluorescence. Fluorescence was measured in three phases: a 1-h adaptation phase without zinc, 3 h with 25 μ M zinc for zinc uptake, and 1 h with 1 μ M TPEN to bind free zinc and stop zinc uptake. FluoZin-3AM assays showed a reduced uptake of zinc by MS $\Delta\Delta$ 4 compared to MSwt (Fig. 5). Complementation with all MSMEG and MAP zinc importers restored uptake to the wild-type level or above (Fig. 5), and zinc uptake was significantly higher in all MS $\Delta\Delta$ 4 strains complemented with MAP transporters than in the deletion mutant. Hence, this experiment provides strong evidence for zinc uptake by MSMEG transporters ZnuABC^{MSMEG}, ZnuABC2^{MSMEG}, and MAP transporters MptABC, MAP3774-76, and ZnuABC^{MAP}.

In silico analyses of mycobacterial zinc transporters. Having established the impact of all predicted MSMEG and MAP transporters in zinc uptake, we were interested in their prevalence in other bacterial species. Therefore, we analyzed these transporters in more detail *in silico* to identify common or conserved structural elements of mycobacterial zinc transporters. By such analyses, we were able to distinguish two types of ZnuABC transporters. They were termed “type 1” for the classical ZnuABC, which is present in numerous bacteria, and “type 2,” which was found to be present exclusively in actinobacteria. The transporter MptABC appeared to be specific to MAP.

ZnuABC type 1. We first analyzed MSMEG and MAP zinc transporters by NCBI blastp to identify homologous proteins/transport systems in other bacteria. ZnuABC^{MSMEG}, ZnuABC^{MAP}, and surprisingly MAP3774-76 were highly similar to ZnuABC type 1 of MVAN (MVAN5318-20, ZnuABC^{MVAN}) and ZnuABC type 1 of SCO (SCO2505-07, ZnuABC^{SCO}); similarity was indicated by high amino acid identity and query coverage (Table 1). This implies the presence of two ZnuABC type 1 importers in MAP. Interestingly, MAP3774-76 was much more similar to ZnuABC^{MVAN} (88 to 99% query coverage and 53 to 74% identity) than the annotated ZnuABC^{MAP} (83 to 91% query coverage and 34 to 42% identity). Further, structural similarities of ZnuABC type 1 transporters could be shown by comparison of MSMEG and MAP transporters with known ZnuABC transporters of SCO, MVAN, MTB, *E. coli* and *S. enterica* serovar

TABLE 1 NCBI blastp analysis of MSMEG and MAP putative zinc transporters against MVAN and SCO ZnuABC transporters^a

MSMEG			MVAN			SCO				
Transporter	Old locus tag	New locus tag ^b	Protein length (aa)	Locus tag	Protein length (aa)	% query coverage/ % identity ^c	Locus tag	Protein length (aa)	% query coverage/ % identity ^c	Putative function ^d
ZnuABC1	<i>msmeg_6045</i>	MSMEG_RS29145	290	<i>mvan_5318</i>	285	85/77.54	SCO0475	283	42/30.89	Metal ABC transporter permease ZnuB
	<i>msmeg_6046</i>	MSMEG_RS29150	259	<i>mvan_5319</i>	262	98/68.36	SCO2506	256	78/37.20	ABC transporter ATP-binding protein ZnuC
	<i>msmeg_6047</i>	MSMEG_RS29155	335	<i>mvan_5320</i>	321	91/64.91	SCO2505	328	96/22.51	Zinc ABC transporter substrate-binding protein ZnuA
ZnuABC2	<i>msmeg_6049</i>	MSMEG_RS29165	388	<i>mvan_5323</i>	397	97/67.10	SCO0472	402	99/49.74	Hypothetical protein
	<i>msmeg_6050</i>	MSMEG_RS29170	300	<i>mvan_5324</i>	297	98/63.18	SCO0473	329	91/59.01	Zinc ABC transporter substrate-binding protein ZnuA2
	<i>msmeg_6051</i>	MSMEG_RS29175	387	<i>mvan_5325</i>	663	97/44.16	SCO0474	415	91/36.68	Inner membrane permease protein ZnuB2
ZnuABC1	<i>msmeg_6052</i>	MSMEG_RS29180	274	<i>mvan_5325</i>	663	95/64.89	SCO0475	283	97/61.28	Inner membrane permease protein ZnuB2
	<i>map_0487</i>	MSMEG_RS29185	236	<i>mvan_5326</i>	218	83/60.10	SCO0476	252	84/48.76	Metal ABC transporter ATP-binding protein ZnuC2
	<i>map_0488</i>	MAP_RS02500	286	<i>mvan_5318</i>	285	91/38.02	SCO2506	256	79/36.65	Metal ABC transporter permease ZnuB
Other	<i>map_0489</i>	MAP_RS02495	273	<i>mvan_5319</i>	262	86/41.95	SCO2506	256	79/36.65	Metal ABC transporter ATP-binding protein ZnuC
	<i>map_3734</i>	MAP_RS19145	593	<i>mvan_5320</i>	321	83/34.35	SCO0473	329	45/30.67	Zinc ABC transporter substrate-binding protein ZnuA/A2
	<i>map_3735</i>	MAP_RS19150	468	<i>mvan_5007</i>	663	82/31.14				ABC transporter ATP-binding protein
ZnuABC1	<i>map_3736</i>	MAP_RS22990	319							
	<i>map_3774</i>	MAP_RS19345	285	<i>mvan_5318</i>	285	88/74.10				Metal ABC transporter permease ZnuB
	<i>map_3775</i>	MAP_RS19350	262	<i>mvan_5319</i>	262	96/62.20	SCO2506	256	80/34.58	ATP-binding cassette domain-containing protein ZnuC
	<i>map_3776</i>	MAP_RS19355	263	<i>mvan_5320</i>	321	99/53.42	SCO0473	329	38/26.50	Zinc ABC transporter substrate-binding protein ZnuA/A2

^aZnuABC type 1 transporter homologues of MSMEG and MAP are highlighted in gray, aa, amino acids.

^bDue to a reannotation of the MSMEG genome [NC_008596](#) by NCBI, new locus tags were introduced in this study. The ZnuABC^{MSMEG} type 1 transporter is newly annotated with MSMEG_RS29145-55. New locus tags of the reannotation of the MAP genome [NC_002944](#) were also added.

^cSimilarity of proteins is shown by percent length of the query and percent identical amino acid residues.

^dBased on NCBI blastx analysis.

Typhimurium by Phobius domain prediction. All ZnuA and ZnuC proteins exhibited noncytoplasmic domains and signal sequences, whereas ZnuB proteins featured eight transmembrane domains, which is characteristic of type 1 ABC transporter transmembrane permeases (32) (Fig. 6A). Structure analysis also confirmed the strong resemblance of MAP3774-76 with classical ZnuABC type 1 transporters (Fig. 6A), further supporting the hypothesis of two ZnuABC type 1 transporters in MAP. The ZnuABC transporter of MTB, however, was missing the ZnuC protein and some transmembrane domains in ZnuB. Following this, alignments of the single Znu proteins by Clustal Omega analyses showed high homologies of all analyzed ZnuA proteins. They all contained highly conserved histidine (H) residues for zinc binding; however, the histidine/arginine/glutamic acid (H/E/D) rich region, which is often found in zinc transporters (10), was not present in ZnuA of pathogenic mycobacterial species (Fig. 6C). ZnuB and ZnuC proteins were highly similar (except MTB), ZnuC proteins harbored classical Walker A+B and C motifs for ATP binding. However, the switch region was present only in ZnuC of *E. coli* and *S. Typhimurium* (see Fig. S2 in the supplemental material).

ZnuABC type 2. ZnuABC2^{MSMEG} was suggested to resemble ZnuABC2 of MVAN (20). The presence of a second ZnuABC transporter was also described for SCO (24). However, no general classification concerning ZnuABC2 transporters has been published. BLAST analyses confirmed high similarities of *msmeg_6049-51* to ZnuABC2 of SCO (SCO0472-76, ZnuABC2^{SCO}), with a query coverage of 84 to 99% and 36 to 61% amino acid identity, and to ZnuABC2 of MVAN (MVAN5323-26, ZnuABC2^{MVAN}), with a query coverage of 83 to 98% and 44 to 67% amino acid identity (Table 1). Reannotation of the genome NC_08596 by NCBI revealed a novel open reading frame (ORF), MSMEG_RS29180, between *msmeg_6049-51* (MSMEG_RS29165-75) and the adjacent *znuC2*-homologous gene *msmeg_6052* (MSMEG_RS29185) (Table 1). Since the genes MSMEG_RS29180-85 were presumably part of the ZnuABC2^{MSMEG} transporter operon, we analyzed operon organization in more detail. Identification of a continuous transcript spanning MSMEG_RS29165-80 by PCR from cDNA of MSMEG (Fig. S3) and in our previous transcriptome sequencing (RNA-Seq) analyses (20) (Fig. S4) confirmed operon organization. MSMEG_RS29185 (*znuC2*) shows a different orientation, suggesting transcription of *znuABC2* by a bidirectional promoter (Fig. 6B). Moreover, conserved Zur binding motifs were detected 17 bases upstream of the translation start site (TLS) of MSMEG_RS29180 (TCCTTTTGAAAACGATTATCATTA) and 10 bases upstream the TLS of *znuC2* (CATTAATGATAATCGTTTCAAAA) by MEME-FIMO analysis in our recent study (20) (Fig. 6B). These findings together imply that the gene locus *msmeg_6049-52*, now referred to as MSMEG_RS29165-85, is a bidirectionally transcribed ZnuABC type 2 transporter in MSMEG with more than the classical three proteins. However, the role of MSMEG_RS29165 remains to be elucidated.

Phobius domain prediction and protein alignments revealed a high degree of structural analogy of ZnuA2, ZnuB2, and ZnuC2 proteins of MSMEG, MVAN, and SCO to the corresponding proteins of all analyzed ZnuABC type 1 transporters. However, there were some remarkable differences. All ZnuA2 proteins lack the H/E/D-rich region, which was present in ZnuA of MSMEG and MVAN (Fig. 6C), and all ZnuB2 proteins possess an additional noncytoplasmic domain, encoded either by a separate gene (MSMEG, SCO) or by a prolonged gene together with the transmembrane domain (MVAN) (Fig. 6A). This noncytoplasmic domain is unique for ZnuABC type 2 transporters, and no homologies between noncytoplasmic domains of ZnuB2 and ZnuB proteins could be identified (Fig. S2). BLAST analyses confirmed that homologues of ZnuB2 are present only in actinobacteria, mainly the genera *Nocardia*, *Streptomyces*, and *Mycobacterium*. Interestingly, within the genus *Mycobacterium*, only fast-growing species of the recently proposed subgenus *Mycolicibacterium* (33), which comprises MSMEG and MVAN, contain ZnuB2 homologues (Fig. 7).

Others. MptABC did not resemble classical ZnuABC transporters and did not show homologies to other ZnuABC transporters or any structural similarities, except the presence of transmembrane domains.

Altogether, we found that while ZnuABC type 1 transporters are present in all

FIG 6 Structural analysis of ZnuABC transporters. (A) Protein prediction of transmembrane (gray peaks), cytoplasmic (green lines), and noncytoplasmic domains (blue lines) as well as signal sequences (red lines) by Phobius analysis of ZnuABC transporter proteins from MSMEG, SCO, MVAN, MAP, MTB (abbreviated Rv), *S. Typhimurium* (EIL) and *E. coli* (b). (B) Organization of *znuABC* transporter genes of MVAN (top), MSMEG (middle), and MAP (bottom). Genes of an operon are colored the same. Bars indicate predicted Zur binding sites; the colors indicate affiliations with operons. Green arrowheads indicate locations of gene-overlapping primers used in Fig. S2. (C) Clustal Omega alignment of ZnuA/ZnuA2 proteins of MSMEG, SCO, MVAN, MAP, MTB, *S. Typhimurium*, and *E. coli*. Highly conserved histidine (H) residues for zinc binding are highlighted in green; H/E/D loops are highlighted in yellow.

analyzed species, ZnuABC type 2 transporters are present only in actinobacteria and harbor a characteristic additional noncytoplasmic domain. Type 1 transporters in pathogenic mycobacteria and all type 2 transporters lacked classical H/E/D loops in the ZnuA2 protein. Only a reduced form of ZnuABC type 1 was found in MTB, with only three transmembrane domains, while MAP presumably possesses two type 1 ZnuABC transporters and additionally MptABC.

DISCUSSION

In the present study, we characterized zinc transporters in MSMEG and MAP and showed the relevance of ZnuABC^{MSMEG}, ZnuABC2^{MSMEG}, ZnuABC^{MAP}, MAP3774-76, and

C

b1857-EcoliZnuA	AAVVASLKPVGFIAIAIDGVTETEV-LLPDGASEHDYSLRPSDVKRLQNADLVVWVGPE	84
EIL76_09930-SaleZnuA	AAVVASLKPLGFIAIAIDGVTDTQV-LLPDGASEHDYSLRPSDVKRLQGGADLVVWVGPE	84
SCO2505-ZnuA	FDVVASFYPMELAEQIGGDHVNVT-LTEPGQEPHDLELSAKQTARMGEADAVLYLKSL	98
MVAN5320-ZnuA	VNVVVSVDQWGNIVSQLGGACAEVTTVLGSAVDPHDYEPAASDAADFEGAQLVVLNGGH	84
MSMEG6047-ZnuA	VNVVVSVDQWGDIVSELGGACAKVTTVLGSSVDPHDFEPSPSDAAKFEGAQLVVLNGGH	111
MAP0489-ZnuA	TAVVASTDVWGSVARAVAGGHVAVASILSGSDQDPHSYEASPSDAAAADAGLVVFNNGG	102
MAP3776-ZnuA	---MVSVQWGDIVSELGGACANVKTVLASSVDPHDIYEPSPADAADFMNAKLIVVNGAG	57
Rv2059-ZnuA	-PGERKWDVWGSVARAVTGGHVPVKSILTGAHADPHSYQASPADAAAIVDAELVIYNGGG	161
SCO0473-ZnuA2	PRIVVTTNITLGDITRQVVGDAQAEVTV-LMKPNADPHSFGLSAVQAAQLERADLVFNGLG	98
MVAN5324-ZnuA2	GEVIVTTNITLGDVVRNVVGDAAADVVRV-LMQPNADPHSFVSAQDAAMGRAGLIVYNGLG	82
MSMEG6050-ZnuA2	RQIVVTTTTILGDVVSEVVGDTADVVRV-MMKPNADPHSYGLSAADAAMARADLVYNGLG	86
	. : . . . : . *	
b1857-EcoliZnuA	MEAFMQKPVSKLPGAKQV-TIAQLE-DVKPLLMKSIHGDDDDHDHA---E--KSDED--	134
EIL76_09930-SaleZnuA	MEAFMEKSVRNIPDNKQV-TIAQLA-DVKPLLMKGADEDEDEHAHTGADEE--KGDVH--	138
SCO2505-ZnuA	QPAVDEA-IAQSDVTKI-DAASLT---K--LEDHGNVEHDHHDHGGEEHAEDEHEEHS	150
MVAN5320-ZnuA	YDEWAAKLAATSADAPVNVNAVELSGGHAEQGGQEHAGEDGHGHAEEGQEHAQEQQ----	139
MSMEG6047-ZnuA	YDEWAGKLARSSAPNAPVVAADAGAHEDDAHDH-----GHDEHGHHAHA-----	156
MAP0489-ZnuA	YDGWVDDVLAHHP-GVARVDAYALLPDD-----	129
MAP3776-ZnuA	YDSWASKLAGSSASGAPLVSAAAVTT-----	83
Rv2059-ZnuA	YDPWVDQVLGHP-GVQAVDAYSLLGAV-----	188
SCO0473-ZnuA2	LEENVLRHVDAAREAGVA-TFEVGR-AVDPLTF-----H-----SGDDGGP	137
MVAN5324-ZnuA2	LEETVIRNVESAAAEGVP-TLAVTD-HVDPIEF-----A-----EGDSA--	119
MSMEG6050-ZnuA2	LEESMQRHVDAADAAENGVR-TLPVGD-HVDPLQF-----S-----A-DEH--	122
b1857-EcoliZnuA	-HHHGDFNMHLWLSPEIARATAVAIHGKLV-LMPQSRAKLDANLKDFAEQLASTETQVG	192
EIL76_09930-SaleZnuA	-HHHGEYNMHLWLSPEIARATAVAIHEKLV-LMPQSRAKLDANLKDFAEQLAATDKQVG	196
SCO2505-ZnuA	EGGEHALDPHVWLDPVKYAEIAQGVASFEEK-ADPDHAGDYRKNAEALAKKLSGLDTAYK	209
MVAN5320-ZnuA	QELGDEGNPHVWYNPTAVTEFAEAVTAQLGK-LSPDAAGYFAERHAEEFAESMKPYDEVIA	198
MSMEG6047-ZnuA	---EEGVNPHAWYSPAAVTTVADEVTAQLGD-LAPEAKDYFAERRAAFTESMKPYNQLID	212
MAP0489-ZnuA	-G--RPRNEHVFFYHLGVAKAVAAVADRLLA-IDPGNAADYRRNAAAFGRDADAIAIEH	185
MAP3776-ZnuA	--TPDGANPHLWYLPASAVTAVADAVTQELSR-MEPPAAGYFSQRRQFTSATRLVYVNLIA	140
Rv2059-ZnuA	-GDDAPNEHVFFYDPNVAKAVAATIADRLAD-LDPSNSGNRYRANAEEFSRGADAIAIEH	246
SCO0473-ZnuA2	EGESGRPDPHFWTDPDRVRTAAGLITDRVSEHVTGVDHDAIRANAEEYDEQLADLTAFME	197
MVAN5324-ZnuA2	----GTSDPHFWTDPQRMIMAVDAIEEAVRE-IDGIDRDAVARNAESYRAKLRELDTSMT	174
MSMEG6050-ZnuA2	----GAADPHFWTDPQRMILAVDAITGRVAE-LPGIDRAVAANAARYRDELNNLSAHMT	177
	: * :	

FIG 6 Continued

MptABC in zinc uptake. From our *in silico* analyses, we conclude, that ZnuABC transporters can be grouped in classical type 1, present in numerous bacterial species (11–18), and type 2, which is found only in actinobacteria. The noncytoplasmic part of protein ZnuB2 was exclusively present in type 2 transporters, mainly in *Nocardia* spp., *Streptomyces* spp., and fast-growing mycobacteria of the recently proposed subgenus *Mycolicibacterium* (33). This subgenus was described as being ancestral and evolutionarily related to *Nocardia* species (34). Hence, ZnuABC type 2 transporters in *Mycolicibacterium* spp. might originate from this evolutionary ancestor. The lack of

FIG 7 Phylogenetic tree of ZnuB2 homologous proteins. Shown are the results of NCBI blastp analysis with MSMEG_RS29175 and subsequent processing of data by the neighbor-joining method (maximum sequence difference, 0.75), illustrated in iTOL (version 5.7). Mycobacteria are in red, *Nocardia* species are in blue, and *Streptomyces* species are in green.

ZnuABC type 2 transporters in pathogenic mycobacteria, e.g., MTB and MAP might primarily be a result of genome reduction upon adaptation to their ecological niche in the host. However, MAP, in contrast to MTB, seems to have acquired a second copy of ZnuABC type 1 transporter, MAP3774-76, which is located on the MAP-specific LSP15 (29) and much more closely related to ZnuABC type 1 of the nonpathogenic MVAN than the “original” ZnuABC^{MAP} (Table 1). Hence, MAP might have acquired MAP3774-76 by horizontal gene transfer from closely related species, as described for *Vibrio parahaemolyticus* (35). This assumption is supported by the location of MAP3774-76 on a MAP-specific genomic insertion region, which was acquired during the evolution of *M. avium* subspecies (29).

The transporter MAP3774-76, as well as MptABC, has been shown to be related to iron transport by other groups (25–27). In the present study, we clearly showed that all of the above-mentioned putative MAP zinc transporters contributed to zinc uptake in MSMEG when cultured in SM and on MB agar and in the FluoZin-3AM assay, which is in agreement with their zinc starvation-induced gene expression shown in our previous transcriptomic study (21). However, in contrast to ZnuABC^{MAP}, which in all experiments fully restored the growth defect of the MSMEG double-deletion mutant MSΔΔ4, the MAP transporters MptABC and MAP3774-76 showed a reduced complementary

effect. For example, in the Spot-On assay, they complemented the MS $\Delta\Delta$ 4 growth phenotype only in the presence of zinc or after 10 days incubation. Interestingly, in SM, growth of all complemented MAP MS $\Delta\Delta$ 4 strains was restored to wild-type level, even without zinc supplementation. It is conceivable that these transporters are less efficient or are cotransporters for different metals, such as the SloABCR Mn/Fe transporter of *Streptococcus mutans* (36). For example, MAP3774-76 is a predicted Mn/Zn transport system, and the adjacent Fur-like regulator encoded by *map_3773* is a Fe/Zn regulator (25). However, we could not show any relevance of these transporters in manganese or iron transport, and this issue needs further analysis. An additional explanation for the growth differences on MB agar and SM medium might be the accessibility of free zinc. Even though SM is metal depleted by chelation, traces of metal ions remain in the medium. In MB agar, the concentration of zinc should be around 7 μ M (37–39). However, a large amount of zinc might be bound by the bovine serum albumin, which is a component of the oleic acid-albumin-dextrose-catalase (OADC) supplement and which is known to strongly bind zinc and copper with a stoichiometry of 1:1 (40–42). Moreover, accessibility might be further restricted by the growth on the agar surface, while traces of zinc are freely available in the liquid SM.

Overall, MSMEG possesses two and MAP three transporters for zinc. This is an exceptional mechanism for maintenance of zinc homeostasis and allows quick and efficient adaptation to changing zinc concentrations. Environmental mycobacteria such as MSMEG encounter zinc concentrations in a very wide range, from 1 μ g/liter (water) to at least 100 mg/kg (soil) (43–45). However, the level of free zinc is low, as zinc is mainly part of different types of minerals and may eventually be released by physicochemical parameters such as changing pH or temperature (46, 47). The presence of three zinc transporters in MAP suggests that it might also be exposed to strongly changing or low zinc concentrations in the intestine, the entry site of MAP into the host and the main site of zinc absorption in ruminants (48, 49). A restriction might be caused by zinc-binding enzymes such as calprotectin, which are secreted by the host as a reaction upon infection and also upon colonization, as shown for *S. Typhimurium*, *Campylobacter jejuni*, and *Acinetobacter baumannii* (50–52). It is also conceivable that acquisition of additional zinc importers might have occurred to compensate for the loss of a functional mycobactin. This iron siderophore is not functional due to mutations in several *mbt* operon genes in MAP (53), and thus, iron acquisition for MAP appears to be more difficult than that for, e.g., MTB, which has a functional mycobactin. Hence, MAP might have acquired these particular zinc uptake systems to utilize zinc over iron. A substitution of missing metal ions by another “first-row” transition metal was described for *Borrelia burgdorferi*, which uses manganese instead of iron for protein structure, to overcome iron shortage in the host (54). Also, for *S. Typhimurium*, utilization of cobalt upon zinc limitation in a ZnuABC-deficient mutant was described (55). However, whether this substitution mechanisms can be used by MAP needs further research.

ZnuABC^{MSMEG} seems to be the predominant zinc transporter in MSMEG, as deletion of this transporter alone led to transcriptional changes of zinc starvation marker genes, including *znuABC2*. In contrast, the ZnuABC2^{MSMEG}-deficient mutant MS Δ 2 showed no phenotype. Probably expression of *znuABC* occurs earlier, i.e., when zinc is already limited in the cell but not below a threshold which induces expression of *znuABC2*. Hence, ZnuABC^{MSMEG} is presumably sufficient to compensate for the loss of *znuABC2* in MS Δ 2 when zinc is available in the growth medium, and ZnuABC2^{MSMEG} is not necessary to maintain zinc homeostasis under these conditions. ZnuABC2^{MSMEG} might function as a backup transporter which is brought into action, when zinc concentration drops below a certain concentration, e.g., in MS Δ 1 or upon TPEN treatment. Indeed, a graded expression of different zinc-regulated genes in response to zinc starvation was recently proposed for *Bacillus subtilis*. It was shown that Zur-regulated promoters exhibited different affinities to either partially zinc-saturated Zur dimers (Zur₂-Zn₃) and/or fully zinc-saturated Zur dimers (Zur₂-Zn₄) and that decreasing zinc availability led to a stepwise

TABLE 2 Bacterial strains and mutants used in this study

Strain	Description	Source or reference
<i>E. coli</i> 10- β	NEB 10- β competent <i>E. coli</i>	NEB
MSwt	<i>M. smegmatis</i> mc ² 155	66
MS Δ 1	Markerless <i>znuABC</i> (MSMEG_RS29145-55) deletion mutant of MSwt	This work
MS Δ 2	Markerless <i>znuABC2</i> (MSMEG_RS29165-75) deletion mutant of MSwt	This work
MS $\Delta\Delta$ 4	Markerless <i>znuABC</i> and <i>znuABC2</i> double mutant of MSwt	This work
MS Δ 6	Markerless <i>zur</i> (MSMEG_RS21680) deletion mutant of MSwt	21
MS Δ 1::A	MS Δ 1 complemented homologously with pMV306kan-MS <i>znuABC</i>	This work
MS Δ 2::B	MS Δ 2 complemented homologously with pMV306kan-MS <i>znuABC2</i>	This work
MS $\Delta\Delta$ 4::A	MS $\Delta\Delta$ 4 complemented homologously with pMV306kan-MS <i>znuABC</i>	This work
MS $\Delta\Delta$ 4::B	MS $\Delta\Delta$ 4 complemented homologously with pMV306kan-MS <i>znuABC2</i>	This work
MS $\Delta\Delta$ 4::A::B	MS $\Delta\Delta$ 4 complemented homologously with <i>znuABC</i> and <i>znuABC2</i>	This work
MS $\Delta\Delta$ 4::C	MS $\Delta\Delta$ 4 complemented heterologously with pMV306kan-MAP <i>mptABC</i>	This work
MS $\Delta\Delta$ 4::D	MS $\Delta\Delta$ 4 complemented heterologously with pMV306kan-MAP3774-76	This work
MS $\Delta\Delta$ 4::E	MS $\Delta\Delta$ 4 complemented heterologously with pMV306kan-MAP <i>znuABC</i>	This work
MS Δ 6::L	MS Δ 6 complemented homologously with pMV306hyg-MS <i>zur</i> -HA	20

derepression of the Zur regulon (22). Therefore, we assume that zinc limitation in MS Δ 2 under the chosen conditions might not have been sufficient to derepress the *znuABC* operon, while deletion of *znuABC* in MS Δ 1 led to more severe starvation and thus derepression of *znuABC2* expression.

In conclusion, our data clearly reveal the presence of different zinc importer types in mycobacteria. We showed that two types of ZnuABC zinc importers—ZnuABC^{MSMEG} and ZnuABC2^{MSMEG}—are necessary for zinc-dependent growth in MSMEG, while ZnuABC^{MSMEG} seems to be the main importer and ZnuABC2^{MSMEG} is presumably a backup transporter. Moreover, we showed that MAP possesses two ZnuABC type 1 transporters—MAP3774-76 and ZnuABC^{MAP}—and MptABC, which all contribute to zinc uptake in MSMEG, strengthening their role in MAP zinc homeostasis.

MATERIALS AND METHODS

Bacterial strains, growth conditions, and chemicals. All strains used are listed in Table 2, and plasmids and primers are listed in Tables S1 and S2, respectively, in the supplemental material. Chemicals were purchased from Carl Roth (Karlsruhe, Germany) or Sigma-Aldrich (Munich, Germany) if not stated otherwise.

E. coli 10 β was used for plasmid cloning and grown in Luria-Bertani (LB) broth (37°C and 200 rpm) or on LB agar plates (37°C). Competent cells were prepared as described by Sambrook and Russell (56).

M. smegmatis mc²155 wildtype (MSwt), mutant strains MS Δ 1, MS Δ 2, MS $\Delta\Delta$ 4, and MS Δ 6, and complemented mutant strains were grown in Difco 7H9 Middlebrook medium (Becton Dickinson, Franklin Lakes, NJ, USA) containing 10% OADC (0.06% oleic acid, 5% albumin, 2% dextrose, 0.085% NaCl, 0.003% catalase, 2.5% glycerol 0.05% MgSO₄, and 0.025% tyloxapol) (MB) or Sauton's medium [0.05% KH₂PO₄, 0.2% citric acid, 0.005% ammonium iron(III) citrate, 0.4% L-asparagine monohydrate, 6% glycerol, 0.025% tyloxapol; pH 7.4] (SM) at 37°C and 150 rpm or on LB agar or Difco Middlebrook 7H10 agar (Becton Dickinson, Franklin Lakes, NJ, USA) containing 10% OADC and 2.5% glycerol (37°C). SM was prepared in purified Milli-Q water in rinsed plasticware and treated with 1% Chelex overnight. Media were supplemented with kanamycin (broth, 20 μ g/ml; agar plate, 50 μ g/ml) or hygromycin (broth, 50 μ g/ml; agar plate, 100 μ g/ml) if necessary. Electrocompetent MSMEG cells were prepared as described by Parish and Stoker (57).

Growth experiments. For determination of growth defects in the mutant strains, growth curves were performed in MB and SM. In addition, a Spot-On assay on MB agar plates containing different supplements was conducted, to determine the influence of metal ions on bacterial growth. For growth curves in MB and for the Spot-On assays, cryostocks were used for inoculation and prepared as follows. Bacteria were grown to an OD₆₀₀ of approximately 2.0, harvested, and resuspended in the subsequently used medium containing 10% glycerol by vortexing with glass beads; clumps were removed by slow centrifugation (300 $\times$ g and 4°C for 5 min), and the bacteria were stored at -80°C. Cryostocks were thawed on ice prior to use. Main cultures for growth kinetics were inoculated at an optical density at 600 nm (OD₆₀₀) of 0.05 (MB), and growth was monitored daily for 4 days by OD₆₀₀ measurement. For growth kinetics in SM, precultures were grown in MB (supplemented with antibiotics if necessary) to an OD₆₀₀ of approximately 2.0, harvested, washed twice, and resuspended in SM. Main cultures in SM (without antibiotics) were inoculated at a starting OD₆₀₀ of 0.1 and supplemented with 1 μ M N,N,N',N'-tetraakis(2-pyridylmethyl)ethylenediamine (TPEN) or 1 or 10 μ M ZnSO₄ or left untreated (control). OD₆₀₀ was measured daily for 8 days. Spot-On assays were performed as described by Speer et al. (58). Bacteria

from cryostocks were inoculated in $1 \times$ phosphate-buffered saline (PBS; Difco, Detroit, MI, USA) to an OD_{600} of 0.1 (1×10^{-1}) and 10-fold serially diluted to 1×10^{-4} . A 5- μ l portion of each strain and each dilution was spotted on predried 20-ml MB agar plates without supplements or containing 1 or 5 μ M TPEN or 25 or 100 μ M $ZnSO_4$, $CoCl_2$, $MnSO_4$, $MgSO_4$, or $FeNH_3$ -citrate. Sealed agar plates were incubated at 37°C for up to 10 days; pictures were taken after 4 and 10 days.

Preparation of nucleic acids, cDNA synthesis, and qRT-PCR. Extraction of genomic DNA was performed as described by Stratmann et al. (59), with extraction of ChIP-DNA as described below. Plasmid DNA was prepared with a NucleoSpin plasmid extraction or NucleoBond AX kit (Macherey and Nagel GmbH, Düren, Germany) according to the manufacturer's protocol. Total RNA was prepared using a Direct-zol RNA miniprep kit according to the manufacturer's protocol (Zymo Research, Bath, UK). cDNA was synthesized and analyzed by quantitative real-time PCR (qRT-PCR) as described by Eckelt et al. (21), using *msmeg_3084* (*gapdh*, primers 1 and 2) as a housekeeping gene for normalization. Data are presented as $\Delta\Delta C_T$ values.

Generation and complementation of unmarked mutant strains. Mutant strains were generated by use of the p2NIL-pGOAL system published by Parish and Stoker (60) according to the protocol by Goethe et al. (20). In brief, 1.5-kb upstream (A) and downstream (B) flanking fragments of the deleted regions were amplified by PCR with Q5 high-fidelity polymerase (NEB, Frankfurt, Germany) from genomic DNA of MSwt using primers 3 to 6 for the *znuABC* (*msmeg_6045-47*, MSMEG_RS29145-55) deletion mutant MS Δ 1 and primers 7 to 10 for the *znuABC2* (*msmeg_6049-51*, MSMEG_29165-75) deletion mutant MS Δ 2. A and B fragments were then subcloned into pJET1.2 (Thermo Fisher Scientific, Waltham, MA, USA) and sequenced. Correctly sequenced plasmids were digested with HindIII/BciVI (*znuABC* A), BciVI/KpnI (*znuABC* B), HindIII/BbsI (*znuABC2* A), and BbsI/KpnI (*znuABC2* B), and related A/B fragments were ligated simultaneously to HindIII/KpnI-digested p2NIL, resulting in manipulation plasmids p2NIL-MS6045-47AB (*znuABC*) and p2NIL-MS6049-51AB (*znuABC2*). A marker gene cassette, obtained by PacI digestion from pGOAL19, was then ligated into the PacI-digested manipulation plasmids, resulting in deletion plasmids p2NIL-MS6045-47_Del and p2NIL-MS6049-51_Del.

Transformation of electrocompetent MSMEG cells with deletion plasmids and selection of mutants were performed as described before (20). The *znuABC* and *znuABC2* double-mutant MS $\Delta\Delta$ 4 was generated by transformation of MS Δ 1 with deletion plasmid p2NIL-MS6049-51_Del. Mutant strains were then complemented homologously with the deleted genes (MSMEG_RS29145-55, MSMEG_RS29165-75, or both), resulting in strains MS Δ 1::A, MS Δ 2::B, and MS $\Delta\Delta$ 4::A::B, respectively. MS $\Delta\Delta$ 4 was additionally complemented heterologously with the putative zinc transporter gene *map_3734-36* (MptABC), *map_3774-76*, or *map_0487-89* (ZnuABC^{MAP}) of MAP, yielding strains designated MS $\Delta\Delta$ 4::C, MS $\Delta\Delta$ 4::D, and MS $\Delta\Delta$ 4::E, respectively. For construction of complementation plasmids, the mentioned genes were amplified by standard PCR with Q5 high-fidelity polymerase (NEB, Frankfurt, Germany) from genomic DNA of MSwt or MAPwt with primers 11 to 20, as indicated in Tables S1 and S2. HindIII/XbaI-digested PCR products were ligated into HindIII/XbaI-digested pMV306hyg or pMV306kan. Sequenced plasmids were then transformed into the aforementioned mutant strains. Complementation was confirmed and polar effects were excluded by qRT-PCR using primers 21 to 30 (Fig. S1). Construction of the MSMEG Zur deletion mutant MS Δ 6 (*msmeg_4487*, MSMEG_RS21680) and complementation with HA-tagged Zur under the control of the promoter of *msmeg_4486*/MSMEG_RS21675 (MS Δ 6::L) were described previously (20).

Zinc uptake assay. A nonradioactive assay was established as described by Bayle et al. (30) and Chimalapati et al. (31) by use of FluoZin-3AM (Thermo Fisher Scientific, Waltham, MA, USA) to quantify zinc uptake in different MSMEG strains. Precultures of MSwt, MS $\Delta\Delta$ 4, MS $\Delta\Delta$ 4::A::B, MS $\Delta\Delta$ 4::C, MS $\Delta\Delta$ 4::D, and MS $\Delta\Delta$ 4::E were grown in MB to an OD_{600} of 2.0, washed twice, and resuspended in SM. Main cultures in 10 ml SM were inoculated with the preculture at an OD_{600} of 0.1 and incubated for 72 h in a shaking incubator at 37°C and 150 rpm to starve bacteria of zinc. Cells were then harvested, washed twice with 10 ml $1 \times$ PBS, resuspended in 1 to 2 ml $1 \times$ PBS (dependent on the pellet size), and adjusted to an OD_{600} of 2.0. For labeling, FluoZin-3AM was added to a final concentration of 1 μ M in dark reaction tubes to protect the FluoZin-3AM from light. To allow FluoZin-3AM uptake by the cells, samples were subsequently incubated for 30 min at room temperature on a rotator. Cells were then harvested by centrifugation, washed three times, and resuspended in 1 to 2 ml $1 \times$ PBS.

For complete deesterification of intracellular FluoZin-3 esters, cells were incubated for another 30 min at room temperature on a rotator and were subsequently diluted with $1 \times$ PBS to an OD_{600} of 0.1 (input). Two hundred microliters of the samples and $1 \times$ PBS as a blank were transferred to a black 96-well plate (in triplicate), and the samples were incubated at 37°C in a TECAN microtiter plate reader. Excitation and emission (495 and 516 nm) were measured every 10 min. First, a 1-h measurement was performed to obtain the baseline. Next, 25 μ M $ZnSO_4$ was added to each well, and measurement was performed for another 3 h to monitor zinc uptake. Then, 1 μ M TPEN was added, and measurement was performed for 1 h to show zinc specificity. Results were normalized to the negative control (PBS) and adjusted to the precise OD_{600} of the input.

ChIP and ChIP-qRT-PCR. ChIP and ChIP-qRT-PCR of MSwt and MS Δ 6::L were performed as described elsewhere (20). In brief, bacteria were grown in MB to an OD_{600} of approximately 2.0 and treated with 10 μ M TPEN and 500 μ M $ZnSO_4$ for 2 h or left untreated. After cross-linking with 1% formaldehyde for 20 min, cells were harvested and disrupted by treatment with a homogenizer and sonification. Two 250- μ g portions of protein from each sample were then incubated overnight (4°C) with A/G agarose matrix (Santa Cruz Biotechnology, Texas, USA) and either 2 μ g anti-HA antibody or 2 μ g anti-IgG1k antibody (all antibodies were from BioLegend, California, USA). Protein-DNA-A/G agarose complexes were then harvested (supernatant was retained as input control), beads were washed as described by Braunstein et al. (61), and DNA-protein complexes were eluted by heating and

centrifugation. Subsequently, protein and RNA were digested by proteinase K and RNase A treatment. DNA from ChIP and input samples was extracted by standard phenol-chloroform-isoamyl alcohol (25:24:1) extraction and precipitated with glycogen-LiCl-ethanol. DNA from ChIP and input was then subjected to qRT-PCR with primers targeting the promoter regions of *znuABC* (primers 31 and 32), *znuABC2* (primers 33 and 34), and *zitAint* (primers 35 and 36) as an intragenic control. Expression of samples from ChIP with anti-HA and isotype control samples was first normalized to the input, and the relative fold enrichment was calculated from normalized ChIP samples treated with anti-HA compared to normalized ChIP samples treated with the isotype control as described earlier (20).

Bioinformatics, statistics, and sequence analysis. National Center for Biotechnology Information (NCBI) protein BLAST (blastp suite) was performed with protein sequences of MSMEG (NCBI [NC_08596](#)) and MAP ([NC_002944](#)) against MVAN (NCBI BLAST TaxID 110539) and SCO (NCBI BLAST TaxID 1902) (Table 1), with the following settings: database nucleotide collection (nr/nt), models (XM/XP), and uncultured/environmental sample sequences excluded. For MSMEG_RS29075, blastp was performed with the above-mentioned settings for up to 5,000 sequences. Hits (3,027 sequences) were filtered (query coverage, 50% to 100%; protein identity, 40% to 100%; 127 sequences), and one strain of each species was selected for further analysis (45 species). A “distance tree of results” was generated in NCBI (tree method, neighbor joining; maximum sequence difference, 0.75), and the homologous sequences were downloaded and illustrated in iTOL (version 5.7) (62).

Protein sequences of MSMEG, MAP, MVAN ([NC_008726](#)), *S. enterica* subsp. *enterica* serovar Typhimurium ([CP034230](#)), *E. coli* strain K-12 substrain M1655 ([NC_000913](#)), and SCO ([AL_645882](#)) listed in Table S3 were subjected to analysis with the Phobius prediction program (<https://www.ebi.ac.uk/Tools/pfa/phobius/>) for domain analyses (63). Alignments of the above-mentioned proteins were performed using Clustal Omega (<https://www.ebi.ac.uk/Tools/msa/clustalo/>) (64). Statistics were calculated with GraphPad Prism 8.0.1 (version 244) (65).

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 0.6 MB.

ACKNOWLEDGMENTS

This work was supported by a grant from the German Research Foundation (DFG) to RG (Go983/4-1).

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

E.G. and R.G. conceived and planned the experiments. E.G., K.L., and A.G. constructed plasmids, mutants, and complemented strains and performed growth experiments and qRT-PCR. A.G. and E.G. performed FluoZin-3AM assays and Spot-On assays. J.L. performed ChIP assays. E.G. performed *in silico* analyses. E.G. and R.G. wrote the paper.

We declare that we have no competing interests.

REFERENCES

- Kehl-Fie TE, Skaar EP. 2010. Nutritional immunity beyond iron: a role for manganese and zinc. *Curr Opin Chem Biol* 14:218–224. <https://doi.org/10.1016/j.cbpa.2009.11.008>.
- Oутten CE, O'Halloran TV. 2001. Femtomolar sensitivity of metalloregulatory proteins controlling zinc homeostasis. *Science* 292:2488–2492. <https://doi.org/10.1126/science.1060331>.
- Andreini C, Banci L, Bertini I, Rosato A. 2006. Zinc through the three domains of life. *J Proteome Res* 5:3173–3178. <https://doi.org/10.1021/pr0603699>.
- Mertens J, Smolders E. 2013. Zinc, p 465–493. In Alloway BJ (ed), *Heavy metals in soils: trace metals and metalloids in soils and their bioavailability*, 3rd ed, vol 22. Springer, Dordrecht, The Netherlands.
- Babich H, Stotzky G. 1978. Toxicity of zinc to fungi, bacteria, and coliphages: influence of chloride ions. *Appl Environ Microbiol* 36:906–914. <https://doi.org/10.1128/AEM.36.6.906-914.1978>.
- McDevitt CA, Ogunniyi AD, Valkov E, Lawrence MC, Kobe B, McEwan AG, Paton JC. 2011. A molecular mechanism for bacterial susceptibility to zinc. *PLoS Pathog* 7:e1002357. <https://doi.org/10.1371/journal.ppat.1002357>.
- Soldati T, Neyrolles O. 2012. Mycobacteria and the intraphagosomal environment: take it with a pinch of salt(s)! *Traffic* 13:1042–1052. <https://doi.org/10.1111/j.1600-0854.2012.01358.x>.
- Lonergan ZR, Skaar EP. 2019. Nutrient zinc at the host-pathogen interface. *Trends Biochem Sci* 44:1041–1056. <https://doi.org/10.1016/j.tibs.2019.06.010>.
- Chandrangsu P, Rensing C, Helmann JD. 2017. Metal homeostasis and resistance in bacteria. *Nat Rev Microbiol* 15:338–350. <https://doi.org/10.1038/nrmicro.2017.15>.
- Hantke K. 2005. Bacterial zinc uptake and regulators. *Curr Opin Microbiol* 8:196–202. <https://doi.org/10.1016/j.mib.2005.02.001>.
- Gabbianelli R, Scotti R, Ammendola S, Petrarca P, Nicolini L, Battistoni A. 2011. Role of ZnuABC and ZinT in *Escherichia coli* O157:H7 zinc acquisition and interaction with epithelial cells. *BMC Microbiol* 11:36. <https://doi.org/10.1186/1471-2180-11-36>.
- Patzner SI, Hantke K. 1998. The ZnuABC high-affinity zinc uptake system and its regulator Zur in *Escherichia coli*. *Mol Microbiol* 28:1199–1210. <https://doi.org/10.1046/j.1365-2958.1998.00883.x>.
- Ammendola S, Pasquali P, Pistoia C, Petrucci P, Petrarca P, Rotilio G, Battistoni A. 2007. High-affinity Zn²⁺ uptake system ZnuABC is required for bacterial zinc homeostasis in intracellular environments and contributes to the virulence of *Salmonella enterica*. *Infect Immun* 75:5867–5876. <https://doi.org/10.1128/IAI.00559-07>.
- Bobrov AG, Kirillina O, Fosso MY, Fetherston JD, Miller MC, VanCleave TT, Burlison JA, Arnold WK, Lawrenz MB, Garneau-Tsodikova S, Perry RD. 2017. Zinc transporters YbtX and ZnuABC are required for the virulence of *Yersinia pestis* in bubonic and pneumonic plague in mice. *Metallomics* 9:757–772. <https://doi.org/10.1039/c7mt00126f>.
- Bobrov AG, Kirillina O, Fetherston JD, Miller MC, Burlison JA, Perry RD. 2014. The *Yersinia pestis* siderophore, yersiniabactin, and the ZnuABC system both contribute to zinc acquisition and the development of lethal

- septicaemic plague in mice. *Mol Microbiol* 93:759–775. <https://doi.org/10.1111/mmi.12693>.
16. Sheng Y, Fan F, Jensen O, Zhong Z, Kan B, Wang H, Zhu J. 2015. Dual zinc transporter systems in *Vibrio cholerae* promote competitive advantages over gut microbiome. *Infect Immun* 83:3902–3908. <https://doi.org/10.1128/IAI.00447-15>.
 17. Ellison ML, Farrow JM, Parrish W, Danell AS, Pesci EC. 2013. The transcriptional regulator Np20 is the zinc uptake regulator in *Pseudomonas aeruginosa*. *PLoS One* 8:e75389. <https://doi.org/10.1371/journal.pone.0075389>.
 18. Campoy S, Jara M, Busquets N, Perez De Rozas AM, Badiola I, Barbe J. 2002. Role of the high-affinity zinc uptake znuABC system in *Salmonella enterica* serovar Typhimurium virulence. *Infect Immun* 70:4721–4725. <https://doi.org/10.1128/iai.70.8.4721-4725.2002>.
 19. Maciag A, Dainese E, Rodriguez GM, Milano A, Provvedi R, Pasca MR, Smith I, Palu G, Riccardi G, Manganello R. 2007. Global analysis of the *Mycobacterium tuberculosis* Zur (FurB) regulon. *J Bacteriol* 189:730–740. <https://doi.org/10.1128/JB.01190-06>.
 20. Goethe E, Laarmann K, Luhrs J, Jarek M, Meens J, Lewin A, Goethe R. 2020. Critical role of Zur and SmtB in zinc homeostasis of *Mycobacterium smegmatis*. *mSystems* 5:e00880-19. <https://doi.org/10.1128/mSystems.00880-19>.
 21. Eckelt E, Jarek M, Fromke C, Meens J, Goethe R. 2014. Identification of a lineage specific zinc responsive genomic island in *Mycobacterium avium* ssp. *paratuberculosis*. *BMC Genomics* 15:1076. <https://doi.org/10.1186/1471-2164-15-1076>.
 22. Shin JH, Helmann JD. 2016. Molecular logic of the Zur-regulated zinc deprivation response in *Bacillus subtilis*. *Nat Commun* 7:12612. <https://doi.org/10.1038/ncomms12612>.
 23. Shin JH, Oh SY, Kim SJ, Roe JH. 2007. The zinc-responsive regulator Zur controls a zinc uptake system and some ribosomal proteins in *Streptomyces coelicolor* A3(2). *J Bacteriol* 189:4070–4077. <https://doi.org/10.1128/JB.01851-06>.
 24. Kallifidas D, Pascoe B, Owen GA, Strain-Damerell CM, Hong HJ, Paget MS. 2010. The zinc-responsive regulator Zur controls expression of the coelbactin gene cluster in *Streptomyces coelicolor*. *J Bacteriol* 192:608–611. <https://doi.org/10.1128/JB.01022-09>.
 25. Shoyama FM, Janetanakit T, Bannantine JP, Barletta RG, Sreevatsan S. 2020. Elucidating the regulon of a Fur-like protein in *Mycobacterium avium* subsp. *paratuberculosis* (MAP). *Front Microbiol* 11:598. <https://doi.org/10.3389/fmicb.2020.00598>.
 26. Wang J, Moolji J, Dufort A, Staffa A, Domenech P, Reed MB, Behr MA. 2015. Iron acquisition in *Mycobacterium avium* subsp. *paratuberculosis*. *J Bacteriol* 198:857–866. <https://doi.org/10.1128/JB.00922-15>.
 27. Lamont EA, O'Grady SM, Davis WC, Eckstein T, Sreevatsan S. 2012. Infection with *Mycobacterium avium* subsp. *paratuberculosis* results in rapid interleukin-1 β release and macrophage transepithelial migration. *Infect Immun* 80:3225–3235. <https://doi.org/10.1128/IAI.06322-11>.
 28. Riccardi G, Milano A, Pasca MR, Nies DH. 2008. Genomic analysis of zinc homeostasis in *Mycobacterium tuberculosis*. *FEMS Microbiol Lett* 287:1–7. <https://doi.org/10.1111/j.1574-6968.2008.01320.x>.
 29. Alexander DC, Turenne CY, Behr MA. 2009. Insertion and deletion events that define the pathogen *Mycobacterium avium* subsp. *paratuberculosis*. *J Bacteriol* 191:1018–1025. <https://doi.org/10.1128/JB.01340-08>.
 30. Bayle L, Chimalapati S, Schoehn G, Brown J, Vernet T, Durmort C. 2011. Zinc uptake by *Streptococcus pneumoniae* depends on both AdcA and AdcAll and is essential for normal bacterial morphology and virulence. *Mol Microbiol* 82:904–916. <https://doi.org/10.1111/j.1365-2958.2011.07862.x>.
 31. Chimalapati S, Cohen JM, Camberlein E, MacDonald N, Durmort C, Vernet T, Hermans PW, Mitchell T, Brown JS. 2012. Effects of deletion of the *Streptococcus pneumoniae* lipoprotein diacylglycerol transferase gene lgt on ABC transporter function and on growth in vivo. *PLoS One* 7:e41393. <https://doi.org/10.1371/journal.pone.0041393>.
 32. Rees DC, Johnson E, Lewinson O. 2009. ABC transporters: the power to change. *Nat Rev Mol Cell Biol* 10:218–227. <https://doi.org/10.1038/nrm2646>.
 33. Gupta RS, Lo B, Son J. 2018. Phylogenomics and comparative genomic studies robustly support division of the genus *Mycobacterium* into an emended genus *Mycobacterium* and four novel genera. *Front Microbiol* 9:67. <https://doi.org/10.3389/fmicb.2018.00067>.
 34. Bachmann NL, Salamzade R, Manson AL, Whittington R, Sintchenko V, Earl AM, Marais BJ. 2019. Key transitions in the evolution of rapid and slow growing mycobacteria identified by comparative genomics. *Front Microbiol* 10:3019. <https://doi.org/10.3389/fmicb.2019.03019>.
 35. Liu M, Yan M, Liu L, Chen S. 2013. Characterization of a novel zinc transporter ZnuA acquired by *Vibrio parahaemolyticus* through horizontal gene transfer. *Front Cell Infect Microbiol* 3:61. <https://doi.org/10.3389/fcimb.2013.00061>.
 36. Paik S, Brown A, Munro CL, Cornelissen CN, Kitten T. 2003. The sloABCR operon of *Streptococcus mutans* encodes an Mn and Fe transport system required for endocarditis virulence and its Mn-dependent repressor. *J Bacteriol* 185:5967–5975. <https://doi.org/10.1128/jb.185.20.5967-5975.2003>.
 37. Middlebrook G. 1955. Fitzsimons Army Hospital report no. 1. Fitzsimons Army Hospital, Denver, Colorado.
 38. Middlebrook G, Cohn ML. 1958. Bacteriology of tuberculosis: laboratory methods. *Am J Public Health Nations Health* 48:844–853. <https://doi.org/10.2105/ajph.48.7.844>.
 39. Middlebrook G, Cohn M, Schaefer W. 1954. Studies on isoniazid and tubercle bacilli. III. The isolation, drug-susceptibility, and catalase-testing of tubercle bacilli from isoniazid-treated patients. *Am Rev Tuberc* 70:852–872. <https://doi.org/10.1164/art.1954.70.5.852>.
 40. Ohyoshi E, Hamada Y, Nakata K, Kohata S. 1999. The interaction between human and bovine serum albumin and zinc studied by a competitive spectrophotometry. *J Inorg Biochem* 75:213–218. [https://doi.org/10.1016/s0162-0134\(99\)00090-2](https://doi.org/10.1016/s0162-0134(99)00090-2).
 41. Peters T, Jr, Blumenstock FA. 1967. Copper-binding properties of bovine serum albumin and its amino-terminal peptide fragment. *J Biol Chem* 242:1574–1578. [https://doi.org/10.1016/S0021-9258\(18\)96130-2](https://doi.org/10.1016/S0021-9258(18)96130-2).
 42. Singh N, Pagariya D, Jain S, Naik S, Kishore N. 2018. Interaction of copper (II) complexes by bovine serum albumin: spectroscopic and calorimetric insights. *J Biomol Struct Dyn* 36:2449–2462. <https://doi.org/10.1080/07391102.2017.1355848>.
 43. Merian E, Clarkson TW. 1991. Metals and their compounds in the environment. VCH, New York, NY.
 44. Van Sprang P, Verdonck F, Van Assche F, Regoli L, De Schamphelaere K. 2009. Environmental risk assessment of zinc in European freshwaters: a critical appraisal. *Sci Total Environ* 407:5373–5391. <https://doi.org/10.1016/j.scitotenv.2009.06.029>.
 45. Goodwin ID, Howard WR. 2012. Evidence of environmental change from the marine realm, p 181–210. In Matthews JA, Bartlein PJ, Briffa KR, Dawson AG, de Vernal A, Denham T, Fritz SC, Oldfield F (ed), *The SAGE handbook of environmental change*. SAGE Publications, Thousand Oaks, CA.
 46. Li H, Shi A, Li M, Zhang X. 2013. Effect of pH, temperature, dissolved oxygen, and flow rate of overlying water on heavy metals release from storm sewer sediments. *J Chem* 2013:434012. <https://doi.org/10.1155/2013/434012>.
 47. Sims JT. 1986. Soil pH effects on the distribution and plant availability of manganese, copper, and zinc. *Soil Science Soc Am J* 50:367–373. <https://doi.org/10.2136/sssaj1986.03615995005000020023x>.
 48. Hampton D, Miller W, Neathery M, Kincaid R, Blackmon D, Gentry R. 1976. Absorption of zinc from small and large intestine of calves. *J Dairy Sci* 59:1963–1966. [https://doi.org/10.3168/jds.S0022-0302\(76\)84469-4](https://doi.org/10.3168/jds.S0022-0302(76)84469-4).
 49. Flagstad T. 1976. Lethal trait A 46 in cattle. Intestinal zinc absorption. *Nord Vet Med* 28:160–169.
 50. Gielda LM, DiRita VJ. 2012. Zinc competition among the intestinal microbiota. *mBio* 3:e00171-12. <https://doi.org/10.1128/mBio.00171-12>.
 51. Moore JL, Becker KW, Nicklay JJ, Boyd KL, Skaar EP, Caprioli RM. 2014. Imaging mass spectrometry for assessing temporal proteomics: analysis of calprotectin in *Acinetobacter baumannii* pulmonary infection. *Proteomics* 14:820–828. <https://doi.org/10.1002/pmic.201300046>.
 52. Liu JZ, Jellbauer S, Poe AJ, Ton V, Pesciaroli M, Kehl-Fie TE, Restrepo NA, Hosking MP, Edwards RA, Battistoni A, Pasquali P, Lane TE, Chazin WJ, Vogl T, Roth J, Skaar EP, Raffatellu M. 2012. Zinc sequestration by the neutrophil protein calprotectin enhances *Salmonella* growth in the inflamed gut. *Cell Host Microbe* 11:227–239. <https://doi.org/10.1016/j.chom.2012.01.017>.
 53. Li L, Bannantine JP, Zhang Q, Amonsin A, May BJ, Alt D, Banerji N, Kanjilal S, Kapur V. 2005. The complete genome sequence of *Mycobacterium avium* subspecies *paratuberculosis*. *Proc Natl Acad Sci U S A* 102:12344–12349. <https://doi.org/10.1073/pnas.0505662102>.
 54. Posey JE, Gherardini FC. 2000. Lack of a role for iron in the Lyme disease pathogen. *Science* 288:1651–1653. <https://doi.org/10.1126/science.288.5471.1651>.

55. Ammendola S, Ciavardelli D, Consalvo A, Battistoni A. 2020. Cobalt can fully recover the phenotypes related to zinc deficiency in *Salmonella Typhimurium*. *Metallomics* 12:2021–2031. <https://doi.org/10.1039/d0mt00145g>.
56. Sambrook J, Russell DW. 2006. Transformation of *E. coli* by electroporation. *CSH Protoc* 2006:pdb.prot3933. <https://doi.org/10.1101/pdb.prot3933>.
57. Parish T, Stoker NG. 1998. Electroporation of mycobacteria. *Methods Mol Biol* 101:129–144.
58. Speer A, Rowland JL, Haeili M, Niederweis M, Wolschendorf F. 2013. Porins increase copper susceptibility of *Mycobacterium tuberculosis*. *J Bacteriol* 195:5133–5140. <https://doi.org/10.1128/JB.00763-13>.
59. Stratmann J, Strommenger B, Goethe R, Dohmann K, Gerlach GF, Stevenson K, Li LL, Zhang Q, Kapur V, Bull TJ. 2004. A 38-kilobase pathogenicity island specific for *Mycobacterium avium* subsp. *paratuberculosis* encodes cell surface proteins expressed in the host. *Infect Immun* 72:1265–1274. <https://doi.org/10.1128/iai.72.3.1265-1274.2004>.
60. Parish T, Stoker NG. 2000. Use of a flexible cassette method to generate a double unmarked *Mycobacterium tuberculosis* *tlyA* *plcABC* mutant by gene replacement. *Microbiology* 146:1969–1975. <https://doi.org/10.1099/00221287-146-8-1969>.
61. Braunstein M, Rose AB, Holmes SG, Allis CD, Broach JR. 1993. Transcriptional silencing in yeast is associated with reduced nucleosome acetylation. *Genes Dev* 7:592–604. <https://doi.org/10.1101/gad.7.4.592>.
62. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res* 47:W256–W259. <https://doi.org/10.1093/nar/gkz239>.
63. Kall L, Krogh A, Sonnhammer EL. 2004. A combined transmembrane topology and signal peptide prediction method. *J Mol Biol* 338:1027–1036. <https://doi.org/10.1016/j.jmb.2004.03.016>.
64. Goujon M, McWilliam H, Li W, Valentin F, Squizzato S, Paern J, Lopez R. 2010. A new bioinformatics analysis tools framework at EMBL–EBI. *Nucleic Acids Res* 38:W695–W699. <https://doi.org/10.1093/nar/gkq313>.
65. Swift ML. 1997. GraphPad Prism, data analysis, and scientific graphing. *J Chem Inf Comput Sci* 37:411–412. <https://doi.org/10.1021/ci960402j>.
66. Snapper SB, Melton RE, Mustafa S, Kieser T, Jacobs WR, Jr. 1990. Isolation and characterization of efficient plasmid transformation mutants of *Mycobacterium smegmatis*. *Mol Microbiol* 4:1911–1919. <https://doi.org/10.1111/j.1365-2958.1990.tb02040.x>.