Published in final edited form as:

Cancer Immunol Res. 2021 April ; 9(4): 470–485. doi:10.1158/2326-6066.CIR-20-0085.

Immunosuppressive Myeloid Cells Induce Nitric Oxide-dependent DNA Damage and p53 Pathway Activation in CD8⁺ T cells

Adam N. R. Cartwright^{1,2}, Shengbao Suo³, Soumya Badrinath^{1,2}, Sushil Kumar^{1,2}, Johannes Melms^{5,6}, Adrienne Luoma^{1,2}, Archis Bagati^{1,2}, Assieh Saadatpour³, Benjamin Izar^{5,6}, Guo-Cheng Yuan³, Kai W. Wucherpfennig^{1,2,4}

¹Department of Cancer Immunology and Virology, Dana Farber Cancer Institute, Boston, MA 02215, USA

²Department of Immunology, Harvard Medical School, Boston, MA 02115, USA

³Department of Pediatric Oncology, Dana Farber Cancer Institute, Boston, MA 02215, USA

⁴Department of Neurology, Brigham & Women's Hospital & Harvard Medical School, Boston, MA 02115, USA

⁵Columbia Center for Translational Immunology, New York NY, USA

⁶Columbia University Medical Center, Division of Hematology and Oncology, New York, NY, USA

Abstract

Tumor-infiltrating myeloid-derived suppressor cells (MDSCs) are associated with poor survival outcomes in many human cancers. MDSCs inhibit T cell-mediated tumor immunity in part because they strongly inhibit T-cell function. However, whether MDSCs inhibit early or later steps of T-cell activation is not well established. Here we showed that MDSCs inhibited proliferation and induced apoptosis of CD8⁺ T cells even in the presence of dendritic cells (DCs) presenting a high-affinity cognate peptide. This inhibitory effect was also observed with delayed addition of MDSCs to co-cultures, consistent with functional data showing that T cells expressed multiple early activation markers even in the presence of MDSCs. Single-cell RNA-seq analysis of CD8⁺ T cells demonstrated a p53 transcriptional signature in CD8⁺ T cells co-cultured with MDSCs and DCs. Confocal microscopy showed induction of DNA damage and nuclear accumulation of activated p53 protein in a substantial fraction of these T cells. DNA damage in T cells was dependent on the iNOS enzyme and subsequent nitric oxide release by MDSCs. Small molecule-mediated inhibition of iNOS or inactivation of the *Nos2* gene in MDSCs markedly diminished

Corresponding author: Kai W. Wucherpfennig, Dana-Farber Cancer Institute, Smith 736, 450 Brookline Avenue, Boston, MA 02215, phone: (617) 632-3086, kai_wucherpfennig@dfci.harvard.edu.

Author Contributions

A.N.R.C. and K.W.W. conceived the study and designed experiments, A.N.R.C. performed the majority of *in vitro* and all *in vivo* experiments, S.B. performed a subset of co-culture experiments, S.K. performed western blots, A.L. generated libraries for single-cell RNA-seq; S.S., A.S. and G.-C.Y. performed computational analysis of single-cell RNA-seq data, S.K., J.M., and A.B. provided scientific discussion and ideas, A.N.R.C. and K.W.W. wrote the paper with input from all authors.

Competing interests:

K.W.W. serves on the scientific advisory board of TCR2 Therapeutics, T-Scan Therapeutics, SQZ Biotech and Nextechinvest, and receives sponsored research funding from Novartis. He is a co-founder of Immunitas, a biotech company.

DNA damage in CD8⁺ T cells. DNA damage in CD8⁺ T cells was also observed in KPC pancreatic tumors but was reduced in tumors implanted into *Nos2*-deficient mice compared with wild-type mice. These data demonstrate that MDSCs do not block early steps of T-cell activation but rather induce DNA damage and p53 pathway activation in CD8⁺ T cells through an iNOS-dependent pathway.

Keywords

CD8⁺ T cells; Myeloid-derived suppressor cells; DNA damage; P53 activation; Cancer immunology

Introduction:

Microbial infections result in heightened production of myeloid cells that leave the bone marrow before they have fully matured, a process referred to as ‘emergency hematopoiesis’ (1,2). Human cancers also substantially alter hematopoiesis through release of key regulators of myelopoiesis (including GM-CSF, G-CSF, and M-CSF) and pro-inflammatory mediators (including IL-6, IL-1 β , and TNF α) (3,4). Such alterations in myelopoiesis result in accumulation of immature myeloid cells with a pathological activation state in the circulation, lymphoid organs, and tumors. These immature myeloid cells are collectively referred to as myeloid-derived suppressor cells (MDSCs), and the number of circulating MDSCs represents an independent indicator of poor survival outcomes in a number of human cancers (5–7). In both humans and mouse models, MDSCs include populations of monocytic (M-MDSCs) and polymorphonuclear (PMN-MDSCs) that can be distinguished based on morphological characteristics and cell surface markers (8). MDSCs potently inhibit T cell-mediated tumor immunity and circulating MDSCs negatively correlate with clinical responses to checkpoint blockade (9).

In vitro and *in vivo* studies have demonstrated that MDSCs potently inhibit T cell proliferation (10) through multiple mechanisms (Supplementary Table 1) (11–19). A number of studies have shown that the release of nitric oxide (NO) by MDSCs is an important contributor to inhibition of T-cell proliferation (20). NO is synthesized by inducible nitric oxide synthase (iNOS) in MDSCs and NO readily diffuses across cell membranes (10,20). The production of superoxide by NADPH oxidase (Nox2) has also been implicated in inhibition of effector T-cell function (21,22). NO can react with superoxide, resulting in the formation of peroxynitrite. It was previously proposed that peroxynitrite inhibits the first steps of T-cell activation through nitrosylation of TCR and CD8 molecules (26). Other mechanisms implicated in the inhibition of T-cell function by MDSCs include the depletion of essential amino acids (including L-arginine, L-tryptophan, and L-cystine), release of inhibitory cytokines (such as IL10), and expression of inhibitory surface ligands (including PD-L1) (17,27–30).

The present study was designed to further interrogate the molecular changes that occur in T cells interacting with MDSCs. Single-cell RNA-seq analysis demonstrated that T cells co-cultured with dendritic cells (DCs) and MDSCs initially acquired transcriptional features of an activated state. However, the presence of MDSCs induced a p53 transcriptional signature

and DNA damage in T cells at later timepoints, resulting in T cell apoptosis. DNA damage was prevented by inactivation of the *Nos2* gene in MDSCs, implicating production of NO by iNOS in this process. Furthermore, KPC pancreatic tumors from *Nos2*-deficient mice were infiltrated by a greater number of CD8⁺ T cells, and DNA damage was substantially reduced in tumor-infiltrating CD8⁺ T cells from *Nos2* deficient mice compared with those in wild-type mice.

Materials and Methods

Antibody and cell staining reagents

Flow cytometry: Hoechst 33342 (Thermo; #R37165); Vybrant CFDA SE Cell Tracer (Thermo; #V12883); CellTrace CFSE Cell Proliferation Kit (Thermo; #C34553); Zombie UV (BioLegend; #423108); Zombie Violet (BioLegend; #423114); TruStain FcX (BioLegend; anti-mouse CD16/CD32; Clone 93; #101320); Annexin V-Pacific Blue (Thermo; #A35122); CD8α-BV650 (BioLegend; Clone 53-6.7; #100741); CD8α-PE (BioLegend; Clone 53-6.7; #100708); CD69-PE (BioLegend; Clone H1.2F3; #104508); CD25-APC-Cy7 (BioLegend; Clone PC61; #102026); IFNγ-PE (BioLegend; Clone XMG1.2; #505808); TNFα-PE-Dazzle594 (BioLegend; Clone MP6-XT22; #506346); CD11b-APC (BioLegend; Clone M1/70; #101262); Gr-1-PE (BioLegend; Clone RB6-8C5; #108408); Ly6G-FITC (BioLegend; Clone 1A8; #127606); Ly6G-PE-Cy7 (BioLegend; Clone HK1.4; #128018); Ly6C-PerCP-Cy5.5 (BioLegend; Clone HK1.4; #128028); Ly6G-FITC (BioLegend; Clone 1A8; #127606); CD45-PerCP-Cy5.5 (BioLegend; Clone 30-F11; #103130); CD3-BV786 (BioLegend; Clone 17A2; 100232); CD3-BV421 (BioLegend; Clone 17A2; #100228); NK1.1-APC (BioLegend; Clone PK136; 108710); CD107a-BV421 (BioLegend; Clone 1D4B; #121618); Granzyme B-APC (Thermo; Clone NGZB; #17-8898-82).

Western Blotting: Western Blotting anti-mouse p53 (Rat IgG2b; Clone 19764; R&D Systems; #AF1355); anti-phospho-p53 (S15) (Abcam; Polyclonal; #ab1431); Goat anti-rat IgG-HRP (H+L) (Thermo; Polyclonal; #31470); Goat anti-rabbit IgG-HRP (Cell Signaling Technologies; Polyclonal; #7074).

Neutralization: Neutralization anti-mouse IL-10 (Bio X Cell; Clone JES5-2A5; #BE0049); anti-mouse PD1 (CD279) (Bio X Cell; Clone 29F.1A12; #BE0273).

Microscopy: Microscopy anti-γH2AX-AF488 (Abcam; Clone EP854(2)Y; #ab195188); anti-phospho-p53 (S15) (Abcam; Polyclonal; #ab1431); anti-rabbit IgG-AF488 (Abcam; Polyclonal; #ab150077).

Small molecule inhibitors and co-culture supplements

N-acetyl-L-cysteine (NAC; Sigma; #A7250) was prepared in PBS to a working concentration of 1.25 mM. NG-monomethyl-L-arginine (L-NMMA; Sigma; #M7033) was prepared in PBS to a working concentration of 3.125 mM. Inhibitors were added to cultures prior to seeding of cells to final concentrations as indicated in the Results section. L-Arginine (Sigma; #A5006) stock solution was prepared at 5g/L (5X stock) in R10 medium

[RPMI-1640 (Thermo; #11875119), 10% fetal bovine serum (Thermo; #16000044), 2 mM L-glutamine (GlutaMAX; Thermo; #35050061), 100 µg/mL streptomycin, and 100 µg/mL penicillin (Penicillin-Streptomycin; Thermo; #15070063)], and the pH was adjusted to 7.0. L-Arginine was serially diluted in R10 medium and added to co-culture experiments at final concentrations of 0.2g – 1.0g/L.

Mice

C57BL/6 (000664-BJ; B6), C57BL/6-Tg (TcraTcrb)1100Mjb/J (003831; OT-I) and B6.129P2-Nos2^{tm1Lau}/J (002609; *Nos2*^{-/-}) were purchased from Jackson Laboratories. Experiments in mice were performed in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) at DFCI.

Isolation of cell populations for co-culture experiments

CD8⁺ T cells: CD8⁺ T cells were isolated from the spleens of OT-I mice by magnetic negative selection (Murine CD8⁺ negative selection kit; StemCell; #19853). All experiments (except where stated) were performed in R10 medium.

Bone marrow–derived DCs: Bone marrow cells (BMCs) were flushed from the tibias and fibulas of C57BL/6 mice using sterile PBS. Red blood cells were lysed (1 minute at RT) using Red Cell Lysis Buffer (Thermo; #A1049201) and washed with R10 medium and resuspended at 2×10^7 cells/mL. During washing, 10 mL R10 medium supplemented with 10ng/mL GM-CSF (BioLegend; 576306) was added to 10 cm Petri dishes. Once BMCs were resuspended, 5×10^6 cells were added per plate (dropwise into the center of the dish), incubated at RT for 15 minutes to allow to settle, then incubated at 37°C 5% CO₂ for 3 days. On day 3, 10 mL of GM-CSF supplemented pre-warmed R10 medium was gently added followed by culture for another 3 days. On day 6, 90% of the medium was aspirated, replaced with R10 medium supplemented with 10 ng/mL GM-CSF, 1 µg/mL LPS (*Escherichia coli* O111:B4, Sigma Aldrich; #LPS25), and 1nM SIINFEKL peptide (Invivogen; #vac-sin), and BM-DCs were then incubated overnight. On day 7, BM-DCs were gently lifted using a cell scraper, washed ($300 \times g$, 5 mins), and resuspended at a final density of 1×10^5 cells/mL in R10 medium.

Bone marrow–derived MDSCs: BMCs were flushed from the tibias and fibulas of C57BL/6 mice using sterile PBS. Red blood cells were lysed (1 minute at RT) using Red Cell Lysis Buffer, washed with R10 medium, then resuspended at 2×10^7 cells/mL. During washing, 10 mL R10 medium supplemented with 20ng/mL GM-CSF and IL-6 (BioLegend; #575706) was added to 10 cm tissue culture plates. Once BMCs were resuspended, 1×10^7 cells were added per plate (dropwise into the center of the dish then gently swirled around the plate) and incubated at 37°C 5% CO₂ for 3 days. On day 3, 10 mL of GM-CSF/IL-6 supplemented R10 medium was added and incubated overnight. On day 4, BM-MDSCs were gently lifted using a cell scraper, washed ($300 \times g$, 5 mins), and resuspended at a final density of 1×10^6 cells/mL. Typical composition of cultures was ~75% MDSC (of total viable) comprising ~60% PMN-MDSCs and ~30% M-MDSC (% of total MDSCs). For experiments using specific MDSC subsets, cells were stained using anti-CD11b-APC, anti-Gr-1-PE, anti-Ly6G-PE-Cy7, and anti-Ly6C-FITC. Defined cell populations were isolated

by FACS as described in the 'Flow Cytometry and Fluorescence-Activated Cell Sorting (FACS)' section.

Isolation of myeloid-derived suppressor cells from tumors: KPC tumors were cultured, implanted, and harvested as described in the 'Tumor cell lines and tumor models' section. After preparation of a single cell suspension, samples were stained using a live/dead marker (1:1000; Zombie Violet) and anti-murine CD16/32 (TruStain FcX; 1:100) as an Fc receptor block for 15 minutes at 4°C. Samples were then washed with FACS buffer and stained with anti-CD45-PerCP-Cy5.5, anti-CD11b-APC, and anti-Gr-1-PE (all 1:100; 30 mins at 4°C). After staining, cells were washed using FACS buffer and resuspended in FACS buffer supplemented with 0.5 mM EDTA (final concentration) and 100 µg/mL DNase I (Sigma; #11284932001). Cells were then sorted into R10 medium supplemented with 200 ng/mL GM-CSF and IL-6, washed using R10 medium ($300 \times g$, 5 mins), and resuspended at 1×10^6 cells/mL for co-culture assays. Sorted cells were assessed by flow cytometry to confirm purity of >90% for CD11b⁺Gr-1⁺ cells.

In vitro co-culture assays

MDSC suppression co-culture assay: *In vitro* co-culture assays were performed using 96-well U-bottom plates. OT-I T cells, SIINFEKL-pulsed DCs and MDSCs were co-cultured at a ratio of 1 (T cells) to 0.2 (DCs) to 2 (MDSCs), unless otherwise stated. DCs, OT-I T cells, and MDSCs were resuspended to 1×10^6 , 1×10^5 , and 1×10^6 cells/mL respectively and 100, 50, and 100 µL of each cell type was plated in respective order, resulting in a total of 5×10^4 , 1×10^4 , and 1×10^5 cells/well respectively. In DC only conditions, 100 µL R10 medium was added to maintain a total well volume of 250 µL. Where inhibitors/treatments were used, working concentrations (detailed in 'Small molecule inhibitors' section) or solvent controls were added to the wells prior to plating of cells. After addition of cells, co-cultures were mixed gently using a multi-channel pipette, centrifuged at $300 \times g$ for 30 seconds and then incubated for indicated times at 37°C 5% CO₂.

T-cell proliferation was assessed by labeling of CD8⁺ T cells with CFSE (CellTrace); equal volumes of CFSE (2 µM in PBS) and 1×10^7 CD8⁺ T cells/mL in PBS were mixed, the reaction was incubated for 1 min in a 37°C water bath, and cells were then washed twice with TCM. CFSE dilution by T cells was analyzed by flow cytometry at different time points (24 – 96 hours, as noted in the respective figure legends).

MDSC spike-in assay: MDSC cultures were prepared on day 4 (d4), day 3, and day 2 before co-culture so that d4 MDSCs could be added to co-cultures at different time points. Co-culture assays were prepared as described in the 'MDSC suppression co-culture assay' section, where conditions with MDSC spike-in at 24 or 48 hours were given 100 µL R10 in place of MDSCs. On the day of MDSC addition, 100 µL of supernatant was gently aspirated and MDSCs or R10 medium (to compensate for supernatant nutrient changes) were gently pipetted into the relevant wells. All co-cultures were then harvested 96 hours from the start of initial co-cultures for flow cytometric analysis.

Cell co-culture assay: Cells for co-cultures were prepared as described in ‘MDSC suppression co-culture assay’ section for ‘DC stimulated condition’. For ‘Cytokine stimulated condition’, OT-I CD8⁺ T cells and MDSCs were added to wells containing 100 µg/mL IL-15 (BioLegend; #566304) and 5 µg/mL IL-7 (BioLegend; #577806). Once plated, samples were mixed gently with a multi-channel pipette, centrifuged at 300 × *g* for 30 seconds and then incubated for the indicated times at 37°C 5% CO₂.

Cytokine stimulation of MDSCs and MDSC subsets

MDSCs were cultured as described in ‘Isolation of cell populations for co-culture experiments’ section and subsets were sorted as described in ‘Flow cytometry and Fluorescence-Activated Cell Sorting (FACS)’ section. After sorting, MDSCs and M-MDSCs were plated into 12-well plates (300,000 cells/well) and stimulated with either 1ng/mL IFN γ (BioLegend; 575306), 1ng/mL TNF α (BioLegend; #575206) or both cytokines (1ng/mL each) for 24 hours. Expression of *Nos2* was assessed by qRT-PCR as described in the ‘Quantitative RT-PCR’ section.

Apoptosis assays for CD8⁺ T cells

Apoptosis was assessed at the 72-hour time point for co-culture assays. Samples were first washed with PBS (300 × *g*, 5 mins) and then with Annexin Binding Buffer (Thermo; #V13246) (300 × *g*, 5 mins). After washing, cells were resuspended in 100 µL Annexin Binding Buffer containing Zombie UV (1:1000), Annexin V-Pacific Blue (1:250), and anti-CD8 α -BV650 (1:250). Samples were incubated at 37°C for 15 mins, followed by addition of 150 µL of Annexin Binding Buffer and analysis by flow cytometry. Early and late stages of apoptosis were determined using Zombie UV/Annexin V staining: early apoptosis (Zombie UV⁻Annexin V⁺) and late apoptosis (Zombie UV⁺Annexin V⁺).

Flow cytometry and Fluorescence-Activated Cell Sorting (FACS)

Flow cytometry: Flow cytometry analysis was performed using a BD LSR Fortessa instrument. All samples were washed (300 × *g*, 5 mins) and stained using FACS buffer (4% FBS in PBS). All samples were stained at an antibody dilution of 1:150 for 45 minutes at 4°C, then washed (300 × *g*, 5 mins) and resuspended in FACS buffer (4% FBS in PBS) for analysis/sorting. For intracellular staining, Cytofix/Cytoperm (BD) was used as per the manufacturer’s instructions.

FACS: Cell sorting was performed using a BD Aria III instrument. Cells were sorted into complete RPMI medium in 15 mL falcon tubes maintained at 4°C. Immediately after sorting, cells were pelleted (300 × *g*, 5 mins) and processed for downstream analysis as appropriate for each experiment. Cells were stained using a live/dead marker (1:1000; Zombie Violet, BioLegend) and anti-murine CD16/32 (1:100; BioLegend) for 15 mins at RT. Cells were then washed and stained using relevant antibodies for 30 mins at 4°C. After staining, cells were washed using FACS buffer and resuspended in FACS buffer supplemented with 0.5 mM EDTA and 0.1 mg/mL DNase I. All cells were sorted into R10 medium (supplemented as indicated). Once sorted, cells were washed with R10 medium (300 × *g*, 5 mins) and prepared for downstream applications as described in the relevant sections. All antibodies used are detailed in ‘Antibody Reagents’ section.

Quantitative RT-PCR

CD8⁺ T cells were sorted from co-cultures by FACS, then washed with PBS. RNA was isolated from CD8⁺ T cells using a Direct-zol RNA Miniprep Plus kit (Zymogen; #R2071) per the manufacturer's instructions and eluted into nuclease-free H₂O. RNA purity and concentration were determined using a DS-11 spectrophotometer (Denovix). For reverse transcription, 350 ng of RNA was used to transcribe cDNA using Superscript VILO (Thermo) per the manufacturer's instructions, then cDNA was diluted 5–10-fold in ddH₂O. The following assay probes (conjugated to 6-carboxyfluorescein) were purchased from Thermo: *Rps18* (housekeeping control; Mm02601777_g1), *Ccng1* (Mm00438084_m1), *Cdkn1a* (Mm04205640_g1), *Phlda3* (Mm00449846_m1), *Bax* (Mm00432051_m1), *Mdm2* (Mm01233138_m1) and *Nos2* (Mm00440502_m1). Quantitative RT-PCR reactions were prepared using Taqman Gene Expression Master Mix (Thermo; #4369610) per the manufacturer's instructions with a final volume of 10 µL per reaction using 4.5 µL diluted cDNA. All RT-PCR reactions were performed using technical triplicates on biological triplicates. RT-PCR was performed using a QuantStudio 6 (Thermo). Mean Ct values were then used to calculate Ct values to compare gene expression.

Nitrite quantification (Griess Reagent Reaction)

Nitrite quantification was performed using a Griess Reagent Kit (Thermo; #G7921) per manufacturer's instructions. After 72 hours of co-culture, plates were centrifuged (300 × g, 5 mins) and 200 µL supernatant was collected for this assay.

Tumor cell lines and tumor models

Tumor cell preparation and implantation: KPC tumor cells were a kind gift from Dr. Stephanie Dougan (Dana-Farber Cancer Institute). Cells were authenticated in 2017 by transcriptional sequencing. Cells were confirmed to be mycoplasma-free by both Charles River Laboratories Mouse Essential CLEAR panel prior to gifting and by the Universal Mycoplasma Detection Kit (Fisher Scientific; #50189644FP) prior to expansion and storage in liquid nitrogen. KPC cells were cultured in DMEM (Thermo; #11054001) supplemented with 10% fetal bovine serum, 100 µg/mL streptomycin, and 100 µg/mL penicillin. For each experiment, KPC cells were grown to a confluency of 60–80% and passaged two times prior to implantation. For passaging, cells were dissociated using TrypLE Express Enzyme (Thermo; #12604039) per the manufacturer's instructions. Cell dissociation was stopped by addition of KPC culture medium, and cells were then washed three times with sterile PBS. KPC cells were resuspended in PBS to a final cell density of 6×10⁶ cells/mL. Mice were shaved and anesthetized with isoflurane prior to implantation of KPC cells (3×10⁵ cells in 50 µL per mouse) into the subcutaneous layer of the skin along the left flank. Tumors were harvested after 21 days.

Tumor harvest and digestion.—Tumors were excised and placed into 10cm tissue culture dishes containing 10 mL of ice-cold RPMI-1640. Tumors were then diced using scalpels to <2 mm pieces and washed with RPMI-1640 (300 × g, 5 mins). Tumors were then resuspended in 9 mL of tumor dissociation solution [RPMI-1640 containing 1mg/mL collagenase type IV from *Clostridium histolyticum* (Sigma, #C5138), 20 U/mL DNase type

IV from bovine pancreas (Sigma; #D5025) and 0.1 mg/mL hyaluronidase type V from sheep testes (Sigma; #H6254)] and incubated at 37°C with shaking at 220 rpm for 45 mins. Digested tumors were then passed through a 70 µm cell strainer, washed with RPMI-1640 (300 × g, 5 mins), and cells were resuspended in 5 mL ACK lysis buffer followed by incubation at RT for 5 mins. Cells were then washed twice with PBS (300 × g, 5 mins) before downstream applications.

Confocal microscopy analysis of DNA damage and p53 activation in T cells

CD8⁺ T cells were isolated by FACS from co-cultures and plated on 0.01% poly-L-lysine (Sigma; #A-005-C) coated 8-well chamber LabTek slides (Nunc); cells were incubated at 4°C for 20 mins to allow them to adhere to the slides. Cells were then washed twice with PBS-Tween buffer [PBS supplemented with 0.05% Tween-20 (Sigma; #P1379); 200 µL per wash, removal by flicking], then fixed at 4°C for 20 mins in 200 µL Cytotfix/Cytoperm buffer (BD; #554722). Cells were washed twice with PBS-Tween buffer and then permeabilized using PBS containing 0.05% Triton-X-100 (Sigma; #X100) and 1% BSA (Sigma; #A1933) for 20 mins at RT. Following permeabilization, cells were washed twice with PBS supplemented with 1% BSA, 0.05% Tween-20, and slides were blocked for 30 mins at RT using 10% Normal Goat Serum (Thermo; #50062Z) supplemented with 5% FBS. Cells were then washed with PBS-Tween buffer and stained at 4°C overnight with anti-γH2AX-AF488 or anti-phospho-p53 (S15) diluted 1:200 in staining buffer (PBS containing 0.01% Tween 20 and 1% BSA). For detection of phospho-p53, cells were washed three times with PBS-Tween buffer, then stained with secondary Ab goat anti-rabbit IgG-AF488 at RT for 1 hour (1:200 in PBS containing 1% BSA 0.05% Tween-20). After staining, cells were washed twice with PBS and then stained using Hoechst-33342 (1:1,000 in PBS) for 5 mins at RT, followed by two washes with PBS-Tween. After washing, cells were covered with 200 µL of PBS for imaging.

Imaging was performed using a Leica SP5X confocal microscope with a 63x oil immersion lens (NA=1.4) and 2–3X digital zoom. Both γH2AX and phospho-p53 were imaged using an accumulation average of 4. Data were analyzed using ImageJ software. To determine mean intensity of fluorescence signal per cell, the integrated intensity of each cell was measured using the default threshold mask. For counting of γH2AX foci, a negative control sample of freshly isolated CD8⁺ T cells was stained and imaged as described above, and the maximum intensity value of negative samples was used to set a threshold minimum value. These thresholds were then applied to sample images and a threshold mask was applied to each image. Images were then converted to binary and particle measurement parameters which were set to >2 pixels and a circularity 0–1.00. Foci identified using these parameters were then recorded. For cells positive for γH2AX, cells with >1 foci were counted. For phospho-p53 positive cells, any cells with staining above background were counted.

Western blotting

FACS purified CD8⁺ T cells were lysed (30 µL per 1×10⁶ cells) in Pierce RIPA buffer (Thermo; #89901) supplemented with Halt Protease & Phosphatase Inhibitor Cocktail (Thermo; #78444) at a 1X final concentration. Lysates were incubated on ice for 30 mins, sonicated for 20 seconds, incubated on ice for a further 10 minutes, then centrifuged at

$>20,000 \times g$ for 15 minutes. Lysates were stored at -80°C prior to use and defrosted by centrifugation (4°C , $20,000 \times g$, 15 mins). Protein concentration was determined by BCA assay (Thermo; #A53225) and equal amounts of protein were prepared. Lysates were mixed with 4X SDS Loading Buffer (Thermo; #NP0007) supplemented with 60mM DMSO (Sigma; #472301) and heated at 95°C for 20 minutes. Samples were then pulsed in a centrifuge and loaded onto NuPAGE 8% Bis-Tris gels (Thermo; #WG1002BOX) which were run with 1X MOPS running buffer (Sigma; M1254) at 150V for 100 minutes. For molecular weight approximation, 4 μL of SeeBlue2 ladder (Thermo; LC5925) was loaded. Protein was then transferred onto PVDF membranes in Towbin buffer (25 mM Tris, 192 mM glycine, pH 8.3) at 100V for 100 minutes at 4°C . After transfer, membranes were blocked in 5% Blotting Grade Blocker (Bio-Rad; #1706404) for 30 minutes, washed with 1X TBS-T buffer, and probed using relevant antibody and blocking solution (indicated in antibody section) at 4°C overnight. Membranes were then washed extensively (at least 3 washes for 5–10 mins) with 1X TBS-T and then probed with relevant HRP-conjugated secondary antibody in 5% Blotting Grade Blocker for 1 hour at RT. Membranes were then washed extensively (at least 3 washes for 5–10 mins) with 1X TBS-T and exposed using $\sim 800 \mu\text{L}$ Western Lightning Plus-ECL (Perkin Elmer; #NEL105001EA) on a ChemiDoc MP Imaging System (Bio-Rad).

Single-cell RNA-sequencing (scRNA-seq)

Sample and library preparation: For scRNA-seq, naïve CD8^{+} T cells were isolated by negative selection as described in the ‘Cells and cell culture’ section; CD8^{+} T cells from co-culture assays were isolated by FACS at 24 and 72 hours as described in the ‘Flow cytometry and fluorescence-activated cell sorting (FACS)’ section. Immediately following isolation, cells were washed in PBS containing 0.04% RNase-free BSA. A total of 10,000 cells from each condition were targeted for 10X Genomics 3’ V2 single cell assay (10X Genomics). Reverse transcription, cDNA amplification and library preparation were performed per the manufacturer’s instruction. All libraries were sequenced using an Illumina HiSeq 2500 on rapid-run mode, yielding $\sim 31,000$ – $55,000$ reads per cell.

ScRNA-seq data analysis: De-multiplexing, alignment to the mm10 mouse genome and UMI counting were performed using Cell Ranger (version 2.0.1, 10X Genomics). This included read alignment and barcode demultiplexing, followed by UMI and barcode filtering and correction. The resulting gene expression matrix was then filtered to exclude genes with less than one UMI count in at least one cell. UMI counts were normalized by dividing the raw counts by the total counts in each cell. Subsequent clustering and visualization were conducted on a reduced expression matrix, retaining only the 1,000 genes exhibiting the highest dispersion values (defined as the variance over the mean expression). For each remaining gene, UMI counts were log-transformed and z-score normalized. To filter single cells with low quality, we adopted the following steps: (i) building a median-centered median absolute deviation (MAD)-variance normal distribution based on the fraction of mitochondrial genes of all single cells; removal of single cells with a mitochondrial fraction larger than the median of this distribution (31); (ii) removal of single cells with less than 200 detected genes; (iii) use of DoubletFinder (32) to filter potential doublets (set the doublet ratio as 7.5%). Following these steps, the data from different time points and conditions

were integrated and analyzed together; this included 4,050 cells from 0hr, 2,208 cells from 24hrs DC, 3,020 cells from 24hrs DC+MDSC, 2,260 cells from 72hrs DC, and 5,256 from 72hrs DC+MDSC cultures. Only genes that were expressed in more than 10 single cells were included. In order to focus on T cell populations, additional filtering was performed to obtain cells that had at least 5 UMI counts of CD3E, CD8A, or CD8B1, after which 13,629 single cells remained. The Seurat (33,34) package in R was then used for analyses. The data were then normalized and scaled using the “SCTransform” (35) function in Seurat and 2,000 highly variable genes were selected and top 30 principal components were used. Finally, shared nearest neighbor (SNN) was applied for clustering, and uniform manifold approximation and projection (UMAP) was used to visualize the single cell transcriptional profile in 2D space. Differentially expressed genes between different time points, conditions or clusters were identified by “FindAllMarkers” function (use default parameter but set min.pct as 0.3) in Seurat. The sub-clustering analysis was performed using the same strategy described above.

Data and materials availability

The GEO accession number for the scRNA-seq data reported in this paper is GSE145396.

Statistics

All statistical analyses were performed using Prism 8 (GraphPad). *P* value threshold was set at 0.05. All corrections for multiple comparisons are detailed in the figure legends.

Results:

MDSCs inhibit T-cell proliferation and survival.

This study was designed to investigate the mechanisms by which MDSCs interfere with T-cell function in the presence of DCs presenting a cognate peptide. MDSCs were either generated *in vitro* from bone marrow (Supplementary Fig. 1A) or isolated from KPC tumors (Supplementary Fig. 1B), a model of pancreatic cancer. We directly compared co-cultures of CD8⁺ T cells (OT-I T cells specific for Ova peptide) with DCs presenting the relevant high-affinity peptide (SIINFEKL) to cultures with DCs + MDSCs. ScRNA-seq was used to study the impact of MDSCs on the transcriptome of CD8⁺ T cells, and the identified pathways were further investigated by confocal microscopy and other techniques (Fig. 1A). Co-culture of OT-I CD8⁺ T cells with peptide-pulsed DCs induced potent T-cell activation, as evidenced by proliferation and expression of the activation markers CD25 and CD69 (Fig. 1B–E and Supplementary Fig. 1C–E). The addition of MDSCs to these cultures potently inhibited T-cell proliferation (Fig. 1B–C and Supplementary Fig. 1C), but these T cells expressed a high level of the CD25 and CD69 activation markers at 24 hours (Fig. 1B, D–E and Supplementary Fig. 1D–E), indicating that the MDSCs did not block essential early steps of T-cell activation in this system. In co-cultures with DCs + MDSCs, T-cell numbers were greatly reduced at the 72-hour time point (~25-fold) (Fig. 1F) and labeling with the cell death marker Zombie UV demonstrated substantially increased cell death (Supplementary Fig. 1F–G).

MDSCs include populations of PMN-MDSCs and M-MDSCs.

To determine the individual contribution of these subpopulations to inhibition of CD8⁺ T-cell proliferation and survival, we sorted PMN-MDSCs (Ly6G^{hi}Ly6C^{lo}) and M-MDSCs (Ly6C^{hi}Ly6G^{lo}) (Supplementary Fig. 1A, H) and examined their inhibitory activity. Given that the relative frequencies of PMN-MDSC to M-MDSCs in total MDSC cultures were 2:1, we adjusted the ratios of each subset accordingly for the relevant co-culture condition. We found that M-MDSCs potently inhibited CD8⁺ T-cell proliferation (Fig. 1G–J) and survival (Fig. 1K–L). M-MDSCs reduced T-cell numbers at 72 hours to a similar extent as the total MDSC population (Fig. 1G). Labeling with Annexin V and Zombie UV showed that M-MDSCs induced substantial apoptosis of CD8⁺ T cells in co-cultures (Fig. 1K–L). We also tested the suppressive capacity of MDSCs at lower T cell to MDSC ratios (Fig. 1M, N) and found that MDSC-mediated inhibition of CD8⁺ T-cell proliferation was directly correlated with MDSC numbers per culture. Similar findings were made with MDSCs directly isolated from KPC tumors (Supplementary Fig. 1B and Fig. 1O–R): CD8⁺ T-cell proliferation was greatly diminished in co-cultures with DCs + MDSCs (Fig. 1O–Q), and CD8⁺ T-cell numbers were diminished at 72 hours of co-culture (Fig. 1R). These data demonstrate that monocytic MDSCs potently inhibit T-cell proliferation and survival even in the presence of DCs presenting a high-affinity peptide.

MDSCs inhibit T-cell proliferation even following initial activation by DCs.

To investigate whether MDSCs could inhibit T-cell proliferation following initial activation of T cells by DCs, we set up co-culture experiments in which MDSCs were spiked into cultures of OT-I T cells and peptide-pulsed DCs after a delay of 24 or 48 hours. We found that T-cell proliferation and accumulation was attenuated even when MDSCs were added with a delay (Fig. 2A–C).

It has been proposed that MDSCs inhibit early signaling events downstream of the T-cell receptor (TCR) and co-receptors (26,36). We therefore tested whether MDSCs could inhibit CD8⁺ T-cell proliferation in response to the cytokines IL-2 and IL-15. MDSCs indeed inhibited cytokine-mediated T-cell proliferation and reduced T-cell numbers at 72 hours, although not to the same extent compared to co-cultures with peptide-pulsed DCs + MDSCs (Fig. 2D–F). These data demonstrate that MDSCs inhibit T-cell proliferation and accumulation even following initial activation by DCs.

ScRNA-seq analysis of transcriptional programs in CD8⁺ T cells modified by MDSCs.

We performed scRNA-seq analysis of CD8⁺ T cells sorted at 24 or 72 hours from co-cultures with DCs or DCs + MDSCs to study how MDSCs affect the transcriptional programs of CD8⁺ T cells. CD8⁺ T cells from 24-hour co-cultures with DCs and DCs + MDSCs tended to co-cluster in a global analysis of the scRNA-seq data (Fig. 3A; Supplementary Figure 2A); this 24-hour cluster was clearly distinct from the cluster representing input naïve CD8⁺ T cells. In contrast, at the 72-hour time point, CD8⁺ T cells from DCs versus DCs+ MDSCs were present in separate clusters.

Sub-clustering of CD8⁺ T cells from the 24-hour time point was performed to study potential differences of the transcriptional programs in T cells from DC versus DC + MDSC

co-cultures (Fig. 3B–D and Supplementary Fig. 2A). A total of seven sub-clusters were identified, and CD8⁺ T cells from DC + MDSC co-cultures were under-represented in two of these clusters (clusters 2 and 6). However, CD8⁺ T cells from cultures with DCs + MDSCs were well represented in other clusters, including cluster 0 (differential expression of cytotoxicity markers *Gzmb* and *Eomes* compared to other clusters), cluster 1 (differential expression of genes associated with DNA replication, including *Mki67* and *Cdk1*) and cluster 3 (differential expression of chemokine and cytokine genes, including *Ccl4*, *Il2*, and *Ifng*) (Fig. 3C–F). Consistent with these data, flow cytometric analysis showed a similar percentage of CD8⁺ T cells expressing granzyme B and IFN γ at the 24-hour but not the 72-hour time point in DC + MDSC co-cultures compared to DC co-cultures (Fig. 3G and Supplementary Fig. 2B–D). Gene set enrichment analysis (GSEA) of KEGG pathways demonstrated activation and effector related programs for all these clusters (Supplementary Fig. 2E). These results indicate that MDSCs did not inhibit all of the transcriptional changes associated with T-cell activation at this early time point.

MDSCs induce transcriptional activation of the p53 pathway in CD8⁺ T cells.

We next investigated gene expression programs in CD8⁺ T cells at the 72-hour time point. Comparison of highly variable genes highlighted transcriptional targets of p53 (*Dcxr*, *Ccng1*, *Rps27l*, *Phlda3*, *Bax*, *Ldhd*, *Cdkn1a*, *H2afj*, *Pmm1*, *Aen*, *Perp*, *Zmat3* and *Ei24*) that were more highly expressed in CD8⁺ T cells from DC + MDSC versus DC co-cultures (Fig. 4A, B). We therefore performed GSEA of highly variable genes expressed in T cells from DC + MDSC versus DC co-cultures at the 72-hour time point. This analysis again highlighted that p53 pathway genes were more highly expressed by T cells from DC + MDSC versus DC only co-cultures (Fig. 4C–D).

Sub-clustering of CD8⁺ T cells identified four clusters from the 72-hour time point, with clusters 2–4 representing CD8⁺ T cells from DC + MDSC co-cultures (Fig. 4E). GSEA showed enrichment of p53 pathway genes in all three T cell clusters from 72-hour DC + MDSC co-cultures (clusters 2–4); the p53 pathway ranked among the top two pathways for clusters 3 and 4 and was also identified for cluster 2 (Fig. 4F–G). T cells in clusters 3 and 4 may therefore represent cells with the highest level of transcriptional activation of the p53 pathway, which is consistent with the expression pattern of p53 pathway genes across T-cell clusters in UMAP plots (Fig. 4B). Using qPCR, we validated significantly increased expression of p53 pathway genes (*Ccng1*, *Cdkn1a*, *Phlda3*, *Bax*, and *Mdm2*) at the 72-hour timepoint in CD8⁺ T cells from DC + MDSC versus DC co-cultures (Fig. 4H). These data demonstrated that the presence of MDSCs induced transcriptional activation of the p53 pathway in CD8⁺ T cells.

Expression of iNOS by MDSCs is required for p53 pathway activation in CD8⁺ T cells.

We used genetic and pharmacological approaches to investigate the role of iNOS (*Nos2* gene) on activation of the p53 pathway in T cells. First, we compared the capacity of MDSCs derived from the bone marrow of *Nos2*^{+/+} and *Nos2*^{−/−} mice (Supplementary Fig. 3A–C) to inhibit proliferation and induce p53 pathway activation in CD8⁺ T cells. Second, we evaluated the effects of the iNOS inhibitor N^G-Monomethyl-L-arginine (L-NMMA) and the antioxidant N-acetyl-L-cysteine (NAC) on this phenotype. MDSCs from *Nos2*^{−/−} bone

marrow failed to inhibit T-cell proliferation, consistent with the reported inhibitory effect of NO on T cells (37–39) (Fig. 5A–D). Addition of the iNOS inhibitor L-NMMA partially restored the ability of T cells to proliferate in DC + MDSC co-cultures, while the antioxidant NAC was less effective. No additive or synergistic advantage to CD8⁺ T-cell proliferation in the presence of MDSCs was observed by combining NAC and L-NMMA, and L-NMMA at a concentration of 125 μ M remained the most effective treatment for blocking MDSC-mediated inhibition of CD8⁺ T cells (Supplementary Fig. 4A–D). Furthermore, addition of L-NMMA to DC + *Nos2*^{-/-} MDSC co-cultures had no additional effects on OT-I T-cell activation (Supplementary Fig. 4E–I). To further investigate the importance of NO production by MDSCs as a major mechanism of CD8⁺ T-cell inhibition, we isolated MDSCs directly from KPC tumors implanted into *Nos2*^{+/+} or *Nos2*^{-/-} mice. *Nos2*-deficient MDSCs were unable to inhibit CD8⁺ T-cell proliferation compared to *Nos2*-proficient MDSCs (Fig. 5A–D and Supplementary Fig. 5A–C). We also evaluated other proposed mechanisms by which MDSCs inhibit T cell function, including the expression of PD-L1 and secretion of IL10, but found that blocking these pathways did not restore CD8⁺ T-cell proliferation in DC + MDSC co-cultures (Supplementary Fig. 5D–H). One mechanism of MDSC-mediated suppression is the catabolism of L-arginine (18–22), but we found that supplementation of co-cultures with L-arginine did not improve CD8⁺ T-cell survival or proliferation in the presence of MDSCs (Supplementary Fig. 5I–J).

Our data strongly suggest that iNOS-mediated NO production is the major mechanism of MDSC-mediated CD8⁺ T-cell suppression. We therefore proceeded to investigate whether expression of iNOS by MDSCs could contribute to p53 pathway activation in CD8⁺ T cells. RT-qPCR analysis of OT-I T cell sorted from co-cultures demonstrated that p53 pathway genes were only overexpressed in CD8⁺ T cells from co-cultures with DCs + *Nos2*^{+/+} but not *Nos2*^{-/-} MDSCs (Fig. 5E). Also, addition of the iNOS inhibitor L-NMMA reduced mRNA levels of p53 pathway genes in DC + MDSC co-cultures (Fig. 5F). Finally, Western blot analysis demonstrated a higher degree of phosphorylation of p53 at serine 15 (S15) in CD8⁺ T cells from co-cultures of DCs + *Nos2*^{+/+} MDSCs compared to DCs + *Nos2*^{-/-} MDSCs (Fig. 5G and Supplementary Fig. 5K–M). We also found that expression of p53 target genes was induced in CD8⁺ T cells co-cultured with purified M-MDSCs (Fig. 5H), consistent with functional data demonstrating that the presence of M-MDSCs inhibited CD8⁺ T-cell proliferation and induced apoptosis (Fig. 1G–L).

We used confocal microscopy to further define the kinetics of p53 activation (Fig. 5I–K and Supplementary Fig. 6A–C). CD8⁺ T cells were isolated at three time points (24, 48 and 72 hours) from co-cultures with DCs, DC + *Nos2*^{+/+} MDSC or DC + *Nos2*^{-/-} MDSCs. At the 48-hour time point, we observed an increase in the fraction of phospho-p53 positive CD8⁺ T cells from DC + *Nos2*^{+/+} MDSC co-cultures (Fig. 5I–K); this increase was absent when MDSCs lacked expression of iNOS (DCs only: 8 $\pm$ 2% positive cells, DCs + *Nos2*^{+/+} MDSCs: 58 $\pm$ 11% positive cells, DCs + *Nos2*^{-/-} MDSC: 13 $\pm$ 3% positive cells). The increased phospho-p53 signal in CD8⁺ T cells was maintained at the 72-hour time point in T cells from DC + *Nos2*^{+/+} MDSC co-cultures. These data demonstrate that p53 activation in CD8⁺ T cells is dependent on iNOS expression by MDSCs.

Our data showed few MDSC-mediated effects at 24 hours and accumulation of activated p53 in CD8⁺ T cells after 48 hours following initiation of co-cultures, suggesting that there was a delay in the induction of MDSC-mediated inhibition. Previous studies show IFN γ /TNF α -mediated induction of the *Nos2* gene in stromal cells (38). Therefore, we tested the hypothesis that release of these cytokines by activated CD8⁺ T cells could induce iNOS expression in MDSCs by treating total MDSCs or FACS-sorted M-MDSCs with IFN γ , TNF α , or both cytokines for 24 hours, quantifying *Nos2* mRNA levels by qRT-PCR (Fig. 5L). Stimulation of MDSCs with IFN γ specifically and significantly induced *Nos2* mRNA in both total MDSCs (IFN γ : 6 $\pm$ 1.9-fold increase) and FACS-enriched M-MDSCs (IFN γ : 3.8 $\pm$ 0.8-fold increase). Using the Griess Reaction, we detected nitrites only in supernatants from *Nos2*^{+/+} MDSCs following stimulation with IFN γ (Fig. 5M). We also detected nitrites in co-cultures of CD8⁺ T cells with DCs + MDSCs; again, nitrites were only detected when MDSCs originated from *Nos2*^{+/+} mice (Fig. 5N). MDSC numbers correlated with their capacity to inhibit CD8⁺ T-cell proliferation (Fig. 1N), which also directly correlated with the detected concentrations of nitrites in co-cultures (Fig. 5N). These data show that IFN γ , a cytokine released by CD8⁺ T cells following activation, induced MDSC-intrinsic *Nos2* mRNA levels, resulting in the release of NO. The induction of iNOS expression in MDSCs by T cell-derived IFN γ explains why MDSCs do not block the initial steps of T-cell activation but rather induce T-cell apoptosis following IFN γ release.

iNOS-dependent induction of DNA damage in CD8⁺ T cells.

The observed activation of the p53 pathway (in addition to ‘DNA Repair’ and ‘Reactive Oxygen Species’ pathways, Fig. 4C) suggested the presence of DNA damage in CD8⁺ T cells co-cultured with MDSCs. The histone protein H2AX is rapidly phosphorylated at residue S139 (γ H2AX) by the kinases ATM and ATR in response to double-strand DNA breaks; γ H2AX therefore serves as a sensitive marker for DNA damage. Confocal microscopy of FACS-enriched CD8⁺ T cells labeled with a γ H2AX-AF488 specific antibody detected multiple nuclear γ H2AX foci at the 48-hour time point in a substantial fraction of CD8⁺ T cells from DC + *Nos2*^{+/+} MDSC co-cultures (39 $\pm$ 8% positive cells with 7 $\pm$ 3 foci), but minimally in CD8⁺ T cells from co-cultures with DCs + *Nos2*^{-/-} MDSCs (13 $\pm$ 1% positive cells with 2 $\pm$ 1 foci) or DCs (7 $\pm$ 6% positive cells with 2 $\pm$ 1 foci) (Fig. 6A–F and Supplementary Fig. 6D–F). Similar findings were made for the 72-hour time point. We also performed this analysis for co-cultures with purified M-MDSCs (Fig. 6G–I and Supplementary Fig. 7). Confocal imaging showed that both total MDSCs and M-MDSCs induced DNA damage in CD8⁺ T cells. Taken together, these data show that iNOS expression by M-MDSCs is required for the induction of DNA damage and activation of the p53 pathway in CD8⁺ T cells.

Inactivation of the *Nos2* gene reduces DNA damage in tumor-infiltrating CD8⁺ T cells.

We next examined whether iNOS expression by host cells could induce DNA damage in tumor-infiltrating T cells. KPC tumor cells (*Nos2*^{+/+} genotype) were implanted subcutaneously into *Nos2*^{+/+} or *Nos2*^{-/-} mice on a C57BL/6 background. *Nos2* deficiency reduced tumor burden and increased tumor infiltration by CD8⁺ T cells (Fig. 7A–D) and NK cells (Fig. 7E). A larger percentage of tumor-infiltrating T cells in *Nos2*^{-/-} compared to *Nos2*^{+/+} mice expressed IFN γ , TNF α , and granzyme B (Fig. 7F–I).

Confocal microscopy of sorted CD8⁺ T cells from KPC tumors that had been implanted into *Nos2*^{+/+} or *Nos2*^{-/-} mice showed that a larger fraction of cells from *Nos2*^{+/+} compared to *Nos2*^{-/-} mice were positive for nuclear γ H2AX foci (*Nos2*^{+/+} 74.8 $\pm$ 11.8% of cells; *Nos2*^{-/-} 26.9 $\pm$ 9.7% of cells) (Fig. 7J–M and Supplementary Fig. 8). This difference was also reflected by the number of γ H2AX foci per positive cells (*Nos2*^{+/+} 6 $\pm$ 2 foci/cell; *Nos2*^{-/-} 3 $\pm$ 1 foci/cell) and a decrease in the integrated fluorescence intensity/cell (*Nos2*^{+/+} 1762 $\pm$ 918 AU/cell; *Nos2*^{-/-} 777 $\pm$ 564 AU/cell). These data provide *in vivo* evidence for DNA damage in tumor-infiltrating CD8⁺ T cells that is dependent on iNOS expression by host cells (Fig. 7N).

Discussion:

These studies demonstrate that MDSCs potently inhibit CD8⁺ T cells, even in the presence of DCs presenting a high-affinity cognate peptide. ScRNA-seq analysis demonstrated that the presence of MDSCs did not prevent the expression of key activation-induced genes at the 24-hour time point. Rather, CD8⁺ T cells expressed several genes associated with p53 pathway activation at a later time point, including those associated with cell cycle arrest (*Cdk1na*, *Phlda3*) and apoptosis (*Bax*). Confocal microscopy demonstrated induction of DNA damage in CD8⁺ T cells, which was dependent on expression of the *Nos2* gene by MDSCs. IFN γ , a key cytokine released by activated CD8⁺ T cells, increased expression of *Nos2* and release of NO by MDSCs, explaining why initial T-cell activation remained intact but T-cell proliferation and survival were profoundly impaired. This finding was relevant to the *in vivo* setting because tumor-infiltrating CD8⁺ T cells from *Nos2*-deficient compared to *Nos2*-proficient mice had significantly less DNA damage.

Several previous studies demonstrate that NO released by MDSCs is important for inhibition of T-cell proliferation, and our data are consistent with these reports (10,20,26,36). We also show that T-cell proliferation is strongly dependent on the number of MDSCs and the concentration of released NO. However, previous studies proposed that MDSCs block initial T-cell activation by nitrosylation of the TCR, CD8 co-receptor, or signaling molecules (26,36). NO produced by iNOS can freely diffuse across cellular membranes and react with cysteine residues (40); it can also react with superoxide produced by NADPH oxidase to form peroxynitrite, a highly reactive compound that can oxidize cysteine and nitrosylate tyrosine residues (24). Such protein modifications can be identified by Western blotting in T cells co-cultured with MDSCs (26,36). However, our data demonstrate that MDSCs are inhibitory even when added to co-cultures with T cells + DCs after a delay of 24 or 48 hours. Furthermore, they inhibit T-cell proliferation induced by cytokines rather than TCR stimulation. The scRNA-seq data clearly demonstrate the induction of an activation-related transcriptional signature in T cells from co-cultures with DCs + MDSCs, although a shift in the distribution of T cells across some of the sub-clusters was noted at the 24-hour time point. The presence of MDSCs could certainly modify early T-cell activation processes, but we did not observe a block in initial T-cell activation in our experiments based on assessment of early T-cell activation markers and cytokine production.

Several suppressive mechanisms have been described for MDSCs, including release of IL-10 and catabolism of arginine (see Supplementary Table 1 for details) (18–22). However, we

found that inhibition of NO release alone was sufficient to block MDSC-mediated inhibition of CD8⁺ T-cell proliferation. Furthermore, induction of DNA damage and activation of the p53 pathway in T cells was only observed with MDSCs from *Nos2*^{+/+} but not *Nos2*^{-/-} mice, indicating a dependence on NO synthesis. Consistent with this, we observed increased T-cell apoptosis in the presence of MDSCs concurrent with a decline in T-cell numbers at later time points. These findings are consistent with the hypothesis that DNA damage and p53 pathway activation are relevant for the loss of T cells in these cultures. NO and peroxynitrite can induce deamination of DNA bases (including guanine and cytosine) and DNA breaks (41). However, these highly reactive compounds can also modify proteins and membrane lipids, and we therefore do not propose that the induction of DNA damage is the sole mechanism for the observed inhibition of T-cell proliferation and the drastic decline in T-cell numbers. Other mechanisms including depletion of key amino acids such as L-arginine, release of inhibitory cytokines and expression of inhibitory surface molecules are likely to be important contributors to the *in vivo* function of MDSCs (17,27,28). The relative importance of these mechanisms may depend on features of the tumor microenvironment.

We observed reduced tumor weight when KPC cells were implanted into *Nos2*^{-/-} compared to *Nos2*^{+/+} mice. Also, KPC tumors from *Nos2*^{-/-} mice showed greater infiltration by CD8⁺ T cells, including IFN γ ⁺ and Gzmb⁺ CD8⁺ T cells. Confocal imaging demonstrated that a substantially smaller percentage of tumor-infiltrating CD8⁺ T cells from *Nos2*^{-/-} hosts had evidence of DNA damage based on γ H2AX staining, consistent with our conclusion that MDSCs can induce DNA damage and apoptosis in activated T cells. Reduced tumor growth in *Nos2*^{-/-} compared *Nos2*^{+/+} mice may also reflect the dynamic relationship between T cell-mediated immunity and MDSC-mediated suppression: a reduced tumor mass may also decrease secretion of growth factors that promote differentiation of myeloid cells to immunosuppressive MDSCs, thus favoring T cell-mediated tumor immunity.

Tumors with substantial infiltration by MDSCs are characterized by impaired T cell-mediated tumor immunity (42–48). The profound block in proliferation and induction of apoptosis in T cells by MDSCs is therefore a significant barrier to tumor-immunity. Strategies that inhibit the release of NO by MDSCs, block its detrimental effects on T cells, or reduce MDSC accumulation into tumors could therefore help to promote protective T cell-mediated tumor immunity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We would like to thank Dr. Stephanie Dougan for providing the murine KPC tumor cell line. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health/NCI.

Funding information:

This work was supported by NIH grants R01 CA238039, R01 CA251599, P01 CA163222 (to K.W.W.), T32 CA207021 (to A.L., A.B. and K.W.W.), fellowships from AACR (to S.K.), and the Cancer Research Institute (CRI, to ANRC).

References

1. Meyer C, Sevko A, Ramacher M, Bazhin AV, Falk CS, Osen W, et al. Chronic inflammation promotes myeloid-derived suppressor cell activation blocking antitumor immunity in transgenic mouse melanoma model. *Proc Natl Acad Sci U S A* 2011;108(41):17111–6 doi 10.1073/pnas.1108121108. [PubMed: 21969559]
2. Norris BA, Uebelhoefer LS, Nakaya HI, Price AA, Grakoui A, Pulendran B. Chronic but not acute virus infection induces sustained expansion of myeloid suppressor cell numbers that inhibit viral-specific T cell immunity. *Immunity* 2013;38(2):309–21 doi 10.1016/j.immuni.2012.10.022. [PubMed: 23438822]
3. Metcalf D Hematopoietic cytokines. *Blood* 2008;111(2):485–91. [PubMed: 18182579]
4. Ueha S, Shand FH, Matsushima K. Myeloid cell population dynamics in healthy and tumor-bearing mice. *Int Immunopharmacol* 2011;11(7):783–8 doi 10.1016/j.intimp.2011.03.003. [PubMed: 21406269]
5. Ai L, Mu S, Wang Y, Wang H, Cai L, Li W, et al. Prognostic role of myeloid-derived suppressor cells in cancers: a systematic review and meta-analysis. *BMC Cancer* 2018;18(1):1220 doi 10.1186/s12885-018-5086-y. [PubMed: 30518340]
6. Wang Z, Liu Y, Zhang Y, Shang Y, Gao Q. MDSC-decreasing chemotherapy increases the efficacy of cytokine-induced killer cell immunotherapy in metastatic renal cell carcinoma and pancreatic cancer. *Oncotarget* 2016;7(4):4760–9 doi 10.18632/oncotarget.6734. [PubMed: 26716894]
7. Weide B, Martens A, Zelba H, Stutz C, Derhovanessian E, Di Giacomo AM, et al. Myeloid-derived suppressor cells predict survival of patients with advanced melanoma: comparison with regulatory T cells and NY-ESO-1- or melan-A-specific T cells. *Clin Cancer Res* 2014;20(6):1601–9 doi 10.1158/1078-0432.Ccr-13-2508. [PubMed: 24323899]
8. Bronte V, Brandau S, Chen SH, Colombo MP, Frey AB, Greten TF, et al. Recommendations for myeloid-derived suppressor cell nomenclature and characterization standards. *Nat Commun* 2016;7:12150 doi 10.1038/ncomms12150. [PubMed: 27381735]
9. Weber R, Fleming V, Hu X, Nagibin V, Groth C, Altevogt P, et al. Myeloid-Derived Suppressor Cells Hinder the Anti-Cancer Activity of Immune Checkpoint Inhibitors. *Front Immunol* 2018;9:1310 doi 10.3389/fimmu.2018.01310. [PubMed: 29942309]
10. Gabrilovich DI. Myeloid-Derived Suppressor Cells. *Cancer Immunol Res* 2017;5(1):3–8 doi 10.1158/2326-6066.Cir-16-0297. [PubMed: 28052991]
11. Park MJ, Lee SH, Kim EK, Lee EJ, Baek JA, Park SH, et al. Interleukin-10 produced by myeloid-derived suppressor cells is critical for the induction of Tregs and attenuation of rheumatoid inflammation in mice. *Sci Rep* 2018;8(1):3753 doi 10.1038/s41598-018-21856-2. [PubMed: 29491381]
12. Raber P, Ochoa AC, Rodriguez PC. Metabolism of L-arginine by myeloid-derived suppressor cells in cancer: mechanisms of T cell suppression and therapeutic perspectives. *Immunol Invest* 2012;41(6–7):614–34 doi 10.3109/08820139.2012.680634. [PubMed: 23017138]
13. Stiff A, Trikha P, Mundy-Bosse B, McMichael E, Mace TA, Benner B, et al. Nitric Oxide Production by Myeloid-Derived Suppressor Cells Plays a Role in Impairing Fc Receptor-Mediated Natural Killer Cell Function. *Clin Cancer Res* 2018;24(8):1891–904 doi 10.1158/1078-0432.CCR-17-0691. [PubMed: 29363526]
14. Aarts CEM, Hiemstra IH, Beguin EP, Hoogendijk AJ, Bouchmal S, van Houdt M, et al. Activated neutrophils exert myeloid-derived suppressor cell activity damaging T cells beyond repair. *Blood Adv* 2019;3(22):3562–74 doi 10.1182/bloodadvances.2019031609. [PubMed: 31738831]
15. Noman MZ, Desantis G, Janji B, Hasmim M, Karray S, Dessen P, et al. PD-L1 is a novel direct target of HIF-1 α , and its blockade under hypoxia enhanced MDSC-mediated T cell activation. *J Exp Med* 2014;211(5):781–90 doi 10.1084/jem.20131916. [PubMed: 24778419]
16. Hanson EM, Clements VK, Sinha P, Ilkovitch D, Ostrand-Rosenberg S. Myeloid-derived suppressor cells down-regulate L-selectin expression on CD4⁺ and CD8⁺ T cells. *J Immunol* 2009;183(2):937–44 doi 10.4049/jimmunol.0804253. [PubMed: 19553533]

17. Srivastava MK, Sinha P, Clements VK, Rodriguez P, Ostrand-Rosenberg S. Myeloid-derived suppressor cells inhibit T-cell activation by depleting cystine and cysteine. *Cancer Res* 2010;70(1):68–77 doi 10.1158/0008-5472.Can-09-2587. [PubMed: 20028852]
18. Mao Y, Sarhan D, Steven A, Seliger B, Kiessling R, Lundqvist A. Inhibition of tumor-derived prostaglandin-e2 blocks the induction of myeloid-derived suppressor cells and recovers natural killer cell activity. *Clin Cancer Res* 2014;20(15):4096–106 doi 10.1158/1078-0432.CCR-14-0635. [PubMed: 24907113]
19. Zhu J, Powis de Tenbossche CG, Cane S, Colau D, van Baren N, Lurquin C, et al. Resistance to cancer immunotherapy mediated by apoptosis of tumor-infiltrating lymphocytes. *Nat Commun* 2017;8(1):1404 doi 10.1038/s41467-017-00784-1. [PubMed: 29123081]
20. Mazzoni A, Bronte V, Visintin A, Spitzer JH, Apolloni E, Serafini P, et al. Myeloid suppressor lines inhibit T cell responses by an NO-dependent mechanism. *J Immunol* 2002;168(2):689–95 doi 10.4049/jimmunol.168.2.689. [PubMed: 11777962]
21. Aydin E, Johansson J, Nazir FH, Hellstrand K, Martner A. Role of NOX2-Derived Reactive Oxygen Species in NK Cell-Mediated Control of Murine Melanoma Metastasis. *Cancer Immunol Res* 2017;5(9):804–11 doi 10.1158/2326-6066.Cir-16-0382. [PubMed: 28760732]
22. Corzo CA, Cotter MJ, Cheng P, Cheng F, Kusmartsev S, Sotomayor E, et al. Mechanism regulating reactive oxygen species in tumor-induced myeloid-derived suppressor cells. *J Immunol* 2009;182(9):5693–701 doi 10.4049/jimmunol.0900092. [PubMed: 19380816]
23. Caulfield JL, Wishnok JS, Tannenbaum SR. Nitric Oxide-induced Deamination of Cytosine and Guanine in Deoxynucleosides and Oligonucleotides. *The Journal of Biological Chemistry* 1998;273(21):12689–95. [PubMed: 9582291]
24. Galatro A, Gonzalez PM, Malanga G, Robello E, Piloni NE, Puntarulo S. Nitric oxide and membrane lipid peroxidation in photosynthetic and non-photosynthetic organisms under several stress conditions. *Front Physiol* 2013;4:276 doi 10.3389/fphys.2013.00276. [PubMed: 24146649]
25. Keyer K, Imlay JA. Superoxide accelerates DNA damage by elevating free-iron levels. *Proc Natl Acad Sci U S A* 1996;93:13635–40. [PubMed: 8942986]
26. Nagaraj S, Schrum AG, Cho HI, Celis E, Gabrilovich DI. Mechanism of T cell tolerance induced by myeloid-derived suppressor cells. *J Immunol* 2010;184(6):3106–16 doi 10.4049/jimmunol.0902661. [PubMed: 20142361]
27. Rodriguez PC, Zea AH, Culotta KS, Zabaleta J, Ochoa JB, Ochoa AC. Regulation of T cell receptor CD3zeta chain expression by L-arginine. *J Biol Chem* 2002;277(24):21123–9 doi 10.1074/jbc.M110675200. [PubMed: 11950832]
28. Munn DH, Sharma MD, Baban B, Harding HP, Zhang Y, Ron D, et al. GCN2 kinase in T cells mediates proliferative arrest and anergy induction in response to indoleamine 2,3-dioxygenase. *Immunity* 2005;22(5):633–42 doi 10.1016/j.immuni.2005.03.013. [PubMed: 15894280]
29. Ibrahim ML, Klement JD, Lu C, Redd PS, Xiao W, Yang D, et al. Myeloid-Derived Suppressor Cells Produce IL-10 to Elicit DNMT3b-Dependent IRF8 Silencing to Promote Colitis-Associated Colon Tumorigenesis. *Cell Rep* 2018;25(11):3036–46 e6 doi 10.1016/j.celrep.2018.11.050. [PubMed: 30540937]
30. Youn JI, Nagaraj S, Collazo M, Gabrilovich DI. Subsets of myeloid-derived suppressor cells in tumor-bearing mice. *J Immunol* 2008;181(8):5791–802 doi 10.4049/jimmunol.181.8.5791. [PubMed: 18832739]
31. Pijuan-Sala B, Griffiths JA, Guibentif C, Hiscock TW, Jawaid W, Calero-Nieto FJ, et al. A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature* 2019;566(7745):490–5 doi 10.1038/s41586-019-0933-9. [PubMed: 30787436]
32. McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* 2019;8(4):329–37 e4 doi 10.1016/j.cels.2019.03.003. [PubMed: 30954475]
33. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* 2018;36(5):411–20 doi 10.1038/nbt.4096. [PubMed: 29608179]

34. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM 3rd, et al. Comprehensive Integration of Single-Cell Data. *Cell* 2019;177(7):1888–902 e21 doi 10.1016/j.cell.2019.05.031. [PubMed: 31178118]
35. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. *bioRxiv* 2019 doi 10.1101/576827.
36. Feng S, Cheng X, Zhang L, Lu X, Chaudhary S, Teng R, et al. Myeloid-derived suppressor cells inhibit T cell activation through nitrating LCK in mouse cancers. *Proc Natl Acad Sci U S A* 2018;115(40):10094–9 doi 10.1073/pnas.1800695115. [PubMed: 30232256]
37. Sato K, Ozaki K, Oh I, Meguro A, Hatanaka K, Nagai T, et al. Nitric oxide plays a critical role in suppression of T-cell proliferation by mesenchymal stem cells. *Blood* 2007;109(1):228–34 doi 10.1182/blood-2006-02-002246. [PubMed: 16985180]
38. Lukacs-Kornek V, Malhotra D, Fletcher AL, Acton SE, Elpek KG, Tayalia P, et al. Regulated release of nitric oxide by nonhematopoietic stroma controls expansion of the activated T cell pool in lymph nodes. *Nat Immunol* 2011;12(11):1096–104 doi 10.1038/ni.2112. [PubMed: 21926986]
39. Niedbala W, Cai B, Liew FY. Role of nitric oxide in the regulation of T cell functions. *Ann Rheum Dis* 2006;65 Suppl 3:iii37–40 doi 10.1136/ard.2006.058446. [PubMed: 17038470]
40. Marino SM, Gladyshev VN. Analysis and functional prediction of reactive cysteine residues. *J Biol Chem* 2012;287(7):4419–25 doi 10.1074/jbc.R111.275578. [PubMed: 22157013]
41. Caulfield JL, Wishnok JS, Tannenbaum SR. Nitric oxide-induced deamination of cytosine and guanine in deoxynucleosides and oligonucleotides. *J Biol Chem* 1998;273(21):12689–95 doi 10.1074/jbc.273.21.12689. [PubMed: 9582291]
42. Binsfeld M, Muller J, Lamour V, De Veirman K, De Raeve H, Bellahcene A, et al. Granulocytic myeloid-derived suppressor cells promote angiogenesis in the context of multiple myeloma. *Oncotarget* 2016;7(25):37931–43 doi 10.18632/oncotarget.9270. [PubMed: 27177328]
43. Bunt SK, Sinha P, Clements VK, Leips J, Ostrand-Rosenberg S. Inflammation induces myeloid-derived suppressor cells that facilitate tumor progression. *J Immunol* 2006;176(1):284–90 doi 10.4049/jimmunol.176.1.284. [PubMed: 16365420]
44. Kather JN, Suarez-Carmona M, Charoentong P, Weis CA, Hirsch D, Bankhead P, et al. Topography of cancer-associated immune cells in human solid tumors. *Elife* 2018;7 doi 10.7554/eLife.36967.
45. Wang Y, Ding Y, Guo N, Wang S. MDSCs: Key Criminals of Tumor Pre-metastatic Niche Formation. *Front Immunol* 2019;10:172 doi 10.3389/fimmu.2019.00172. [PubMed: 30792719]
46. Wang L, Chang EW, Wong SC, Ong SM, Chong DQ, Ling KL. Increased myeloid-derived suppressor cells in gastric cancer correlate with cancer stage and plasma S100A8/A9 proinflammatory proteins. *J Immunol* 2013;190(2):794–804 doi 10.4049/jimmunol.1202088. [PubMed: 23248262]
47. Solito S, Marigo I, Pinton L, Damuzzo V, Mandruzzato S, Bronte V. Myeloid-derived suppressor cell heterogeneity in human cancers. *Ann N Y Acad Sci* 2014;1319:47–65 doi 10.1111/nyas.12469. [PubMed: 24965257]
48. Zhu H, Gu Y, Xue Y, Yuan M, Cao X, Liu Q. CXCR2+ MDSCs promote breast cancer progression by inducing EMT and activated T cell exhaustion. *Oncotarget* 2017;8(70):114554–67. [PubMed: 29383101]

Synopsis:

MDSCs inhibit T cell-mediated tumor immunity. The data show that MDSCs do not block early steps of T-cell activation but rather induce nitric oxide-dependent DNA damage and subsequent p53 pathway activation in T cells.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. MDSCs do not prevent initial activation of CD8⁺ T cells but interfere with subsequent proliferation and survival.

A, Schematic diagram of experimental design. CD8⁺ T cells (OT-I cells) and SIINFEKL peptide-pulsed DCs were co-cultured with or without MDSCs for up to 72 hours. The impact of MDSCs on T-cell function was studied by scRNA-seq, confocal microscopy, and flow cytometry. **B-F**, Co-cultures were set up for CFSE-labeled OT-I T cells (50,000/well) with peptide-pulsed DCs or DCs + MDSCs (ratio of OT-I: DC: MDSC = 1: 0.2: 2). **B**, Analysis of CFSE dilution and expression of CD69/CD25 activation markers by CD8⁺ T cells at 24 and 72 hours. **C-E**, Summary of data as fraction of divided (C), CD69⁺ (D), and CD25⁺ CD8⁺ T cells (E). **F**, Number of CD8⁺ T cells recovered from co-culture assays at 24 and 72 hours normalized to DC only condition. **G-L**, M-MDSCs and PMN-MDSCs were isolated by FACS. Co-cultures with OT-I T cells (50,000/well) were set up for DC only

(black), DC + MDSC (red), DC + PMN-MDSC (orange), and DC + M-MDSC (magenta) conditions (OT-I: DC = 1: 0.2; OT-I: DC: MDSC = 1: 0.2: 2; OT-I: DC: PMN-MDSC = 1: 0.2: 2; OT-I: DC: M-MDSC = 1: 0.2: 1). **G**, Number of CD8 T cells recovered from co-cultures at 72 hours. **H-J**, Analysis of T-cell proliferation. Histograms showing CFSE dilution of CD8⁺ T cells (**H**), percentage of divided CD8⁺ T cells (**I**), and mean fluorescence intensity (MFI) of CFSE (**J**) at 72 hours. **K-L**, Analysis of CD8⁺ T-cell apoptosis at 72 hours for the indicated conditions. Flow cytometry contour plots show Annexin V/Zombie UV staining (**K**). Percentage of CD8⁺ T cells that were viable (green), early apoptotic (orange) or late apoptotic (red) at 72 hours (**L**). **M-N**, Co-cultures with different ratios of T cells to MDSCs. Co-culture assays were performed with decreasing numbers of MDSCs (relative to OT-I cells) and OT-I proliferation was assessed after 72 hours. Graph shows the percentage of divided OT-I cells in each condition. **O-R**, MDSCs were sorted by FACS from subcutaneous KPC pancreatic tumors. CD8⁺ T cells (50,000/well) were co-cultured with SIINFEKL-pulsed DCs with and without tumor-derived MDSCs for 72 hours (OT-I: DC: *ex vivo* MDSC = 1: 0.2: 2). **P-R**, Analysis of CD8⁺ T cell proliferation based on CFSE dilution for DCs only (left) or DC + *ex vivo* MDSC (right) conditions. Histogram of CFSE dilution (**M**), summary of data as percentage of divided CD8⁺ T cells (**N**), CFSE MFI (**O**), and the number of recovered CD8⁺ T cells (**P**) from the two indicated conditions. Data are representative of 2 (**G-R**) or 3 (**B-F**) independent experiments. Bars indicate mean $\pm$ s.d. Statistical analysis was performed using two-way ANOVA with Sidak's multiple comparisons (C-G), one-way ANOVA with Dunnett's multiple comparisons (I-J, M), two-way ANOVA with Dunnett's multiple comparisons (L), or an unpaired two-tailed t-test (P-R). ns = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 2. MDSC-mediated inhibition occurs even following initial T-cell activation.

A-C, MDSCs were added to co-cultures of CFSE-labeled OT-I CD8⁺ T cells (50,000/well) and SIINFEKL peptide-pulsed DCs at the indicated time points; T-cell proliferation was assessed after 96 hours (ratio of OT-I: DC: MDSC = 1: 0.2: 2). **A**, Histograms showing CFSE dilution following addition of MDSCs at the indicated time points. **B**, MFI of CFSE relative to DC only condition. **C**, Number of CD8⁺ T cells recovered from co-culture assays relative to DC only condition at 96 hours. **D-F**, CFSE-labeled OT-I CD8⁺ T cells were activated by SIINFEKL-pulsed DCs or IL-15/IL-7. These cultures were set up with or without MDSCs (red and black lines, respectively). **D**, Histograms showing CFSE dilution of CD8⁺ T cells in co-culture assays with DCs +/- MDSC (left) or following IL-15/IL-7 stimulation (right) +/- MDSC. **E**, MFI of CFSE relative to conditions without MDSC. **F**, Number of CD8⁺ T cells recovered from co-culture assays relative to activation conditions without MDSCs. Data are representative of 2 independent experiments. Bars indicate mean $\pm$ s.d. Statistical analysis was performed using one-way ANOVA with Dunnett's multiple comparisons (B-C) or two-way ANOVA with Sidak's multiple comparisons (E-F).

***p<0.0001.

Figure 3. Impact of MDSCs on the transcriptome of CD8⁺ T cells at a single-cell level. **A-F**, OT-I CD8⁺ T cells and SIINFEKL-pulsed DCs were co-cultured with or without MDSCs, and after 24 and 72 hours, CD8⁺ T cells were isolated by FACS for scRNA-seq analysis. Naïve CD8⁺ T cells were also analyzed prior to co-culture. **A**, Visualization of CD8⁺ T-cell clusters from different experimental conditions using a UMAP plot. Each dot represents a single cell. Conditions are color-coded as indicated. **B-D**, Analysis of CD8⁺ T cells at 24 hours following co-culture with DCs or DCs + MDSCs. Each dot represents a single cell. **B**, T cells are color-coded in UMAP plot based on culture conditions. **C**, Sub-clustering of CD8⁺ T cells. The indicated genes are differentially expressed in indicated clusters. **D**, Origin of cells from particular clusters for DC only (orange) or DC + MDSC (cyan) culture conditions. **E**, Heat map showing the top ten differentially expressed genes for

each cluster identified in **C**. Expression is normalized to Log₂ fold change ranging from +2 (yellow) to -2 (purple). Selected genes are highlighted with arrows. **F**, Expression of selected genes across clusters visualized using UMAP plots. **G**, OT-I CD8⁺ T cells and SIINFEKL-pulsed DCs were co-cultured with or without MDSCs for up to 72 hours. Expression of granzyme B and IFN γ was assessed by intracellular staining following PMA/Ionomycin stimulation for 6 hours. ScRNA-seq (**A-F**) was performed once using cells from biological triplicate cultures that were pooled prior to T cell isolation. Experiment shown in (**G**) is representative of two independent experiments. Bars represent mean $\pm$ s.d. Statistical analysis was performed using unpaired two-tailed t-test (**G**). ns = not significant, ***p<0.0001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 4. MDSCs induce activation of the p53 pathway in CD8⁺ T cells.

Analysis of scRNA-seq data for the 72-hour time point. **A**, Heat map showing differentially expressed genes for DC only and DC + MDSC conditions at 72 hours. Expression is normalized to Log₂ fold change ranging from +2 (yellow) to -2 (purple). **B**, Feature plots showing expression of genes overexpressed by CD8⁺ T cells from the DC + MDSC condition. Gene expression is scaled from 0 (grey) to 2/3 (purple). **C**, GSEA analysis of Hallmark enriched pathways for highly variable genes in DC + MDSC compared to DC only conditions at 72 hours. **D**, Expression dot plot showing Log₂ fold change (diameter) and *p*-value (color) of genes enriched in pathways (from C) for DC+MDSC compared to DC only conditions. **E**, UMAP projection showing sub-clusters of CD8⁺ T cells for the indicated conditions. Each dot represents a single cell. Clusters are color-coded and relevant clusters

are labeled as indicated. **F**, GSEA analysis of Hallmark pathways enriched in clusters 2, 3, and 4. **G**, Expression dot plot showing Log₂ fold change (diameter) and *p*-value (color) of the indicated p53 pathway genes in clusters 2, 3, and 4. **H**, Analysis of mRNA levels of p53 pathway genes. OT-I CD8⁺ T cells were sorted from co-culture assays and gene expression was assessed by quantitative RT-PCR. Graph shows expression of the indicated genes in DC + MDSC condition relative to DC only condition at 24 and 72 hours. ScRNA-seq was performed once using pooled biological triplicates for each condition (**A-G**); qPCR data are representative of 3 independent experiments (**H**). Bars indicate mean ± s.d. Analysis of single cell data was performed using Monocle v3 (**A, B, E, G**). Statistical analysis was performed using unpaired two-tailed t-test (**H**), **p*<0.05, ***p*<0.01, *****p*<0.0001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 5. p53 pathway activation in CD8⁺ T cells is dependent on iNOS expression by MDSCs. **A-D**, MDSCs were sorted from KPC tumors implanted subcutaneously into *Nos2*^{+/+} or *Nos2*^{-/-} mice. OT-I CD8⁺ T cells (50,000/well) and SIINFEKL-pulsed DCs were co-cultured with and without tumor-derived MDSCs for 72 hours (OT-I: DC: MDSC = 1: 0.2: 2). T cells were analyzed in terms of CFSE dilution (**A**, **B**), CD69 expression (**A**, **C**), and CD25 expression (**A**, **D**). **E-F**, OT-I CD8⁺ T cells were sorted at 72 hours from co-culture assays and assessed for expression of p53 target genes by RT-qPCR (comparison relative to DC only condition). CD8⁺ T cells were co-cultured with DCs + MDSCs from *Nos2*^{+/+} or *Nos2*^{-/-} mice (**E**); alternatively, CD8⁺ T cells were co-cultured with DCs + MDSCs in the presence of the iNOS inhibitor L-NMMA (125 μM) or solvent control (**F**). **G**, Western blot showing total and phosphorylated p53 (serine 15 site) from CD8⁺ T cells co-cultured with

DCs or DCs + MDSCs isolated from *Nos2*^{+/+} or *Nos2*^{-/-} mice. **H**, mRNA levels of p53 target genes at 48 hours for OT-I T cells sorted from co-cultures with DC only (black), DC + total MDSC (red), DC + PMN-MDSC (orange), or DC + M-MDSC (magenta) (comparison relative to DC only condition). **I-K**, Imaging of p53 activation in CD8⁺ T cells. Representative images showing phospho-p53 (serine 15) antibody labeling of CD8⁺ T cells at 72 hours from DC only or DC + MDSC from *Nos2*^{+/+} or *Nos2*^{-/-} mice (**J**). Data were quantified based on the percentage of phospho-p53 positive T cells (**J**) or the integrated phospho-p53 intensity (**K**) at the indicated time points. Number of analyzed cells ranged from 389 to 401 for the different conditions. **L**, Total MDSCs and FACS-enriched M-MDSCs were stimulated with IFN γ , TNF α or both cytokines for 24 hours, and expression of *Nos2* was assessed by qPCR. Expression is shown relative to unstimulated cells. **M**, Griess reaction measurements of nitrite concentrations in the supernatant after 48 hours stimulation of *Nos2*^{+/+} or *Nos2*^{-/-} MDSCs with the indicated concentration of IFN γ . **N**, Griess reaction measurements of nitrite concentrations in the supernatant of co-culture assays with the indicated T cell: MDSC ratios. Data are representative of two (**A-D**, **G-N**) or three (**E-F**) independent experiments. Bars indicate mean $\pm$ s.d. Statistical analysis was performed using one way-ANOVA with Dunnett's multiple comparisons (**B-D**), unpaired two-tailed t-test (**E-F**, **H**, **L**), one-way ANOVA with Holm-Sidak's multiple comparisons (**J-K**, **N**), or two-way ANOVA with Tukey's multiple comparisons (**M**). ns = not significant, **p*<0.05, ***p*<0.01, ****p*<0.001, *****p*<0.0001.

Figure 6. MDSCs induce DNA damage in CD8⁺ T cells.

OT-I T cells and SIINFEKL-pulsed DCs were co-cultured with or without MDSCs derived from *Nos2*^{+/+} or *Nos2*^{-/-} mice. **A-C**, Analysis of CFSE dilution after 24-, 48-, and 72-hours of co-culture. **B**, Number of CD8⁺ T cells recovered from each condition and time point relative to DC only condition. **C**, Percentage of divided CD8⁺ T cells in each condition and time point. **D**, Representative images showing labeling with γH2AX antibody in CD8⁺ T cells after co-culture for 24, 48 or 72 hours with DCs or DCs + MDSCs from *Nos2*^{+/+} or *Nos2*^{-/-} mice. **E**, Fraction of CD8⁺ T cells containing more than one focus positive for γH2AX. **F**, Number of γH2AX positive foci per cell. **G**, DNA damage in T cells following co-culture with different MDSC subpopulations. Representative images showing labeling with γH2AX antibody in CD8⁺ T cells after 48 hours of coculture with DC, DC + total

MDSC, DC + PMN-MDSC or DC + M-MDSC. **H**, Fraction of CD8⁺ T cells containing more than one focus positive for γ H2AX. **I**, Number of γ H2AX positive foci per cell. Analysis was performed on 166 cells for DC only, 98 cells for DC + MDSC, 77 cells for DC + PMN-MDSC, and 137 cells for DC + M-MDSC conditions. Data are representative of two independent experiments. Statistical analysis was performed using two-way ANOVA with Tukey's multiple comparisons (B-C) or one-way ANOVA with Holm-Sidak's multiple comparisons (E-F, H-I). ns = not significant, **p< 0.01, ***p<0.001, ****p<0.0001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 7. DNA damage in tumor-infiltrating CD8⁺ T cells from *Nos2* wild-type mice. KPC tumor cells were implanted subcutaneously into the flanks of *Nos2*^{+/+} (black) or *Nos2*^{-/-} (red) mice and harvested after 24 days. **A**, Quantification of tumor weight. **B-E**, Quantification of CD45⁺ cells (**B**), CD3⁺ cells (**C**), CD8⁺ T cells (**D**), and NK cells (**E**) per milligram of tumor. **F-I**, Percentage of CD8⁺ T cells expressing IFN γ (**F**), granzyme B (**G**), TNF α (**H**) and CD8⁺ T cells co-stained for CD107a and granzyme B (**I**). **J-M**, Tumor-infiltrating CD8⁺ T cells were enriched from KPC tumor single-cell suspensions by FACS. Representative images show labeling with γ H2AX antibody in CD8⁺ T cells from KPC tumors implanted into *Nos2*^{+/+} or *Nos2*^{-/-} mice (**J**). The number of γ H2AX positive foci per cell (**K**), the percentage of CD8⁺ T cells containing more than one focus positive for γ H2AX (**L**), and the integrated γ H2AX intensity per cell (**M**) were quantified. **N**, Schematic

showing the proposed mechanism of MDSC-mediated induction of CD8⁺ T-cell apoptosis. Bars indicate mean $\pm$ s.d; 100 or 87 CD8⁺ T cells were analyzed for *Nos2*^{+/+} and *Nos2*^{-/-} conditions, respectively. Data are representative of two independent experiments. Statistical analysis was performed using unpaired two-tailed t-test. ns = not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript