Published in final edited form as:

Curr Opin Cell Biol. 2020 October ; 66: 51–58. doi:10.1016/j.ceb.2020.05.003.

Organoid models for mammary gland dynamics and breast cancer

Vasudha Srivastava¹, Tyler R. Huycke¹, Kiet T. Phong^{1,2}, Zev J Gartner^{1,3,4,5}

¹Department of Pharmaceutical Chemistry, University of California San Francisco, CA

²University of California Berkeley–University of California San Francisco Graduate Program in Bioengineering, San Francisco, CA

³Helen Diller Family Comprehensive Cancer Center, San Francisco, CA

⁴Chan Zuckerberg BioHub, University of California San Francisco, San Francisco, CA, USA.

⁵Center for Cellular Construction, University of California San Francisco, San Francisco, CA, USA

Abstract

The mammary gland is a highly dynamic tissue that undergoes repeated cycles of growth and involution during pregnancy and menstruation. It is also the site from which breast cancers emerge. Organoids provide an in vitro model that preserves several of the cellular, structural and microenvironmental features that dictate mammary gland function in vivo, and have greatly advanced our understanding of glandular biology. Their tractability for genetic manipulation, live imaging and high throughput screening have facilitated investigation into the mechanisms of glandular morphogenesis, structural maintenance, tumor progression, and invasion. Opportunities remain to enhance cellular and structural complexity of mammary organoid models, including incorporating additional cell types and hormone signaling.

Introduction

Unlike other tissues in the body, the mammary gland undergoes the majority of its development after birth (Fig 1a) when a highly branched ductal epithelium emerges from a rudimentary embryonic ductal tree. The branched ductal epithelium comprises three distinct epithelial lineages. Lining the luminal space are two populations of luminal cells (LEP), which are responsible for milk production and hormone sensing. LEPs are surrounded by an outer population of contractile myoepithelial cells (MEP) that serve in milk ejection during breast-feeding (Fig. 1b). The epithelium is surrounded by a basement membrane and is embedded in a complex stroma containing fibroblasts, nerves, vasculature, lymphatics, immune cells and adipocytes (Fig. 1b). Signals from the surrounding non-epithelial cells, extracellular matrix (ECM) and hormones further guide the structure and function of the mammary gland. The epithelium is also the site where most breast cancers originate (Fig. 1c), a disease that affects 1 in 8 women in the United States.

The mammary gland has been studied extensively as a model tissue, encompassing multiple cell types that undergo large-scale morphological changes in response to stimuli. Early histological studies established changes in tissue composition and structure during development, reproductive cycle and breast cancer [1], but these were only snapshots of highly dynamic processes. In contrast, lineage tracing studies in genetically engineered mouse models (GEMM) provided insight into lineage specification and developmental dynamics [2], but may not accurately reflect human biology [1]. Three-dimensional cell culture of immortalized human mammary epithelial cells can provide insight into human-specific cell biology and have established the importance of structural cues and paracrine signaling in the breast [3]. Frequently referred to as “acini” or “cysts”, these tissue models, however, lack the multiple cell types found in vivo, and suffer from molecular and physical changes associated with long-term cell culture and immortalization. Therefore, the several primary tissue-derived mammary organoid models, which retain more in vivo-like cellular heterogeneity and a more physiologically relevant structure, represent an important advance for understanding the dynamic remodeling in the breast [4–6]. Like immortalized cell lines cultured in vitro, organoids are amenable to advanced imaging and perturbation techniques, providing an exciting venue for studying cell and tissue dynamics. In this review, we will summarize key aspects of mammary gland biology and corresponding organoid models (Fig. 2). Additionally, we will discuss emerging applications where organoids will be transformative for studying both the biology and diseases of the breast. We restrict the scope to organoids sourced from primary human or mouse tissue and stem/progenitor cells that are capable of self-organization and at least limited self-renewal [7].

Establishment of mammary organoid models

Mammary organoids are most commonly established from murine or human mammary epithelial fragments that were micro-dissected or mechanically and enzymatically digested, and then embedded in a reconstituted ECM [6,8]. While murine organoids have been popular due to the availability of powerful genetically engineered mouse models, human-derived organoids sourced from reduction mammoplasty, prophylactic mastectomy, breast biopsy, or resected cancerous tissues are increasingly providing important insight into human biology. Additionally, organoids have been successfully established from dissociated single or reaggregated primary mammary epithelial cells [5,6,9–14]. Importantly, different regions and cell types differ in their organoid-forming potential [9,15,16]. Human pluripotent stem cell (hPSC)-derived mammary-like organoids present the potential for a human model for mammary development and cancer whilst circumventing the requirement for primary tissue [17]. Each of these methods for organoid generation has distinct advantages and disadvantages pertaining to cell compositional control, structural integrity and functionality that must be considered before choosing the right model system (Fig. 3). These models further require additional validation to establish appropriate tissue architecture, cell composition and signaling states [16]. In particular, many of these organoid systems should be benchmarked against primary tissue using techniques like single-cell RNA- or ATAC-seq to better establish in vivo-like cell-type and cell-state diversity [18–21].

Organoid models for mammary gland development and morphogenesis

The earliest stages of mammary development occur prenatally, starting as an ectodermal placode that subsequently gives rise to a rudimentary ductal tree [1]. During puberty, hormonal signaling induces ductal elongation and branching morphogenesis. In the adult, cycling hormone levels accompanying the menstrual cycle cause repeated but small-scale growth and regressions of this ductal network. During pregnancy, however, as hormone levels spike, there is pronounced elaboration of the network prior to lactation. Upon weaning, the network involutes to return to a resting, but somewhat altered ductal structure [22]. Some organoids maintain hormone receptor expression (which is typically lost in 2D culture) over long periods [6,23], but their ability to drive growth and involution in response to hormone levels has not been firmly demonstrated. The primary function of the mammary gland is milk production and delivery to the nipple, and organoid models have begun to recapitulate elements of this process in vitro. While milk protein production by mouse LEP has previously been induced *in vitro* [12], the generation of human organoid models with the correct morphology and cell composition that model lactation and involution has only recently been described [24]. As the expulsion of milk from the gland requires the contractile function of MEP, organoids can additionally be used as a screening system for identifying the Ca²⁺-dependent mechanisms which drive contractility and milk secretion [25].

Organoids undergo dynamic shape changes in response to growth factors and ECM cues, similar to ductal elongation, branching and invasion seen in vivo [8,26,27]. GEMM-derived organoids expressing fluorescent reporters or Cre-drivers from lineage-specific promoters allow cell type-specific genetic manipulation and facilitate spatiotemporal characterization of ductal elongation and branching dynamics in exquisite detail. Fluorescence time lapse microscopy of organoids established that ductal elongation is driven by collective migration of the epithelial cells, reinforced by radial intercalation, and balanced by interfacial tensions along the migrating cell's edges [28,29]. Further, it was shown that cell stratification at the terminal end bud occurs through the loss of apico-basal polarity and oriented cell divisions [30–35].

The three main cell lineages of the mammary epithelium emerge during development and considerable debate surrounds how these lineages are maintained in the adult. Whether there exists a dedicated (as opposed to facultative) bipotent stem cell, the existence of cell trans-differentiation, and the physical and molecular basis for lineage specification remain unclear, in part because in vivo and ex vivo models sometimes provide contradictory results [2,36]. Organoids will be a powerful tool for resolving this debate as they provide precise control on cell composition and microenvironment, and are amenable to live imaging to track cell and tissue dynamics. As pointed out previously, however, more rigorous characterization of cell lineages within different organoid models will be necessary before their potential can be realized in this area.

Organoid models for breast cancer

Breast cancer is a constellation of molecularly distinct diseases. The major molecular subtypes – including luminal A, luminal B, basal, and her2(+) – are also heterogeneous with respect to their gene expression and cellular composition [37]. While recurring mutations in breast cancers are described [38], their impact on cellular signaling, tumor heterogeneity, cancer progression and treatment outcome is not well understood. Organoids present a unique platform to address some of these existing gaps in knowledge. Breast cancer subtype-specific GEMMs exist [39], and organoids derived from these have revealed the importance of collective motility, cell adhesion and matrix interactions on morphogenesis and invasion [10,40–43]. Gain and loss of function studies in organoids, using small molecules and genetics, further established the role of specific genes (e.g. Twist1, Trps1, and E-cadherin) or cell behaviors (cell adhesion and epithelial to mesenchymal transition) in tumor invasion and metastasis [10,42–47].

Recently, organoid biobanks were established from normal and tumor-derived human breast tissue. This resource is amenable to live imaging, as well as genetic and chemical perturbations. These organoids retain their distinct molecular subtypes and elements of the cellular heterogeneity found in vivo [6,11,48]. While they can be cultured for extended periods, it is important to be conscious of molecular and phenotypic drift after passaging, which might alter tumor characteristics or drug response [49]. Human tissue-derived tumor organoids will enable the characterization of dynamic changes to cellular composition, tissue morphology, mechanics, marker expression, and drug susceptibility that coincide with the progression of different tumor types. Additionally, these organoids could allow quantitative analysis of how environmental factors such as aging, injury, hormone therapy, nutrition, and radiation exposure can impact breast cancer risk [47,50]. With sufficient validation, these systems will ultimately be useful guides to develop personalized treatment strategies for patients.

Organoid models for structural dynamics in the breast

The bilayered structure of the mammary epithelium is not only critical to milk secretion and movement, but is a key structural and dynamic barrier to breast cancer progression [51]. Hence, understanding the physical and molecular processes that maintain this structure is an ongoing topic of research. Primary tissue fragments maintain the bilayered structure in vitro, enabling quantitative characterization of changes in tissue organization and shape in response to genetic and microenvironmental perturbations [52,53]. Reconstituted organoids from human reduction mammoplasty tissue established that this bilayered structure arises from a robust program of self-organization [5,54]. Like in many other tissues [55], lineage-specific cell-cell and cell-ECM interfacial tensions in the mammary gland determine the relative mechanical energy of these interfaces and are primarily responsible for driving cells toward the correct luminal or basal position within the tissue. Specifically, the highly unfavorable LEP-ECM interface excludes LEP from the basal tissue layer, while a highly favorable MEP-ECM interface maintains MEP in the basal layer [5].

Breast cancers typically originate in the luminal compartment, and progress through a series of defined structural changes (Fig. 1c), identified as inflection points in risk for breast cancer patients [56]. Less than 30% of ductal carcinoma in situ (DCIS) lesions progress to a more dangerous invasive ductal carcinoma (IDC), characterized by the translocation of transformed LEP past MEP [57,58]. The observation that LEP must translocate past MEP to invade led to the hypothesis that MEP represent a physical barrier to invasion. Using murine organoid models, MEP were recently shown to drastically reduce the dissemination of invasive cancer cells in a dose-dependent manner by undergoing dynamic rearrangement to exclude LEP from the basal layer when they contact ECM [10]. In addition to validating the long-standing hypothesis regarding MEP as structural tumor suppressors, these studies further established the importance of cell dynamics in the process. Understanding the physical and molecular underpinnings of structural maintenance and breakdown may provide key insight into disease progression.

Organoid models for microenvironmental and stromal interactions

Microenvironmental cues from the ECM, stromal and immune cells tightly regulate the mammary epithelium [59]. Alterations to the tissue microenvironment during cancer progression are well established, including changes in ECM stiffness and organization, as well as the composition of the immune cell infiltrate [60]. Previously, patient-derived xenografts and mouse models have been popular for studying these cancer-associated microenvironmental changes, but they provide only minimal control over tumor composition and structure, take time to develop, and are challenging for studying dynamic responses to drugs or stimuli. Many of these challenges can be overcome using organoids that incorporate microenvironmental stimuli from stromal cells and hormone signaling, and are an important frontier for both normal and breast cancer organoids.

Organoids have also proven useful for interrogating the role of epithelial-stromal interactions during glandular morphogenesis [53]. For example, remodeling of the collagenous ECM by the epithelium [61] can reinforce the orientation of collective migration independent of stromal cells, though this is dependent on ECM composition, orientation and adhesion [53,62,63]. Additionally, organoids can be used to study how external signals from hormones or stromal cells can induce ECM changes observed in breast cancer. In one recent example, hormone stimulation of patient-derived organoids identified changes in ECM-related gene expression in patients with BRCA mutations [64]. Considerable effort is being devoted to adding key stromal cell types to co-culture models with epithelial organoids, including fibroblasts, immune cells, and endothelial cells. These efforts are particularly important because the stromal compartment differs in structure and composition between human and mouse (e.g., mouse fibroblasts express estrogen receptors) [1], and cannot be recapitulated in xenograft models. Some progress has been made in this direction. For example, FGF derived from fibroblasts was shown to directly impact epithelial branching and proliferation [29,65–67]. Moving forward, a major goal of organoid-based research will be to add additional stromal cell types, such as immune and endothelial cells, to better mimic the paracrine signaling networks that characterize glandular biology in vivo. A functional immune compartment is also of great interest to the immuno-oncology research community.

Conclusions

Modeling mammary morphogenesis necessitates the correct epithelial structure, ECM organization, cellular composition, as well as multiple stromal cell types. Existing organoid models have begun to bring together the necessary components. These emerging models are shedding light on how signals and forces in the epithelial microenvironment collaborate to generate the complex tissue dynamics observed in vivo, and will undoubtedly be useful in understanding how genomic and microenvironmental changes alter these processes in breast cancer. Looking forward, research should be focused toward the goal of increasing organoid reproducibility, uniformity, complexity and structural accuracy. Further emphasis should be placed on carefully characterizing newly developed organoid models using single cell analysis tools (e.g. scRNA-seq, ATAC-seq, CyTOF, etc.), and benchmarking any findings against in vivo data. These will be important for establishing an optimal human experimental model for studying the signaling and mechanical interactions that occur between mammary epithelial cells, transformed cells, and the surrounding stroma.

Acknowledgements:

Work in the Gartner lab is supported in part by Barbara and Gerson Bakar Foundation and the UCSF Center for Cellular Construction (DBI-1548297), an NSF Science and Technology Center. Z.J.G is a Chan-Zuckerberg BioHub Investigator.

References

1. Parmar H, Cunha GR: Epithelial-stromal interactions in the mouse and human mammary gland in vivo. *Endocr Relat Cancer* 2004, 11:437–458. [PubMed: 15369447]
2. Rios AC, Fu NY, Cursons J, Lindeman GJ, Visvader JE: The complexities and caveats of lineage tracing in the mammary gland. *Breast cancer research : BCR* 2016, 18:116–116. [PubMed: 27887631]
3. Simian M, Bissell MJ: Organoids: A historical perspective of thinking in three dimensions. *The Journal of Cell Biology* 2017, 216:31–40. [PubMed: 28031422]
4. Shamir ER, Ewald AJ: Three-dimensional organotypic culture: experimental models of mammalian biology and disease. *Nat Rev Mol Cell Biol* 2014, 15:647–664. [PubMed: 25237826]
5. Cerchiari AE, Garbe JC, Jee NY, Todhunter ME, Broaders KE, Peehl DM, Desai TA, LaBarge MA, Thomson M, Gartner ZJ: A strategy for tissue self-organization that is robust to cellular heterogeneity and plasticity. *Proceedings of the National Academy of Sciences of the United States of America* 2015, 112:2287–2292. [PubMed: 25633040]
6. Sachs N, de Ligt J, Kopper O, Gogola E, Bounova G, Weeber F, Balgobind AV, Wind K, Gracanin A, Begthel H, et al.: A Living Biobank of Breast Cancer Organoids Captures Disease Heterogeneity. *Cell* 2018, 172:373–386.e310 [PubMed: 29224780] ** The study established a robust method for organoid generation and long-term maintenance from normal and cancerous human breast tissue fragments. The authors demonstrated they can generate hundreds of organoid lines that can be maintained over multiple passages, while retaining hormone receptor expression. These organoids maintain cellular heterogeneity, retain distinct molecular subtypes and exhibit different drug responses that can be used for predicting tumor response to interventions.
7. Lehmann R, Lee CM, Shugart EC, Benedetti M, Charo RA, Gartner Z, Hogan B, Knoblich J, Nelson CM, Wilson KM: Human organoids: a new dimension in cell biology. *Mol Biol Cell* 2019, 30:1129–1137. [PubMed: 31034354]
8. Nguyen-Ngoc KV, Shamir ER, Huebner RJ, Beck JN, Cheung KJ, Ewald AJ: 3D culture assays of murine mammary branching morphogenesis and epithelial invasion. *Methods Mol Biol* 2015, 1189:135–162. [PubMed: 25245692]

9. Linnemann JR, Miura H, Meixner LK, Irmeler M, Kloos UJ, Hirschi B, Bartsch HS, Sass S, Beckers J, Theis FJ, et al.: Quantification of regenerative potential in primary human mammary epithelial cells. *Development* 2015, 142:3239–3251. [PubMed: 26071498]
10. Sirka OK, Shamir ER, Ewald AJ: Myoepithelial cells are a dynamic barrier to epithelial dissemination. *J Cell Biol* 2018, 217:3368–3381 [PubMed: 30061105] ** The study demonstrated that MEP form a dynamic barrier to tumor invasion, providing direct evidence towards a long suggested role for MEP as tumor suppressors. Using reconstituted organoids, containing highly invasive tumor cells and normal MEP, the authors show that MEP can rearrange dynamically to prevent tumor cells from making sustained contact with the ECM. This emphasized the importance of tissue dynamics in cancer progression, for which organoid models are particularly well adept.
11. Dekkers JF, Whittle JR, Vaillant F, Chen HR, Dawson C, Liu K, Geurts M, Herold MJ, Clevers H, Lindeman GJ, et al.: Modeling breast cancer using CRISPR/Cas9-mediated engineering of human breast organoids. *J Natl Cancer Inst* 2019** Described a protocol for engineering human breast organoids using CRISPR/Cas9. The authors further assess post-implantation tumor growth, clonal landscape and drug sensitivity of these engineered organoids. These techniques set the stage for quantitative characterization of single or combination of tumor drivers and signaling perturbations on tumor biology, which is not feasible using tumor-derived models that already contain many accompanying passenger mutations.
12. Jamieson PR, Dekkers JF, Rios AC, Fu NY, Lindeman GJ, Visvader JE: Derivation of a robust mouse mammary organoid system for studying tissue dynamics. *Development* 2017, 144:1065–1071. [PubMed: 27993977]
13. Plaks V, Brenot A, Lawson DA, Linnemann JR, Van Kappel EC, Wong KC, de Sauvage F, Klein OD, Werb Z: Lgr5-expressing cells are sufficient and necessary for postnatal mammary gland organogenesis. *Cell Rep* 2013, 3:70–78. [PubMed: 23352663]
14. Zhang L, Adileh M, Martin ML, Klingler S, White J, Ma X, Howe LR, Brown AMC, Kolesnick R: Establishing estrogen-responsive mouse mammary organoids from single Lgr5(+) cells. *Cellular signalling* 2017, 29:41–51. [PubMed: 27511963]
15. Sokol ES, Miller DH, Breggia A, Spencer KC, Arendt LM, Gupta PB: Growth of human breast tissues from patient cells in 3D hydrogel scaffolds. *Breast Cancer Research* 2016, 18:19. [PubMed: 26926363]
16. Rosenbluth JM, Schackmann RCJ, Gray GK, Selfors LM, Li CM-C, Boedicker M, Kuiken HJ, Richardson A, Brock J, Garber J, et al.: Organoid cultures from normal and cancer-prone human breast tissues preserve complex epithelial lineages. *Nature Communications* 2020, 11:1711* Evaluation of heterogeneity in lineage marker expression and cell type proportions across different tissue sources and culture conditions using CyTOF is a significant step towards benchmarking emerging organoid models against primary tissue. The study further emphasized the importance of organoid models in potentially recapitulating signaling states (e.g. hormone receptor expression) and lineage hierarchies that are quickly lost in 2D cell culture models.
17. Qu Y, Han B, Gao B, Bose S, Gong Y, Wawrowsky K, Giuliano AE, Sareen D, Cui X: Differentiation of Human Induced Pluripotent Stem Cells to Mammary-like Organoids. *Stem cell reports* 2017, 8:205–215. [PubMed: 28132888]
18. Nguyen QH, Pervolarakis N, Blake K, Ma D, Davis RT, James N, Phung AT, Willey E, Kumar R, Jabart E, et al.: Profiling human breast epithelial cells using single cell RNA sequencing identifies cell diversity. *Nature Communications* 2018, 9:2028.
19. Pal B, Chen Y, Vaillant F, Jamieson P, Gordon L, Rios AC, Wilcox S, Fu N, Liu KH, Jackling FC, et al.: Construction of developmental lineage relationships in the mouse mammary gland by single-cell RNA profiling. *Nature Communications* 2017, 8:1627.
20. Bach K, Pensa S, Grzelak M, Hadfield J, Adams DJ, Marioni JC, Khaled WT: Differentiation dynamics of mammary epithelial cells revealed by single-cell RNA sequencing. *Nature Communications* 2017, 8:2128.
21. Pervolarakis N, Sun P, Gutierrez G, Nguyen QH, Jhutti D, XY Zheng G, Nemec CM, Dai X, Watanabe K, Kessenbrock K: Integrated Single-Cell Transcriptomics and Chromatin Accessibility Analysis Reveals Novel Regulators of Mammary Epithelial Cell Identity. *CELL-REPORTS-D-19-03157*. 2019.

22. Russo J, Rivera R, Russo IH: Influence of age and parity on the development of the human breast. *Breast Cancer Res Treat* 1992, 23:211–218. [PubMed: 1463860]
23. Jardé T, Lloyd-Lewis B, Thomas M, Kendrick H, Melchor L, Bougaret L, Watson PD, Ewan K, Smalley MJ, Dale TC: Wnt and Neuregulin1/ErbB signalling extends 3D culture of hormone responsive mammary organoids. *Nature Communications* 2016, 7:13207.
24. Sumbal J, Chiche A, Charifou E, Koledova Z, Li H: Primary mammary organoid model of lactation and involution. *Front. Cell Dev. Biol* 2020, 8.
25. Stewart TA, Davis FM: A Primary Cell and Organoid Platform for Evaluating Pharmacological Responses in Mammary Epithelial Cells. *ACS Pharmacology & Translational Science* 2020* Characterized the dynamics of lineage-specific Ca²⁺ response in mammary organoids to various ligands, further demonstrating how these dynamic processes might differ in 3D compared to 2D. This study shows how the use of genetically coded reporters in engineered organoids enables quantitative characterization of signaling dynamics in the mammary gland, which are structural context-dependent and have been largely inaccessible using traditional cell culture models.
26. Darcy KM, Black JD, Hahm HA, Ip MM: Mammary organoids from immature virgin rats undergo ductal and alveolar morphogenesis when grown within a reconstituted basement membrane. *Exp Cell Res* 1991, 196:49–65. [PubMed: 1879471]
27. Barcellos-Hoff MH, Aggeler J, Ram TG, Bissell MJ: Functional differentiation and alveolar morphogenesis of primary mammary cultures on reconstituted basement membrane. *Development* 1989, 105:223–235. [PubMed: 2806122]
28. Neumann NM, Perrone MC, Veldhuis JH, Huebner RJ, Zhan H, Devreotes PN, Brodland GW, Ewald AJ: Coordination of Receptor Tyrosine Kinase Signaling and Interfacial Tension Dynamics Drives Radial Intercalation and Tube Elongation. *Developmental Cell* 2018, 45:67–82.e66 [PubMed: 29634937] * The study used murine organoid models to demonstrate how cell migration and radial intercalation during tube elongation is achieved through a coordination of interfacial forces and biochemical signaling at a tissue level. Time lapse imaging using fluorescent reporters allowed quantification of cell shape, position and motility over time, that helped develop a physical model for explaining the dynamics of tube elongation based on a balance of mechanical stress and cell protrusion in the epithelium.
29. Chatterjee SJ, Halaoui R, Deagle RC, Rejon C, McCaffrey L: Numb regulates cell tension required for mammary duct elongation. *Biol Open* 2019, 8.
30. Ewald AJ, Brenot A, Duong M, Chan BS, Werb Z: Collective epithelial migration and cell rearrangements drive mammary branching morphogenesis. *Dev Cell* 2008, 14:570–581. [PubMed: 18410732]
31. Ewald AJ, Huebner RJ, Palsdottir H, Lee JK, Perez MJ, Jorgens DM, Tauscher AN, Cheung KJ, Werb Z, Auer M: Mammary collective cell migration involves transient loss of epithelial features and individual cell migration within the epithelium. *J Cell Sci* 2012, 125:2638–2654. [PubMed: 22344263]
32. Huebner RJ, Lechler T, Ewald AJ: Developmental stratification of the mammary epithelium occurs through symmetry-breaking vertical divisions of apically positioned luminal cells. *Development (Cambridge, England)* 2014, 141:1085–1094.
33. Xu W, Gulvady AC, Goreczny GJ, Olson EC, Turner CE: Paxillin-dependent regulation of apical-basal polarity in mammary gland morphogenesis. *Development* 2019, 146.
34. Kessenbrock K, Smith P, Steenbeek SC, Pervolarakis N, Kumar R, Minami Y, Goga A, Hinck L, Werb Z: Diverse regulation of mammary epithelial growth and branching morphogenesis through noncanonical Wnt signaling. *Proceedings of the National Academy of Sciences* 2017:201701464.
35. Nguyen-Ngoc K-V, Silvestri VL, Georgess D, Fairchild AN, Ewald AJ: Mosaic loss of non-muscle myosin IIA and IIB is sufficient to induce mammary epithelial proliferation. *Journal of Cell Science* 2017, 130:3213–3221. [PubMed: 28821574]
36. Seldin L, Le Guelte A, Macara IG: Epithelial plasticity in the mammary gland. *Current opinion in cell biology* 2017, 49:59–63. [PubMed: 29232628]
37. Polyak K: Heterogeneity in breast cancer. *The Journal of Clinical Investigation* 2011, 121:3786–3788. [PubMed: 21965334]

38. Cancer Genome Atlas Research N, Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, Shmulevich I, Sander C, Stuart JM: The Cancer Genome Atlas Pan-Cancer analysis project. *Nature genetics* 2013, 45:1113–1120. [PubMed: 24071849]
39. Hutchinson JN, Muller WJ: Transgenic mouse models of human breast cancer. *Oncogene* 2000, 19:6130–6137. [PubMed: 11156526]
40. Nguyen-Ngoc K-V, Cheung KJ, Brenot A, Shamir ER, Gray RS, Hines WC, Yaswen P, Werb Z, Ewald AJ: ECM microenvironment regulates collective migration and local dissemination in normal and malignant mammary epithelium. *Proceedings of the National Academy of Sciences* 2012, 109:E2595–E2604.
41. Cheung Kevin J, Gabrielson E, Werb Z, Ewald Andrew J: Collective Invasion in Breast Cancer Requires a Conserved Basal Epithelial Program. *Cell* 2013, 155:1639–1651. [PubMed: 24332913]
42. Shamir ER, Pappalardo E, Jorgens DM, Coutinho K, Tsai WT, Aziz K, Auer M, Tran PT, Bader JS, Ewald AJ: Twist1-induced dissemination preserves epithelial identity and requires E-cadherin. *J Cell Biol* 2014, 204:839–856. [PubMed: 24590176]
43. Padmanaban V, Krol I, Suhail Y, Szczerba BM, Aceto N, Bader JS, Ewald AJ: E-cadherin is required for metastasis in multiple models of breast cancer. *Nature* 2019, 573:439–444. [PubMed: 31485072]
44. Cornelissen LM, Drenth AP, van der Burg E, de Bruijn R, Pritchard CEJ, Huijbers IJ, Zwart W, Jonkers J: TRPS1 acts as a context-dependent regulator of mammary epithelial cell growth/differentiation and breast cancer development. *Genes Dev* 2019.
45. Michalak EM, Milevskiy MJG, Joyce RM, Dekkers JF, Jamieson PR, Pal B, Dawson CA, Hu Y, Orkin SH, Alexander WS, et al.: Canonical PRC2 function is essential for mammary gland development and affects chromatin compaction in mammary organoids. *PLoS biology* 2018, 16:e2004986–e2004986. [PubMed: 30080881]
46. Villeneuve C, Lagoutte E, de Plater L, Mathieu S, Manneville J-B, Maître J-L, Chavrier P, Rossé C: aPKC α triggers basal extrusion of luminal mammary epithelial cells by tuning contractility and vinculin localization at cell junctions. *Proceedings of the National Academy of Sciences* 2019, 116:24108–24114.
47. Williams KE, Lemieux GA, Hassis ME, Olshen AB, Fisher SJ, Werb Z: Quantitative proteomic analyses of mammary organoids reveals distinct signatures after exposure to environmental chemicals. *Proc Natl Acad Sci U S A* 2016, 113:E1343–1351. [PubMed: 26903627]
48. Padmanaban V, Tsehay Y, Cheung KJ, Ewald AJ, Bader JS: Between-tumor and within-tumor heterogeneity in invasive potential. *PLoS Comput Biol* 2020, 16:e1007464. [PubMed: 31961880]
49. Goldhammer N, Kim J, Timmermans-Wielenga V, Petersen OW: Characterization of organoid cultured human breast cancer. *Breast Cancer Research* 2019, 21:141. [PubMed: 31829259]
50. Hu JL, Todhunter ME, LaBarge MA, Gartner ZJ: Opportunities for organoids as new models of aging. *J Cell Biol* 2018, 217:39–50. [PubMed: 29263081]
51. Sternlicht MD, Kedeshian P, Shao ZM, Safarians S, Barsky SH: The human myoepithelial cell is a natural tumor suppressor. *Clin Cancer Res* 1997, 3:1949–1958. [PubMed: 9815584]
52. Tanos T, Sflomos G, Echeverria PC, Ayyanan A, Gutierrez M, Delaloye JF, Raffoul W, Fiche M, Dougall W, Schneider P, et al.: Progesterone/RANKL is a major regulatory axis in the human breast. *Sci Transl Med* 2013, 5:182ra155.
53. Nguyen-Ngoc KV, Ewald AJ: Mammary ductal elongation and myoepithelial migration are regulated by the composition of the extracellular matrix. *J Microsc* 2013, 251:212–223. [PubMed: 23432616]
54. Chanson L, Brownfield D, Garbe JC, Kuhn I, Stampfer MR, Bissell MJ, LaBarge MA: Self-organization is a dynamic and lineage-intrinsic property of mammary epithelial cells. *Proc Natl Acad Sci U S A* 2011, 108:3264–3269. [PubMed: 21300877]
55. Brodland GW: The Differential Interfacial Tension Hypothesis (DITH): a comprehensive theory for the self-rearrangement of embryonic cells and tissues. *J Biomech Eng* 2002, 124:188–197. [PubMed: 12002128]
56. Allred DC: Ductal carcinoma in situ: terminology, classification, and natural history. *J Natl Cancer Inst Monogr* 2010, 2010:134–138. [PubMed: 20956817]

57. Sanders ME, Schuyler PA, Dupont WD, Page DL: The natural history of low-grade ductal carcinoma in situ of the breast in women treated by biopsy only revealed over 30 years of long-term follow-up. *Cancer* 2005, 103:2481–2484. [PubMed: 15884091]
58. Sagara Y, Julia W, Golshan M, Toi M: Paradigm Shift toward Reducing Overtreatment of Ductal Carcinoma In Situ of Breast. *Frontiers in Oncology* 2017, 7.
59. Schedin P, Keely PJ: Mammary gland ECM remodeling, stiffness, and mechanosignaling in normal development and tumor progression. *Cold Spring Harb Perspect Biol* 2011, 3:a003228. [PubMed: 20980442]
60. Place AE, Jin Huh S, Polyak K: The microenvironment in breast cancer progression: biology and implications for treatment. *Breast Cancer Res* 2011, 13:227. [PubMed: 22078026]
61. Gomes AM, Bhat R, Correia AL, Mott JD, Ilan N, Vlodavsky I, Pavao MS, Bissell M: Mammary Branching Morphogenesis Requires Reciprocal Signaling by Heparanase and MMP-14. *J Cell Biochem* 2015, 116:1668–1679. [PubMed: 25735873]
62. Brownfield DG, Venugopalan G, Lo A, Mori H, Tanner K, Fletcher DA, Bissell MJ: Patterned collagen fibers orient branching mammary epithelium through distinct signaling modules. *Curr Biol* 2013, 23:703–709. [PubMed: 23562267]
63. Beck JN, Singh A, Rothenberg AR, Elisseeff JH, Ewald AJ: The independent roles of mechanical, structural and adhesion characteristics of 3D hydrogels on the regulation of cancer invasion and dissemination. *Biomaterials* 2013, 34:9486–9495. [PubMed: 24044993]
64. Davaadelger B, Choi M-R, Singhal H, Clare SE, Khan SA, Kim JJ: BRCA1 mutation influences progesterone response in human benign mammary organoids. *Breast Cancer Research* 2019, 21:124. [PubMed: 31771627]
65. Koledova Z: 3D Coculture of Mammary Organoids with Fibrospheres: A Model for Studying Epithelial–Stromal Interactions During Mammary Branching Morphogenesis. In *3D Cell Culture: Methods and Protocols*. Edited by Koledova Z: Springer New York; 2017:107–124.
66. Sumbal J, Koledova Z: FGF signaling in mammary gland fibroblasts regulates multiple fibroblast functions and mammary epithelial morphogenesis. *Development* 2019, 146:dev185306 [PubMed: 31699800] ** The study characterizes the effect of fibroblasts on the branching morphogenesis of mouse mammary organoids through FGF signaling. This is one of the first studies developing co-culture models of primary-derived mammary organoid with non-epithelial stromal cells.
67. Koledova Z, Zhang X, Streuli C, Clarke RB, Klein OD, Werb Z, Lu P: SPRY1 regulates mammary epithelial morphogenesis by modulating EGFR-dependent stromal paracrine signaling and ECM remodeling. *Proc Natl Acad Sci U S A* 2016, 113:E5731–5740. [PubMed: 27621461]

Figure 1:

a. The mammary gland undergoes extensive remodeling during a woman's lifetime, particularly during pubertal development and pregnancy. During puberty, the rudimentary epithelial tree elongates and branches to infiltrate the stroma (mainly adipocytic in mouse and fibroblastic in human), resulting in a lobular tree. Further maturation of the gland occurs during pregnancy, when hormones drive the growth of the terminal lobular ductal units into alveoli, wherein the luminal epithelial cells can differentiate into milk-secreting cells for lactation. After weaning, the process of involution remodels the mammary gland, including the collapse of the alveoli and extensive apoptosis. **b.** The structure of the mammary gland comprises a bilayered epithelium surrounded by a variety of stromal cell types. How tissue homeostasis is maintained remains an active area of research, including questions about

lineage specification, tissue composition and structure, differences in the ducts vs alveoli as well as how they change with time and age, autocrine and paracrine signaling, and response to stimuli (e.g. hormones, drugs). **c.** Histologically, breast cancer progresses through distinct stages, beginning with the filling of the lumen in ductal carcinoma in situ (DCIS), to invasion of the proliferative cancerous cells past the myoepithelium, and finally to metastasis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2:

Many organoid models of the mammary epithelium have been established to study the biology of the mammary gland, both in development and tissue homeostasis (a), as well as breast cancer progression (b). We also highlighted emerging models that could leverage the capabilities of organoid cultures to address active questions in the mammary field (c).

Figure 3:
a. Mammary organoids have been established from various tissue sources including human and murine primary tissue, breast tumors and stem cells, which are processed as tissue fragments, cellular aggregates or single cells. Each method for organoid preparation has its distinct benefits and limitations as summarized here, which must be considered when choosing the appropriate organoid model for the study. Darker colors represent higher similarity to the corresponding feature of the in vivo gland. **b.** Representative images of digested human mammary tissue (**i**), organoids derived from tissue fragments (**ii**) and reconstituted organoids made by aggregating mammary epithelial cells (**iii**). These methods have been commonly used for mammary organoid preparation. The tissue and organoids have been stained for luminal (Keratin-19) and myoepithelial (Keratin-14) markers to highlight incorporation of multiple cell types and maintenance of the in vivo-like bilayered tissue structure. Scale bar: 100 μ m.