Responsiveness of *Aromatoleum aromaticum* EbN1^T to Lignin-Derived Phenylpropanoids

Jannes Vagts,^a Kristin Kalvelage,^a Arne Weiten,^a Ramona Buschen,^a Julian Gutsch,^a Sabine Scheve,^a Lars Wöhlbrand,^a Stefan Diener,^b Heinz Wilkes,^b Michael Winkhofer,^{c,d} Ralf Rabus^a

^aGeneral and Molecular Microbiology, Institute for Chemistry and Biology of the Marine Environment (ICBM), Carl von Ossietzky University of Oldenburg, Oldenburg, Germany

^bOrganic Geochemistry, Institute for Chemistry and Biology of the Marine Environment (ICBM), Carl von Ossietzky University of Oldenburg, Oldenburg, Germany

^cResearch Center Neurosensory Science, Carl von Ossietzky University of Oldenburg, Oldenburg, Germany

^dSensory Biology of Animals, Institute of Biology and Environmental Sciences (IBU), Carl von Ossietzky University of Oldenburg, Oldenburg, Germany

ABSTRACT The betaproteobacterial degradation specialist *Aromatoleum aromaticum* EbN1^T utilizes several plant-derived 3-phenylpropanoids coupled to denitrification. *In vivo* responsiveness of *A. aromaticum* EbN1^T was studied by exposing nonadapted cells to distinct pulses (spanning 100 μ M to 0.1 nM) of 3-phenylpropanoate, cinnamate, 3-(4-hydroxyphenyl)propanoate, or *p*-coumarate. Time-resolved, targeted transcript analyses via quantitative reverse transcription-PCR of four selected 3-phenylpropanoid genes revealed a response threshold of 30 to 50 nM for *p*-coumarate and 1 to 10 nM for the other three tested 3-phenylpropanoids. At these concentrations, transmembrane effector equilibration is attained by passive diffusion rather than active uptake via the ABC transporter, presumably serving the studied 3-phenylpropanoids as well as benzoate. Highly substrate-specific enzyme formation (EbA5316 to EbA5321 [EbA5316-21]) for the shared peripheral degradation pathway putatively involves the predicted TetR-type transcriptional repressor PprR. Accordingly, relative transcript abundances of *ebA5316-21* are lower in succinate- and benzoate-grown wild-type cells than in an unmarked in-frame $\Delta pprR$ mutant. In *trans*-complementation of *pprR* into the $\Delta pprR$ background restored wild-type-like transcript levels. When adapted to *p*-coumarate, the three genotypes had relative transcript abundances similar to those of *ebA5316-21* despite a significantly longer lag phase of the *pprR*-complemented mutant ($\sim$ 100-fold higher *pprR* transcript level than the wild type). Notably, transcript levels of *ebA5316-21* were $\sim$ 10- to 100-fold higher in *p*-coumarate- than succinate- or benzoate-adapted cells across all three genotypes. This indicates the additional involvement of an unknown transcriptional regulator. Furthermore, physiological, transcriptional, and (aromatic) acyl-coenzyme A ester intermediate analyses of the wild type and $\Delta pprR$ mutant grown with binary substrate mixtures suggest a mode of catabolite repression of superior order to PprR.

IMPORTANCE Lignin is a ubiquitous heterobiopolymer built from a suite of 3-phenylpropanoid subunits. It accounts for more than 30% of the global plant dry material, and lignin-related compounds are increasingly released into the environment from anthropogenic sources, i.e., by wastewater effluents from the paper and pulp industry. Hence, following biological or industrial decomplexation of lignin, vast amounts of structurally diverse 3-phenylpropanoids enter terrestrial and aquatic habitats, where they serve as substrates for microbial degradation. This raises the question of what signaling systems environmental bacteria employ to detect these nutritionally attractive compounds and to adjust their catabolism accordingly. Moreover, determining *in vivo* response thresholds of an anaerobic degradation specialist such as *A. aromaticum* EbN1^T for these aromatic compounds provides insights into the environmental fate of the latter, i.e., when they could escape biodegradation due to too low ambient concentrations.

Citation Vagts J, Kalvelage K, Weiten A, Buschen R, Gutsch J, Scheve S, Wöhlbrand L, Diener S, Wilkes H, Winkhofer M, Rabus R. 2021. Responsiveness of *Aromatoleum aromaticum* EbN1^T to lignin-derived phenylpropanoids. *Appl Environ Microbiol* 87:e03140-20. <https://doi.org/10.1128/AEM.03140-20>.

Editor Rebecca E. Parales, University of California, Davis

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Ralf Rabus, rabus@icbm.de.

Received 23 December 2020

Accepted 9 March 2021

Accepted manuscript posted online 19 March 2021

Published 11 May 2021

KEYWORDS anaerobic degradation, phenylpropanoids, regulation, responsiveness, sensory system, deletion mutation, transcript profiling, physiology, diauxie, (aromatic) acyl-CoA ester, *Aromatoleum aromaticum* EbN1^T

Lignin accounts for more than 30% of the global plant dry mass (1), providing compressive strength and protection against pathogens (2, 3). The high heterogeneity of this biopolymer arises from the structural diversity of its phenylpropanoid building blocks, which, moreover, are interconnected by various bond types at multiple positions (4). Due to its global abundance and renewability, lignocellulosic biomass is regarded as a promising alternative to fossil fuels in the production of biofuels (3) and industrial bulk chemicals (5). Hence, pronounced research interest is focused on the identification of novel bacterial enzymes for depolymerization of lignin and respective metabolic engineering (3, 4). In addition to its natural abundance, millions of tons of lignin and lignin-related products are presently released into the environment through waste effluents of the paper and pulping industries (6). White-rot basidiomycetes were found to be mainly accountable for natural lignin depolymerization, employing different ligninolytic enzymes (7–9). However, recent studies showed that bacterial consortia also play a role in degradation of heteropolymeric lignin (4, 10–12). According to the so-called rhizosphere effect, phenylpropanoids are exuded by plant roots (13, 14), enriching the complex soluble organic matter in the rhizosphere soil. This may be rationalized by phenylpropanoids belonging to the dominant metabolites of root core cell types (15). Furthermore, members of the intestinal microbiome of animals can convert dietary flavonoids into phenylpropanoids (16), which may be excreted into the environment. Thus, phenylpropanoids enter terrestrial and aquatic habitats from a variety of sources. Apart from their utilization as a carbon source, recent studies have proven environment-derived 3-phenylpropanoids (i.e., *p*-coumarate) to also play a role in quorum-sensing mechanisms, allowing bacteria to communicate intercellularly and to adjust global gene expression according to environmental changes and cell density (1, 17).

Many different soil and sediment bacteria have been shown to harbor the metabolic capacity to degrade 3-phenylpropanoids under oxic or anoxic conditions (18–21). The betaproteobacterial degradation specialist *Aromatoleum aromaticum* EbN1^T is a well-studied model organism capable of oxidizing >20 different aromatic growth compounds completely to CO₂ coupled to denitrification (22–24). Differential proteogenomic analyses revealed a complex catabolic network composed of distinct peripheral degradation pathways subject to highly substrate-specific regulation (23, 25–27). The peripheral degradation pathway for 3-phenylpropanoids in *A. aromaticum* EbN1^T comprises 5 enzymes and one predicted solute binding protein (SBP), with the coding genes (*ebA5316* to *ebA5321* [*ebA5316-21*]; *pprA* to *pprF* [*pprA-F*]) arranged in an operon-like structure (Fig. 1) (20, 23). The latter is part of a larger cluster comprising genes for an ABC-type transport system and the central anaerobic benzoyl-coenzyme A (CoA) pathway (20).

Substrate-specific pathway regulation in *A. aromaticum* EbN1^T is proposed to be mediated on the transcriptional level by a variety of one- or two-component signaling systems (23), with recent studies pointing to remarkably high *in vivo* sensitivities in the low-nanomolar range (28, 29). Gene expression for anaerobic degradation of *p*-ethylphenol was shown to be regulated by the σ^{54} -dependent one-component transcriptional activator EtpR (30). The two-component system PcrSR was demonstrated to be essential for the expression of genes encoding enzymes of the anaerobic *p*-cresol catabolism (29). For the substrate-dependent regulation of the peripheral 3-phenylpropanoid degradation pathway, the one-component system EbA5314 (here renamed PprR, affiliated with the TetR family) was proposed to act as a transcriptional repressor (23). Similar modes of regulation have been identified for the degradation of *p*-cymene in *Pseudomonas putida* (CymR) (31) and *p*-coumarate in *Rhodopseudomonas palustris* (CouR) and *Acinetobacter* species (21, 32, 33). Based on these studies, one may assume that the catabolic intermediate *p*-coumaroyl-CoA binds to the C-terminal TetR domain of PprR, which thereupon disengages from the promoter upstream region, allowing

FIG 1 Scheme of the proposed transcriptional regulation of anaerobic 3-phenylpropanoid degradation in the denitrifying bacterium *Aromatoleum aromaticum* EbN1^T. Gene expression is proposed to be mediated by the predicted one-component transcriptional repressor PprR. PprR is assumed to bind shared degradation intermediates of different 3-phenylpropanoids to its N-terminal TetR domain. Upon effector binding, PprR disengages from the promoter region of the 3-phenylpropanoid catabolic gene cluster, allowing initiation of transcription (*ebA5316-21*; *pprA-F*). Compound names are the following: 1, benzoate; 2, 3-phenylpropanoate or 3-(4-hydroxyphenyl)propanoate; 3, cinnamate or *p*-coumarate; 4, hydrocinnamoyl-CoA or 3-(4-hydroxyphenyl)propanoyl-CoA; 5, cinnamoyl-CoA or *p*-coumaroyl-CoA; 6, 3-hydroxy-3-phenylpropanoyl-CoA or 3-hydroxy-3-(4-hydroxyphenyl)propanoyl-CoA; 7, benzoylacetate or 4-hydroxybenzoyl-CoA; 8, 4-hydroxybenzoyl-CoA; 9, benzoyl-CoA; 10, cyclohexa-1,5-diene-1-carboxyl-CoA. R, H or OH. green—, presence or absence of double bond. OM, outer membrane; PP, periplasm; IM, inner membrane.

DNA-dependent RNA polymerase to transcribe the 3-phenylpropanoid gene cluster (Fig. 1).

The present study investigates the transcriptional regulation of 3-phenylpropanoid utilization by *A. aromaticum* EbN1^T on several levels. To begin with, the *in vivo* responsiveness towards four different 3-phenylpropanoids across a concentration range from 100 μ M down to 0.1 nM was determined by means of time-resolved, targeted transcript profiling (Fig. 2). Differential proteomics allocated a nearby encoded ABC transporter to 3-phenylpropanoid uptake (Fig. 3). However, theoretical considerations suggest this system does not play a role in threshold concentrations, where transmembrane effector equilibrium is attained by diffusion (see Fig. 8). Furthermore, we used unmarked in-frame deletion to verify the proposed role of PprR as an appendant transcriptional repressor: a Δ *pprR* mutant showed an increased transcript level of 3-phenylpropanoid genes compared to the wild type when anaerobically grown with succinate or benzoate (see Fig. 5A and B). In accordance with this, in *trans*-complementation of *pprR* in the Δ *pprR* background restored wild-type-like transcript levels. When grown with *p*-coumarate, equal transcript levels were observed for all three genotypes (see Fig. 5C). Finally, we provide

FIG 2 Time-resolved, quantitative transcript profiles of *A. aromaticum* EbN1^T in response to different extracellular effector concentrations. Tested 3-phenylpropanoid effectors were 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, cinnamate, and *p*-coumarate. The selected transcripts represent genes (Fig. 1) coding for enzymes involved in the anaerobic degradation of 3-phenylpropanoids (*ebA5316*, *ebA5317*, *ebA5319*, and *ebA5321*). Relative transcript abundances were determined by means of qRT-PCR, with the time point of 5 min prior to effector addition serving as a reference. Each data point is based on 3 biological replicates with 5 technical replicates analyzed for each. Growth data of the cultures providing the RNA samples for all tested conditions are shown in detail in Fig. S1 to S4.

evidence that 3-phenylpropanoid utilization is, besides being controlled by PprR, also subject to catabolite repression, revealing a multilayered transcriptional regulation of the respective gene cluster (see Fig. 6 and 7).

RESULTS

In vivo response threshold toward 3-phenylpropanoids. To assess the response threshold for structural variants of 3-phenylpropanoids, first benzoate-adapted cells of *A. aromaticum* EbN1^T were anaerobically grown with a limiting supply of benzoate (1 mM). Upon its complete consumption, a single pulse of one 3-phenylpropanoid was immediately administered per individual culture, yielding initial concentrations from 100 μ M down to 0.1 nM. Tested 3-phenylpropanoids were 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, cinnamate, and *p*-coumarate. Highly reproducible growth curves with sampling time points for each of the eight tested effector concentrations

FIG 3 Relationship between 3-phenylpropanoid and benzoate utilization by *A. aromaticum* EbN1^T in terms of uptake, catabolism, and transcriptional regulation. (A) Structural and functional representation of the chromosomal locus (27) comprising the genes related to anaerobic catabolism of 3-phenylpropanoids and benzoate (20). Note the proposed shared role of the predicted ABC-type transporter for the uptake of these aromatic carboxylates. (B) Differential profiles of the encoded proteins are in accord with the proposed polyspecificity of this ABC uptake system. Proteomic profiling was based on membrane protein-enriched as well as soluble protein fractions. In case of protein identification in both fractions, the one yielding the highest MASCOT score is shown here; further details are provided in Table S7.

and a negative control (no addition of effector) are provided in Fig. S1 to S4 in the supplemental material and fold changes of transcript abundances in Tables S1 to S4. For transcript profiling, samples were retrieved 5 min prior to as well as 5, 15, 30, 60, and 120 min after the pulse (zoom-ins in Fig. S1 to S4). Differential expression of the four selected genes was related to the individual expression levels directly prior to the effector pulse. Analyzed genes were selected according to their location within the 3-phenylpropanoid gene cluster, covering its start (*ebA5316*), middle (*ebA5317* and *ebA5319*), and end (*ebA5321*).

For all tested 3-phenylpropanoids, only the highest concentration of 100 μ M was sufficient to support an increase in optical density. At all other tested concentrations, the optical density of the cultures remained constant after addition of the respective effector. Time-resolved transcript profiles revealed throughout a correlation between transcriptional response time and the gene position within the operon-like structure (Fig. 2). Accordingly, genes near the end of the gene cluster had delayed and lower transcript abundances than genes located closer to the putative transcriptional start. Moreover, an apparent correlation between effector concentration and overall response intensity was observed.

Upon addition of 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, or cinnamate at final concentrations of ≥ 10 nM, an increase in relative *ebA5316* transcript abundance was already observed after 5 min (≥ 30 nM for *p*-coumarate). Altogether, the maximal

transcript abundance increase of ~300-fold was observed for *ebA5317* in response to 100 μ M 3-phenylpropanoate. For the three other 3-phenylpropanoids, the maxima were likewise recorded at 100 μ M effector: 100-fold for cinnamate and 135-fold for *p*-coumarate and 3-(4-hydroxyphenyl)propanoate. The lowest effector concentration allowing for a transcriptional response was 10 nM in the case of 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, and cinnamate. Therefore, an apparent *in vivo* response threshold between 1 nM and 10 nM is obvious for these three effectors. In the case of *p*-coumarate, the lowest concentration yielding a transcriptional response was 50 nM, suggesting a higher *in vivo* response threshold in the range of 30 to 50 nM.

3-Phenylpropanoid uptake presumably via ABC transporter. The first gene in the phenylpropanoid gene cluster of *A. aromaticum* EbN1^T, *ebA5316*, encodes a solute-binding protein (SBP) of ABC-type transporters that shows highest similarity to SBPs from various betaproteobacteria (Table S5). Furthermore, comparison to experimentally verified aromatic compound-binding SBPs from *R. palustris* revealed higher similarity of EbA5316 to *p*-coumarate-binding RPA1789 (identity 49%) than to benzoate- or *p*-hydroxybenzoate-binding SBPs (RPA0668, 24% identity; RPA4029, 25% identity) (34) (Table S6). This suggests EbA5316 acts as a 3-phenylpropanoid-specific SBP serving an ABC-type transport system. Notably, such a system is encoded in direct proximity of the 3-phenylpropanoid gene cluster, comprising permease (EbA5304/6) and ATP-binding subunits (EbA5307/9) (Fig. 3A). Since an SBP (EbA5303) is encoded directly upstream of *ebA5304-9* on the same strand, this system was previously assigned to benzoate uptake (20). In accordance with this, EbA5303 is markedly more similar to benzoate-binding RPA0668 (identity, 54%) than to *p*-coumarate-binding RPA1789 (identity, 26%) of *R. palustris*. Finally, this ABC transporter (EbA5304-9), together with the two SBPs (EbA5316 and EbA5303 [EbA5316/03]), is encoded between the gene clusters for anaerobic catabolism of 3-phenylpropanoids and benzoate (Fig. 3A).

To scrutinize the implicated polyspecificity of this ABC transporter, the membrane protein-enriched fractions of 3-(4-hydroxyphenyl)propanoate-, benzoate-, and acetate-adapted cells of *A. aromaticum* EbN1^T were analyzed comparatively. The predicted ABC transporter permease (EbA5304/6) as well as ATP-binding subunits (EbA5307/9) were similarly abundant in the case of the two aromatic substrates but not with acetate (Fig. 3B, Table S7). While the benzoate-specific SBP (EbA5303) showed similar abundance with both aromatic substrates, the 3-phenylpropanoid-specific SBP (EbA5316) was abundantly formed only in 3-(4-hydroxyphenyl)propanoate-adapted cells. Thus, the predicted permease (EbA5304/6) apparently accommodates both types of aromatic carboxylates, while the two SBPs should confer substrate specificity (EbA5316 for 3-phenylpropanoids and EbA5303 for benzoate).

Generation of Δ *pprR* and *pprR*-complemented mutants. The predicted one-component transcriptional repressor PprR is encoded directly upstream of the 3-phenylpropanoid gene cluster (Fig. 3A) on the complementary strand. To validate its function in repressing expression of *ebA5316-21*, an unmarked, in-frame Δ *pprR* mutant (Δ *pprR*) was generated. In this mutant, the start and stop codons of *pprR* as well as a BamHI restriction site (6 nucleotides) were preserved to maintain the reading frame (Fig. S5A). Accordingly, gene-specific primers for *pprR* did not yield a PCR product in the deletion mutant, whereas primers hybridizing up- and downstream of the knockout region resulted in a small 316-bp region compared to a large 1,207-bp product in the wild type (Fig. S5B). Deletion of *pprR* led to derepression of the 3-phenylpropanoid gene expression in cells anaerobically grown with benzoate (see below). The Δ *pprR* mutant was in *trans*-complemented (Δ *pprR*-compl.) using a *pprR*-containing pBBR1MCS-2 broad-host-range vector (Fig. S5B), and this complementation restored repression of the 3-phenylpropanoid-specific gene expression in benzoate-grown cells (see below). Moreover, the correctness of both mutants was verified by nucleotide sequencing.

PprR involved in substrate-specific expression of 3-phenylpropanoid genes. To investigate the Δ *pprR*-specific phenotype, all three genotypes (wild type, Δ *pprR*, and Δ *pprR*-compl.) were anaerobically grown with succinate, benzoate, or *p*-coumarate (Fig. 4A to C). Furthermore, for all three genotypes and substrate conditions (at half-

FIG 4 Characterization of the wild type (blue), $\Delta pprR$ mutant (pink), and *pprR*-complemented mutant (gray-blue) of *A. aromaticum* EbN1^T. All three genotypes were grown with 8 mM succinate (A), 4 mM benzoate (B), or 2 mM *p*-coumarate (C). (D) Increased transcript level of *pprR* in the *pprR*-complemented mutant (gray-blue) compared to the wild type (blue) of *A. aromaticum* EbN1^T; transcript levels are normalized against *gyrB*.

maximal optical density, or $\frac{1}{2}OD_{max}$), abundance profiles of transcripts (via quantitative reverse transcription-PCR [qRT-PCR], normalizing against *gyrB*) (Fig. 5) and proteins (via shotgun proteomics; Fig. S6) of the 3-phenylpropanoid gene cluster were determined.

(i) Succinate and benzoate. All three genotypes showed highly similar growth behavior, maximum growth rates (μ_{max}), and substrate consumption profiles (Fig. 4AB). In accordance with the prediction that PprR acts as a transcriptional repressor of the 3-phenylpropanoid gene cluster, transcript abundance of the respective genes was markedly increased (derepressed) in the $\Delta pprR$ mutant compared to the wild type (Fig. 5AB). The strongest relative increase in transcript abundance (10-fold) was observed for *ebA5316*, which is localized closest to the putative transcriptional start. Analogous increases in relative transcript abundance (2.3- to 4.1-fold) were observed for *ebA5317*, *ebA5318*, and *ebA5319*. In the case of *ebA5320* and *ebA5321*, located at the end of the 3-phenylpropanoid gene cluster, such an increase was only observed for succinate-grown cells. The described effects on gene expression in the $\Delta pprR$ mutant could be canceled by in *trans*-complementation of the *pprR* gene, restoring relative transcription to levels similar to those of the wild type, except for *ebA5316* (2.8-fold increase with benzoate).

(ii) *p*-Coumarate. μ_{max} as well as OD_{max} were similar for all three genotypes (Fig. 4C). However, the *pprR*-complemented mutant had a significantly longer lag phase than the wild type and $\Delta pprR$ mutant. Correspondingly, complete substrate consumption by the *pprR*-complemented mutant was achieved approximately 30 h later. This could be due to the markedly higher transcript level of *pprR* (~100-fold; Fig. 4D) in the *pprR*-complemented mutant, since it carries the *pprR* gene on a high expression vector. This further supports PprR to function as a transcriptional repressor, with higher *pprR* expression levels resulting in prolonged repression of 3-phenylpropanoid gene cluster expression.

FIG 5 Relative transcript abundances of 3-phenylpropanoid genes in *A. aromaticum* EbN1^T grown with succinate (A), benzoate (B), or *p*-coumarate (C). Cells were harvested at half-maximum optical density, and transcripts of target genes were normalized against *gyrB*. Each data point is based on 3 biological replicates with 3 technical replicates analyzed for each. Colors: blue, wild type; pink, $\Delta pprR$ mutant; gray-blue, *pprR*-complemented mutant.

In contrast to succinate- and benzoate-grown cells, transcript profiles of *p*-coumarate-grown cells revealed no differences between the three genotypes. Interestingly, transcript levels in all three genotypes were at least 1 order of magnitude higher with *p*-coumarate than those (derepressed) of the $\Delta pprR$ mutant with either succinate or benzoate (Fig. 5). Accordingly, the respective proteins were exclusively detected (Fig. 3B and Fig. S6B and C) in *p*-coumarate cells of all three genotypes.

Benzoate-induced diauxie. The above-described transcript profiles of the three genotypes (Fig. 5) indicated the presence of additional regulatory mechanisms controlling transcription of the 3-phenylpropanoid gene cluster (*ebA5316*–21). To investigate this assumption, cells of *A. aromaticum* EbN1^T (wild type and $\Delta pprR$ mutant) were supplied with a binary mixture of *p*-coumarate (1 mM) and either benzoate (1 mM) or succinate (3 mM). With the mixture of *p*-coumarate and benzoate, both genotypes clearly showed diauxic growth accompanied by preferential utilization of benzoate (Fig. 6A and Fig. S7A). For both genotypes, maximum relative *ebA5316*–21 transcript abundances were observed shortly after benzoate was completely consumed (Fig. 6A and Fig. S7A).

Interestingly, the wild type and $\Delta pprR$ mutant showed monophasic growth and concomitant utilization of substrates when supplied with a mixture of *p*-coumarate and succinate

FIG 6 Anaerobic cultivation of *A. aromaticum* EbN1^T (wild type) for targeted transcript analysis in response to a binary mixture of 1 mM *p*-coumarate and either 1 mM benzoate (A) or 3 mM succinate (B). Growth was monitored by measuring the optical density at 660 nm (OD₆₆₀). Sampling time points for transcript analysis are indicated by gray dashed lines. Fold changes of each gene at each time point were normalized against *gyrB*. Each data point is based on 3 biological replicates with 3 technical replicates analyzed for each.

(Fig. 6B and Fig. S7B). However, both genotypes consumed succinate at a higher rate than *p*-coumarate. The wild type reached maximum relative *ebA5316-21* transcript abundances only after complete consumption of succinate (Fig. 6B). In contrast, this was achieved by the $\Delta pprR$ mutant already when only half of the initially supplied succinate was consumed (Fig. S7B).

Benzoyl-CoA was previously reported as the effector of transcriptional repressor BzdR in *Azoarcus* sp. strain CIB (which needs reclassification as *Aromatoleum* species) (24). Therefore, the formation of CoA-ester intermediates involved in anaerobic degradation of *p*-coumarate and benzoate was analyzed in cells of *A. aromaticum* EbN1^T

FIG 7 Quantification of (aromatic) acyl-CoA metabolites in cells of *A. aromaticum* EbN1^T (wild type and $\Delta pprR$ mutant) adapted to anaerobic growth with *p*-coumarate, benzoate, or succinate. Shades of red represent the number of molecules per cell. For each substrate condition and genotype at least three biological replicates were analyzed. Compound names: 5 to 10, see the legend to Fig. 1; 11, 6-hydroxycyclohex-1-ene-1-carbonyl-CoA; 12, 6-oxocyclohex-1-ene-1-carbonyl-CoA; 13, 3-hydroxypimelyl-CoA.

(wild type and $\Delta pprR$ mutant). Under these substrate conditions, both genotypes contained equal amounts of benzoyl-CoA, $\sim 700 \times 10^3$ molecules cell⁻¹, corresponding to $\sim 1.5 \mu\text{mol g}_{\text{CDW}}^{-1}$, respectively (CDW, cell dry weight) (Fig. 7 and Tables S8 and S9). Notably, the amount of benzoyl-CoA in these cells is ~ 50 -fold higher than that of all other identified (aromatic) acyl-CoA esters (Fig. 7). In contrast, succinate-grown cells of both genotypes harbored significantly smaller amounts of benzoyl-CoA ($\sim 30 \times 10^3$ molecules cell⁻¹ or $0.05 \mu\text{mol g}_{\text{CDW}}^{-1}$). As expected, the *p*-coumarate-specific CoA-ester intermediates could only be detected in cells of the wild type and $\Delta pprR$ mutant upon growth with this phenylpropanoid (Fig. 7).

DISCUSSION

In vivo response threshold toward 3-phenylpropanoids. Recently, Vagts et al. (29) revealed highly sensitive responsiveness of *A. aromaticum* EbN1^T toward alkylphenols. There, it was concluded that such low (lower nanomolar range) response thresholds could have implications for microbial viability at low substrate availability as well as for the mechanisms underlying the persistence of dissolved organic matter (DOM) in a variety of environmental settings. The diverse chemical structures and properties of the tested 3-phenylpropanoids prompted a 2-fold reflection on their determined *in vivo* response thresholds, as discussed below.

(i) Factors controlling attainment of transmembrane effector equilibrium at response threshold concentrations. In contrast to uncharged alkylphenols, 3-phenylpropanoids represent charged molecules (dissociated carboxyl group) at ambient pH. Hence, besides passive diffusion across the cell envelope, there is another means of uptake, in which an ABC transporter in conjunction with the predicted SBP EbA5316 (see above) is assumed to be involved (Fig. 1). In the following paragraphs, we evaluate to what degree active uptake versus passive diffusion of membrane-permeable solutes contributes to attaining transmembrane equilibrium of 3-phenylpropanoids at the response threshold concentrations determined in this study.

As a first step, we estimated the abundance of EbA5316 in *A. aromaticum* EbN1^T actively growing with ample provision of *p*-coumarate. For this purpose, we used the proteomic data of the mutant characterization, generated at $\frac{1}{2}\text{OD}_{\text{max}}$, when approximately 1 mM *p*-coumarate was still available (Fig. 4C). We evaluated these data by referring our measured peptide counts of selected ribosomal proteins against reported copy numbers per cell of the respective homologues in *E. coli* growing in minimal medium with 22 mM glucose and harvested in early logarithmic phase (35) (see Fig. S8 in the supplemental material). We obtained an estimate of 5×10^3 to 10×10^3 copies of EbA5316 per actively growing cell of *A. aromaticum* EbN1^T. Given the limit of accuracy

of our approach (50% variability in multiplier $\bar{M}$; see Fig. S8), our estimate is on the same order of magnitude as the copy numbers (22×10^3 and 45×10^3) determined for the maltose-binding protein (MalE) in logarithmically growing *E. coli* supplemented with 5.8 mM (36) and 11.6 mM (37) maltose, respectively. In *A. aromaticum* EbN1^T, 10,000 copies of EbA5316 in the periplasm (estimated volume, 10^{-16} liters [29]) correspond to a concentration of ~ 0.14 mM.

From dialysis experiments (38, 39) and theoretical reasoning (40), it was recognized early on that SBP copy numbers per cell in the range of 10^3 to 10^4 afford two major functions for the cell. To begin with, they allow for an efficient collection of ligands from the solute flux impinging on the cell. Accordingly, 3×10^3 SBPs already well dispersed in the periplasmic space, only covering 0.1% of the cell surface, would suffice to collect 50% of the flux, thereby enhancing the effective absorption cross section 1,000-fold (40). Furthermore, SBPs at high periplasmic concentrations relative to their dissociation constant (K_d) values act as a retention buffer to maintain the periplasmic solute concentration for a long time (38, 41). Lastly, a highly abundant SBP promotes the transfer of the solute from its SBP-bound state to the permease-mediated uptake and avoids unproductive dissociation of the solute back into the periplasm or extracellular space.

As a second step, we estimated the basal level of EbA5316 in benzoate-adapted cells of *A. aromaticum* EbN1^T, viz. under noninduced conditions. For this purpose, we are considering two observations: (i) fully induced cells of *A. aromaticum* EbN1^T have an ~ 100 -fold increased *ebA5316* transcript level (Table S4), and (ii) according to the study by Lu et al. (35), the ratio of synthesized protein per mRNA molecule typically is at least 100 in *E. coli*. Taken together, we estimate the EbA5316 abundance in noninduced cells of *A. aromaticum* EbN1^T to be $\sim 10^4$ -fold lower than that of *p*-coumarate-adapted cells. Considering that we estimated the latter to carry about 5×10^3 to 10×10^3 copies of EbA5316, this SBP might be present in single-figure copy numbers, if at all, in noninduced cells. Such an estimate is in accord with the lack of proteomic detection of EbA5316 in succinate- or acetate-adapted cells (Fig. 3B and Fig. S6). With benzoate-adapted cells, we have detected EbA5316, but not consistently (absence, Fig. S6; presence, Fig. 3B). This may be due to a certain degree of unspecificity of the involved regulators for aromatic acyl-CoA effectors.

As a third step, we assessed whether an SBP present at low copy numbers in noninduced cells could contribute to attaining effector equilibrium at the low-nanomolar threshold concentrations. In this context, the affinity of a given SBP to its effector molecule is decisive. Therefore, we compiled, from the literature, 89 experimentally determined K_d values of purified SBPs associated with ABC and TRAP transporters specific for different classes of solutes (Fig. 8 and Table S10). For growth-supporting organic substrates, K_d values range from 8 nM to 716 μ M with a median value between 0.5 and 1.0 μ M. At threshold substrate concentrations, $[S]$, of 10 nM, as determined in the present study (Fig. 2), the ratio of unbound SBP ($[P]$) to bound SBP ($[P \cdot S]$) ranges between 50:1 and 100:1 according to the following equation:

$$\frac{K_d}{[S]} = \frac{[P]}{[P \cdot S]} \quad (1)$$

At such ratios, an SBP-mediated uptake is not efficient. Even when considering the lowest K_d value reported for an aromatic carboxylate (95 nM for *p*-hydroxybenzoate [34]), the ratio of free to bound SBP is still 10:1. For EbA5316 to play a role in substrate uptake at response threshold concentrations, its K_d value would have to be distinctly lower than the experimentally determined threshold concentrations (Fig. 8A), i.e., in the sub-nanomolar range, which is currently known only for compounds such as vitamins and trace elements (Fig. 8B). Therefore, we conclude that passive diffusion of 3-phenylpropanoids across the cell envelope apparently drives the attainment of the transmembrane effector equilibrium, as recently described for alkylphenols (29).

FIG 8 Correlation of response threshold with K_d values of known SBPs. (A) Response threshold (red) for 3-phenylpropanoids determined in this study. K_d values (gray) theoretically inferred for the predicted SBP EbA5316 from the *in vivo* response thresholds. Abbreviations: 3PP, 3-phenylpropanoate; 3HP, 3-(4-hydroxyphenyl)propanoate; Cin, cinnamate; *p*-Cou, *p*-coumarate. (B) K_d values of known SBPs as retrieved from published experimental studies (K_d values, further details, and references are compiled in Table S10). Note that the category “others” comprises SBPs for vitamins, cofactors, trace elements, and signaling molecules. Black lines, medians.

(ii) Rationalizing higher threshold observed for *p*-coumarate. Among the four tested 3-phenylpropanoids, 3-phenylpropanoate yields the highest overall transcriptional response (~ 300 -fold for *ebA5317*). Additional structural features [ring hydroxylation in 3-(4-hydroxyphenyl)propanoate versus alkenoyl chain in cinnamate] lead to a lower overall transcriptional response intensity (~ 100 -fold for *ebA5316*), albeit to a response threshold (1 to 10 nM) similar to that for 3-phenylpropanoate (Fig. 2). In contrast, *p*-coumarate, characterized by a *para*-hydroxyl group as well as an alkenoyl chain, delivers a lower overall transcriptional response as well as a higher response threshold (30 to 50 nM).

Even though one may speculate that EbA5316 has a lower affinity to *p*-coumarate, this should be irrelevant for its threshold concentration, since SBP-mediated active uptake does not play a role in attainment of the effector equilibrium, as outlined above. Alternatively, the higher threshold for *p*-coumarate may reflect a lower turnover by CoA-ligase EbA5317, yielding a lower pool size of *p*-coumaroyl-CoA than the CoA-esters formed from the other three structurally different 3-phenylpropanoids. This may be plausible, since EbA5317 represents the first common enzyme during degradation of all four tested 3-phenylpropanoids, and the respective CoA-esters most likely act as the true effectors for the repressor PprR. The latter would be in accord with a previous report on the analogous transcriptional repressor CouR (*R. palustris*), which is not responsive to *p*-coumarate but instead binds *p*-coumaroyl-CoA as its effector (32).

Multilayered transcriptional regulation of 3-phenylpropanoid genes. Characterization of the $\Delta pprR$ mutant underpinned the predicted function of PprR as a transcriptional repressor of *ebA5316-21* expression in a 3-phenylpropanoid-dependent manner. Unexpectedly, however, even in the $\Delta pprR$ mutant, maximum relative transcript abundances of *ebA5316-21* were only observed with *p*-coumarate rather than with benzoate or succinate (Fig. 5). This suggests the involvement of additional transcriptional control circuits, namely, a

3-phenylpropanoid-specific activation and/or an additional benzoate/succinate-mediated repression. A specific transcriptional activator for *ebA5316-21* expression could not be inferred from analyzing the genome of *A. aromaticum* EbN1^T. Furthermore, while some TetR-family members are known to act as repressor as well as activator (42), such a dual function does not appear likely for PprR of *A. aromaticum* EbN1^T, according to our present data and the close relatedness to the proven TetR-type repressors KstR and KstR2, controlling “cholesterol degradation” genes in mycobacteria (43). One can speculate about catabolite repression, since anaerobic utilization of C₄-dicarboxylates by *A. aromaticum* EbN1^T is repressed in the presence of benzoate (44). In accordance with this, catabolite repression was previously reported to play a role in the catabolism of aromatic compounds in *Thauera aromatica* K172^T (45), *Azoarcus* sp. strain CIB (46), *Acinetobacter baylyi* (47), and *Escherichia coli* (48, 49).

(i) Benzoate-induced diauxie. While the two applied binary substrate mixtures (*p*-coumarate and benzoate versus *p*-coumarate and succinate) yielded converse growth courses (diauxic versus monophasic) with the wild type under both substrate conditions, maximal *ebA5316-21* transcript abundances were observed upon complete consumption of benzoate and succinate (Fig. 6). In the case of the Δ *pprR* mutant, the growth courses were more pronounced and maximal transcript abundances were reached when benzoate was completely and succinate half-completely consumed (Fig. S7). These findings suggest the presence of a regulatory mechanism hindering full transcription of 3-phenylpropanoid genes in the presence of both benzoate and succinate. Furthermore, this mechanism seems to be superior to PprR, as its effect was observed not only with the wild type but also with the Δ *pprR* mutant. However, benzoate appears to more strongly repress *ebA5316-21* transcription than succinate, as it not only delayed full *ebA5316-21* transcription but also completely repressed *p*-coumarate utilization until benzoate depletion.

(ii) Role of benzoyl-CoA in diauxie? Anaerobic degradation of most monoaromatic compounds feeds into the central benzoyl-CoA pathway. Benzoyl-CoA represents not only the entry point into this central degradation pathway but also the effector for BzdR (*K_d* of ~150 μ M), a transcriptional repressor controlling gene expression of the anaerobic benzoyl-CoA pathway in *Azoarcus* sp. strain CIB (50, 51). Thus, intracellular benzoyl-CoA levels might function as a control switch to impose preferential utilization of benzoate in the presence of other aromatic or aliphatic growth substrates in *A. aromaticum* EbN1^T (44) and *Magnetospirillum* sp. strain pMbN1 (52). However, the observed similar intracellular pool sizes of benzoyl-CoA in wild-type and Δ *pprR* mutant cells of *A. aromaticum* EbN1^T grown with either benzoate or *p*-coumarate (Fig. 7) argues against this aromatic acyl-CoA conveying benzoate-induced diauxie. Furthermore, these congruent benzoyl-CoA levels suggest a previously reported preference of benzoate over succinate in *A. aromaticum* EbN1^T (44) likewise to not be mediated by benzoyl-CoA, since *p*-coumarate and succinate were concomitantly utilized (Fig. 6 and Fig. S7). The ~25-fold lower but nevertheless detectable level of intracellular benzoyl-CoA in succinate-grown cells most likely reflects its additional role in anabolic processes.

In conclusion, our findings support the assumption of a higher-order transcriptional regulator being responsible for tuning the studied degradation pathways of *A. aromaticum* EbN1^T in a benzoate- and succinate-dependent manner comparable to the ones described for *E. coli* (53) and *Pseudomonas* (54, 55).

MATERIALS AND METHODS

Bacterial strains and cultivation conditions. *A. aromaticum* EbN1^T was cultivated under nitrate-reducing conditions in a defined, ascorbate-reduced, and bicarbonate-buffered mineral medium at 28°C as previously described (22). Organic substrates [acetate, succinate, benzoate, *p*-coumarate, 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, and cinnamate] were added from aqueous stock solutions sterilized by filtration. Each experimental line (response threshold determination, mutant characterization, and differential proteomics) was preceded by the same cultivation steps with benzoate as the substrate for anaerobic growth as described before (28, 29). Essentially, glycerol stocks of *A. aromaticum* EbN1^T were used to inoculate a dilution series (10⁻¹ to 10⁻⁶) as the starting point for two successive

TABLE 1 Oligonucleotide primers used for mutant construction

Primer ^a	Target gene	Nucleotide sequence (5' → 3')	Product length (bp)
Gene-specific primer pair			
pprR_240_F	pprR	GCTGAGTGACAGCCAACG	235
pprR_474_R		CGACGCGAAGTGGTGATA	
Generation of ΔpprR deletion mutation			
benK_PstI_F	benK	GCCGCTGCAGGCTGGCA	1,250
benK_BamHI_R	benK	AAGGATCCTAGCGGGCGCACCCCG	
ebA5316_BamHI_F	ebA5316	GGTGCGCCCGCTAGGATCCCATCCGGAATCCTGCCGC	1,500
ebA5316_BamHI_R	ebA5316	GCTCGGTACCCGGGGATCCGTGCCTCGATGTTACGACC	
Identification of ΔpprR genotype			
ΔpprR_F	pprR	GCTGAGTGACAGCCAACG	292
ΔpprR_R		CGACGCGAAGTGGTGATA	
Generation of in trans-complementation of pprR			
pprR_EcoRI_F	pprR	CGATCGCGGAATTCGCGGTGTC	1,434
pprR_ApaI_R		ACGCACCGGGCCCGG	

^aF, forward primer; R, reverse primer.

precultures (80-ml culture volume), followed by a main culture (400-ml culture volume), which served as the inoculum for each actual experiment. For each experimental line (see below), cultivation was performed in 500-ml flat-bottomed glass bottles (400-ml culture volume) sealed with butyl rubber stoppers. All chemicals were of analytical grade. Growth was monitored by measuring the optical density of the cultures at 660 nm (OD_{660}).

Generation of in-frame $\Delta pprR$ deletion mutation. DNA was isolated according to standard protocols (56). Generation of the in-frame $\Delta pprR$ deletion mutation was carried out as previously described for the *etpR* and *pcrSR* genes of *A. aromaticum* EbN1^T (29, 30) and using the primers listed in Table 1. The $\Delta pprR$ mutant retained the start and stop codons of *pprR* flanking a BamHI restriction site. The final knockout vector, pk19 $\Omega benKebA5316$, for in-frame deletion of *pprR* (*ebA5314*) was constructed in an *E. coli* S17-1 background by following a two-step strategy. First, a 1,250-bp fragment (*benK*) representing the 3'-flanking region of *pprR* including the stop codon of *benK*, and a BamHI restriction site was amplified from genomic DNA of *A. aromaticum* EbN1^T using a Phusion high-fidelity DNA polymerase (ThermoFischer Scientific, Waltham, MA, USA) and cloned into the vector using its PstI and BamHI restriction sites, giving rise to pk19 $\Omega benK$. Second, a 1,500-bp fragment (*ebA5316*) was amplified as described above and cloned into pk19 $\Omega benK$ using the In-Fusion HD cloning plus kit (TaKaRa Bio Inc., Kusatsu, Japan) according to the manufacturer's instructions. This fragment represented the 5'-flanking region of *pprR*, including the start codon of *ebA5316* and a BamHI restriction site, yielding pk19 $\Omega benKebA5316$. Conjugational plasmid transfer to *A. aromaticum* EbN1^T was conducted as described previously (56). PCR-based screening of single-crossover clones (kan^R) involved the $\Delta pprR$ primer pair, yielding 316-bp and 1,207-bp amplicons, respectively (see Fig. S5 in the supplemental material). The second crossover event was achieved by transferring the single-crossover mutant into kanamycin-free liquid medium containing a mixture of benzoate (4 mM), acetate (5 mM), and pyruvate (5 mM). Plating on solid medium containing the same tripartite substrate mixture and 5% (wt/vol) sucrose was used for selection of the $\Delta pprR$ genotype. For identification, colonies were screened as described above, yielding only one amplicon of 316 bp (Fig. S5).

Complementation of *pprR* in *trans* into $\Delta pprR$ deletion mutant. In *trans*-complementation was performed as described in two previous studies (29, 30) and using the primers listed in Table 1. Essentially, a 1,434-bp fragment containing the *pprR* gene together with its ribosomal binding site was amplified from genomic DNA of *A. aromaticum* EbN1^T as described above. The fragment was cloned into the broad-host-range vector pBBR1MCS-2 using its ApaI and EcoRI restriction sites. The final construct was verified by sequencing. The vector was transferred from the *E. coli* S17-1 background to the $\Delta pprR$ mutant of *A. aromaticum* EbN1^T via conjugation, yielding the *pprR*-complemented mutant (genotype of $\Delta pprR$ pBBR1MCS-2 $\Omega pprR$). The conjugational transfer via agar plate mating and the verification of positive clones was performed as previously described (56).

Sequence validation by Sanger sequencing. For sequence validation of the obtained $\Delta pprR$ mutant, a 2,393-bp region of genomic DNA spanning across the entire up- and downstream regions, including the deletion site, was analyzed via Sanger sequencing (57), as previously described (29). Additionally, a 1,892-bp region of the complementation vector (pBBR1MCS-2 $\Omega pprR$) spanning the entire insert was sequenced. Used primer pairs are compiled in Table S11.

Growth experiments. (i) Threshold determination. The responsiveness of *A. aromaticum* EbN1^T to various concentrations of 3-phenylpropanoate, 3-(4-hydroxyphenyl)propanoate, cinnamate, and *p*-coumarate, respectively, was assessed via an experimental setup essentially as described in previous studies (28, 29). In short, benzoate-adapted cells of *A. aromaticum* EbN1^T were exposed to a single pulse of each tested 3-phenylpropanoid, yielding extracellular concentrations ranging from 100 μ M down to 0.1 nM. Subsequently, samples for monitoring growth and time-resolved transcript profiling were retrieved from the cultures as described previously (28, 29): 5 min prior as well as 5, 15, 30, 60, and 120 min after

addition of either 3-phenylpropanoid (effector). The respective detailed growth curves are provided in Fig. S1 to S4.

(ii) Mutant characterization. To assess the effect of the $\Delta pprR$ deletion mutation (see “Generation of in-frame $\Delta pprR$ deletion mutation,” above), the wild type, the $\Delta pprR$ mutant, and the $pprR$ -complemented mutant of *A. aromaticum* EbN1^T were grown with either succinate (8 mM), benzoate (4 mM), or *p*-coumarate (2 mM). Since cultures were inoculated with benzoate-adapted cells (described above), in the case of succinate and *p*-coumarate, substrate adaptation had to be conducted over five passages at first. Cultures of the $pprR$ -complemented mutant contained kanamycin (50 $\mu\text{g ml}^{-1}$). For each substrate condition, three replicate cultures were conducted. These cultures were sampled in short intervals to monitor growth (OD_{660}) and substrate consumption. Additional cultures served to provide samples (harvested at $\frac{1}{2}\text{OD}_{\text{max}}$) for targeted transcript and proteomic analyses (see below).

(iii) Binary substrate mixture experiments. To test a possible involvement of catabolite repression-like processes in the regulation of *ebA5316-21* transcription, benzoate-adapted cells of *A. aromaticum* EbN1^T (wild type and $\Delta pprR$ mutant) were grown with a binary mixture of *p*-coumarate (1 mM) and either benzoate (1 mM) or succinate (3 mM).

(iv) (Aromatic) acyl-CoA ester analyses. For the quantification of intracellular (aromatic) acyl-CoA ester pool sizes, *A. aromaticum* EbN1^T (wild type and $\Delta pprR$ mutant) was adapted to anaerobic growth with benzoate, succinate, or *p*-coumarate. Corresponding cells were anaerobically grown with the respective substrate under nitrate-reducing conditions as described above. Substrate concentrations were 1 mM each for benzoate and *p*-coumarate and 3 mM for succinate. For each substrate condition, cultures were harvested at $\frac{1}{2}\text{OD}_{\text{max}}$ for analyses of (aromatic) acyl-CoA esters.

(v) Differential proteomics. To investigate transport systems possibly involved in the uptake of 3-phenylpropanoids, *A. aromaticum* EbN1^T was grown with acetate (8 mM), benzoate (4 mM), or 3-(4-hydroxyphenyl)propanoate (2 mM). Again, cultures were adapted to each substrate over five passages and harvested at $\frac{1}{2}\text{OD}_{\text{max}}$.

Cell harvesting. (i) Transcript analyses. Samples were retrieved and processed as described previously (28, 29). Essentially, 5 ml of culture broth were withdrawn and immediately added to RNeasy Protect bacterial reagent (Qiagen, Hilden, Germany), mixed thoroughly, incubated for 5 min at ambient temperature, and centrifuged. Pellets were washed with 0.5 ml RNeasy Protect bacterial reagent (Qiagen), shock frozen in liquid N_2 , and stored at -80°C until further analysis.

(ii) (Aromatic) acyl-CoA ester analyses. Substrate-adapted cultures were transferred into N_2 -flushed centrifuge beakers while maintaining a constant flow of N_2 . After centrifugation ($14,334 \times g$, 15 min, 4°C), cell pellets were each resuspended in 1 ml methanol (ULC/MS; Biosolve, Valkenswaard, The Netherlands) while flushing with N_2 and transferred into sterile screw-cap 2-ml polypropylene tubes (Sarstedt, Germany) containing 0.6 g of glass beads (0.1-mm diameter) and 0.4 g zirconia beads (0.7-mm diameter) (Carl Roth, Karlsruhe, Germany). The samples were shock-frozen in liquid N_2 and stored at -80°C until further analysis.

(iii) Proteomic analysis. Samples were retrieved and processed as previously described (58). In short, the entire 400-ml cultures were centrifuged ($14,334 \times g$, 30 min, 4°C), and the pellets were washed in 250 ml washing buffer (100 mM Tris-HCl, 5 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, pH 7.5) and resuspended in 0.8 ml of the same washing buffer. Following centrifugation ($20,000 \times g$, 10 min, 4°C), pellets were shock frozen in liquid N_2 and stored at -80°C until further analysis.

Quantification of growth substrates by (micro)HPLC. Quantitative depletion profiling of benzoate and *p*-coumarate was achieved using a micro-high-performance liquid chromatography (microHPLC) system (UltiMate 3000; ThermoFisher Scientific, Germering, Bavaria, Germany) as previously described (29). Essentially, compounds were separated via reverse-phase chromatography (Hypersil Gold column; ThermoFisher Scientific), applying an acetonitrile gradient and detected by means of a diode array detector. Benzoate was detected at 229 nm, with a retention time of 9.32 min and a dynamic range from 5 nM to 50 μM . *p*-Coumarate was detected at 309 nm, with a retention time of 8.84 min and a dynamic range from 25 nM to 50 μM .

Quantification of succinate was achieved with an HPLC system (UltiMate 3000; ThermoFischer) equipped with an ion exchange column (Eurokat H; 300 by 8 mm, 10- μm inner diameter; Knauer, Berlin, Germany) and an RI detector (RI-101; Shodex, Munich, Germany). The system was operated at 75°C with H_2SO_4 (5 mM) as the eluent provided, at a flow rate of 1.2 ml min^{-1} . Succinate was detected at a retention time of 6.51 min with a dynamic range from 25 μM to 10 mM.

Preparation of total RNA. Preparation of total RNA was performed according to standard methods (50, 59) and as previously described (29). (i) Essentially, stored cell pellets were treated twice with hot acidic phenol (Roti Aqua-Phenol; CarlRoth, Karlsruhe, Germany), and the resultant aqueous phase was transferred into a 2-ml SPRIIME phase-lock gel tube (Quantabio, Beverly, MA, USA). After addition of one volume of phenol-chloroform-isoamyl alcohol (25:24:1), the tube was gently inverted for 5 min and centrifuged. (ii) Nucleic acids were precipitated and washed with ice-cold ethanol. (iii) The pellet was dried and resuspended in RNase-free water. Digestion with DNase I (RNase-free; Qiagen) was verified via PCR. The RNA quality of all 792 samples was assessed with an Experion automated electrophoresis station (Bio-Rad, Hercules, CA, USA). RNA concentration was determined using a TrayCell (Hellma Analytics, Müllheim, Germany) operated in a spectrophotometer (UV-1800; Shimadzu, Duisburg, Germany). RNA samples were stored at -80°C until further analysis. All chemicals used for RNA preparation were of molecular biology grade.

Transcript profiling by qRT-PCR. For the selected target genes, specific primers (Table 2) were designed using the Primer3 software package (version 0.4.0; www.primer3.org). Synthesis of cDNA and real-time PCR was performed with 150 ng of total RNA using the Brilliant III ultrafast SYBR green

TABLE 2 Oligonucleotide primers used for targeted transcript profiling for 3-phenylpropanoid catabolic genes

Primer ^a	Target gene	Nucleotide sequence (5'→3')	Product length (bp)	PCR efficiency ^b
ebA5316_F	ebA5316	GACGCCTACGGTGAAGACTG	266	1.89
ebA5316_R		ACCTTCAGGAAAGCCTGGTT		
ebA5317_F	ebA5317	CCGAAAGCGGTCATCAATAC	266	1.97
ebA5317_R		GGCACGCTGAAATAGATCGT		
ebA5319_F	ebA5319	GAGCTTCGATGCGTATTTC	276	2.06
ebA5319_R		GTGTCCTTCGTCGCTTCC		
ebA5321_F	ebA5321	GCCAGTATGGGAGATGTGGT	246	1.89
ebA5321_R		CCGAAGAGGCGAAAACCTC		

^aF, forward primer; R, reverse primer.^bMean value from all performed qRT-PCR experiments.

quantitative reverse transcription-PCR (qRT-PCR) master mix (Agilent, Santa Clara, CA, USA) and analyzed in a CFX96 real-time system (Bio-Rad). PCR settings comprised initiation for 3 min at 95°C, 40 cycles of 10 s of denaturation at 95°C, 30 s of annealing at 60°C, and 30 s of extension at 60°C, followed by real-time detection for 5 s. For validation of specific product formation, melting-curve analysis was performed with intervals of 0.5°C, ranging from 60°C to 90°C. For time-resolved transcript profiling, all samples retrieved after effector addition represented test states, while a sample retrieved 5 min prior to effector addition served as the reference. For assessing the effect of the $\Delta pprR$ deletion mutation, relative transcript levels of each target gene were normalized against a housekeeping gene (*gyrB*; encoding DNA gyrase subunit B). Three biological replicates per sample were retrieved, for each of which 3 technical replicate measurements were performed. Relative transcript abundances were calculated, as previously described (28), according to the following equation:

$$\text{ratio} = E^{\Delta C_T(\text{reference} - \text{test})} \quad (2)$$

Primer-specific PCR efficiencies (E) were determined as described previously (60).

Identification and quantification of (aromatic) acyl-CoAs. Cell breakage was achieved by vigorous shaking at 10 m s⁻¹ in a bead beater (MP Biomedicals, Schwerte, Germany) for 40 s, followed by cooling on ice for 2 min. This step was repeated twice. Then, the solution was centrifuged at 14,000 × g and 4°C for 10 min, and the supernatant was transferred to a 4-ml opaque vial. The remaining pellet was resuspended in 0.5 ml methanol, homogenized, and centrifuged, and the supernatant was transferred to the same 4-ml vial. This step was repeated once with 0.5 ml methanol. The solution was dried down under a nitrogen stream, and the residue was dissolved in 200 μ l ultrapure water. The extract was transferred to a centrifuge filter (0.22- μ m cellulose acetate filter, 2.0-ml polypropylene tube; Corning, Glendale, AZ, USA) and centrifuged under the same conditions as those described above.

The filtered solutions were analyzed by liquid chromatography on a Vanquish UHPLC system (ThermoFisher Scientific, Waltham, MA, USA) coupled with an Orbitrap Fusion mass spectrometer (ThermoFisher Scientific). The UHPLC was equipped with a Gemini C₁₈ column (150 by 2 mm, 3- μ m pore size; Phenomenex, Torrance, CA, USA) and operated at 35°C with an eluent flow rate of 400 μ l min⁻¹. The amount of injection was 5 μ l. The mobile phase was a gradient composed of 10 mM ammonium formate at pH 8.1 (A) and acetonitrile (ULC-MS; Biosolve) (B). The gradient steps were the following: 0 to 2 min, 100% A; 2 to 20 min, 100% to 80% A; 20 to 23 min, 80% to 0% A; 23 to 27 min, 0% A; 27 to 29 min, 0% to 100% A; and 29 to 34 min, 100% A. The settings of the H-ESI-II source of the mass spectrometer were 4,000 V spray voltage, 350°C vaporizer temperature, 320°C ion transfer tube temperature, 40 arbitrary units sheath gas, and 10 arbitrary units auxiliary gas. The Orbitrap instrument was operated in full scan mode with a resolution of 60,000, a mass range of 750 to 950 Da, a 60% RF lens, an AGC target of 4.0e⁵, and a maximum injection time of 50 ms.

p-Coumaroyl-CoA, 4-hydroxybenzoyl-CoA, and benzoyl-CoA were identified by comparison with reference standards, which were synthesized according to the method of Kawaguchi et al. (61). The other coenzyme A esters were tentatively identified based on relative retention times and exact masses of the molecular ions. The individual coenzyme A thioesters were quantified by external calibration using synthetic benzoyl-CoA.

Proteomic analysis. (i) Whole-cell shotgun analysis. Analysis of the soluble protein fraction was performed as previously described (29). In short, cells were disrupted by means of bead beating, and the soluble protein fraction was obtained via ultracentrifugation. After quantification using the Bradford assay (62), proteins were subjected to tryptic digestion. The peptide mixtures were decomplexed using a nanoLC system run in a trap column mode (UltiMate3000 nanoRSLC; ThermoFisher Scientific) applying an acetonitrile gradient. Continuous analysis proceeded via an online-coupled ion-trap mass spectrometer (amaZon speed ETD; Bruker Daltonik GmbH, Bremen, Germany) equipped with a captive spray ion source (Bruker Daltonik GmbH). Two different lengths of linear gradients were applied: 180 min, samples for mutant characterization (Fig. S6); 280 min, samples for comprehensive pathway analysis (Fig. 3B). Proteins were identified using the ProteinScape platform (version 3.1; Bruker Daltonik GmbH) on an in-house Mascot server (version 2.3; Matrix Science Ltd., London, UK) based on the complete genome sequence of *A. aromaticum* EbN1^T (27), employing a target-decoy strategy (63).

(ii) Membrane protein-enriched fraction. Analysis of the membrane protein-enriched fraction was conducted essentially as described in a previous study (58). Briefly, cell disruption was achieved using a French press (French pressure cell press SLM Aminco; BioSurplus Inc., San Diego, CA, USA), and the resultant extract was treated with Na_2CO_3 (100 mM) prior to ultracentrifugation. Proteins were solubilized from the membrane fraction using SDS (1% [wt/vol]), followed by ultracentrifugation. The supernatant was shock-frozen in liquid N_2 and stored at -80°C until further analysis. Proteins were separated by SDS-PAGE (10 μg load). Then, sample lanes were cut into 8 slices, which were subjected to in-gel digestion. Subsequently, nanoLC electrospray ionization-tandem mass spectrometry analyses were applied using a 130-min linear gradient. Protein identification of the membrane protein-enriched fraction was performed as described above.

(iii) Estimation of EbA5316 (SBP) copy numbers per cell. As a basis for the estimation of EbA5316 copy numbers, we selected six ribosomal proteins (RplADFIJ and RpmC) of the 50S subunit of *A. aromaticum* EbN1^T (see Fig. S8 in the supplemental material), assuming comparable copy numbers of ribosomal proteins in proteobacterial *A. aromaticum* EbN1^T and *E. coli* during active growth. Accordingly, experimentally determined peptide counts of the six selected ribosomal proteins of *A. aromaticum* EbN1^T were set in relation to the quantified copy numbers of the respective *E. coli* homologues previously reported by Lu et al. (35). This relation was calculated as

$$M_i \cdot \left(\frac{c_i}{f_i} \right)_{\text{EbN1}} = (N_i)_{\text{E. coli}} \quad (3)$$

where $(N_i)_{\text{E. coli}}$ is the copy number (APEX) of protein *i* (running over RplADFIJ and RpmC) per cell determined for *E. coli* (35), c_i denotes the peptide counts of the protein *i*, and f_i represents the number of detected tryptic peptides for that protein. We assume that the detection probability of a tryptically digested protein *i* is proportional to the number, N_i , of tryptic peptides (≥ 5 amino acids). This procedure (equation 3) yields six protein-specific multipliers, M_i , from which we take the average value, $\bar{M}$, for converting measured peptide counts into estimated protein copy numbers of EbA5316 in *A. aromaticum* EbN1^T:

$$\bar{M} \cdot \left(\frac{c_i}{f_i} \right)_{\text{EbN1}} = (N_i)_{\text{EbN1}} \quad (4)$$

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 3 MB.

ACKNOWLEDGMENTS

We are grateful to Christina Hinrichs and Kerstin Zdrodowski (both from Oldenburg) for technical assistance. We are also grateful for Sebastian Swirski and John Neidhardt (both from Oldenburg) for performing nucleotide sequencing.

J. Vagts, L. Wöhlbrand, and R. Rabus conceived this study; J. Vagts and J. Gutsch conducted the molecular genetic experiments; J. Vagts, K. Kalvelage, A. Weiten, R. Buschen, and S. Scheve performed the cultivation experiments; S. Scheve conducted the (micro-)HPLC analyses; J. Vagts, K. Kalvelage, A. Weiten, and S. Scheve did the RNA work; M. Winkelhofer performed calculations; S. Diener and H. Wilkes performed the (aromatic) acyl-CoA ester analyses; J. Vagts, M. Winkelhofer, and R. Rabus wrote the manuscript.

This study was supported by the Deutsche Forschungsgemeinschaft (DFG) within the framework of the research training group Molecular Basis of Sensory Biology (GRK 1885).

REFERENCES

- Schaefer AL, Greenberg EP, Oliver CM, Oda Y, Huang JJ, Bittan-Banin G, Peres CM, Schmidt S, Juhaszova K, Sufrin JR, Harwood CS. 2008. A new class of homoserine lactone quorum-sensing signals. *Nature* 454:595–599. <https://doi.org/10.1038/nature07088>.
- Kumar R, Singh S, Singh OV. 2008. Bioconversion of lignocellulosic biomass: biochemical and molecular perspectives. *J Ind Microbiol Biotechnol* 35:377–391. <https://doi.org/10.1007/s10295-008-0327-8>.
- Rubin EM. 2008. Genomics of cellulosic biofuels. *Nature* 454:841–845. <https://doi.org/10.1038/nature07190>.
- Bandounas L, Wierckx NJP, de Winder JH, Ruijsenaars HJ. 2011. Isolation and characterization of novel bacterial strains exhibiting ligninolytic potential. *BMC Biotechnol* 11:94. <https://doi.org/10.1186/1472-6750-11-94>.
- Peinemann JC, Pleissner D. 2020. Continuous pretreatment, hydrolysis, and fermentation of organic residues for the production of biochemicals. *Bioreour Technol* 295:122256. <https://doi.org/10.1016/j.biortech.2019.122256>.
- De los Santos Ramos W, Poznyak T, Chairez I, Córdova R. 2009. Remediation of lignin and its derivatives from pulp and paper industry wastewater by the combination of chemical precipitation and ozonation. *J Hazard Mater* 169:428–434. <https://doi.org/10.1016/j.jhazmat.2009.03.152>.
- Ahmad M, Taylor CR, Pink D, Burton K, Eastwood D, Bending GD, Bugg TDH. 2010. Development of novel assays for lignin degradation: comparative analysis of bacterial and fungal lignin degraders. *Mol Biosyst* 6:815–821. <https://doi.org/10.1039/b908966g>.
- Arantes V, Milagres AMF. 2007. The synergistic action of ligninolytic enzymes (MnP and Laccase) and Fe^{3+} -reducing activity from white-rot fungi for degradation of azure B. *Enzyme Microb Technol* 42:17–22. <https://doi.org/10.1016/j.enzmictec.2007.07.017>.
- Shary S, Kapich AN, Panisko EA, Magnuson JK, Cullen D, Hammel KE. 2008. Differential expression in *Phanerochaete chrysosporium* of membrane-

- associated proteins relevant to lignin degradation. *Appl Environ Microbiol* 74:7252–7257. <https://doi.org/10.1128/AEM.01997-08>.
10. Masai E, Katayama Y, Fukuda M. 2007. Genetic and biochemical investigations on bacterial catabolic pathways for lignin-derived aromatic compounds. *Biosci Biotechnol Biochem* 71:1–15. <https://doi.org/10.1271/bbb.60437>.
  11. Bugg TDH, Ahmad M, Hardiman EM, Singh R. 2011. The emerging role for bacteria in lignin degradation and bio-product formation. *Curr Opin Biotechnol* 22:394–400. <https://doi.org/10.1016/j.copbio.2010.10.009>.
  12. Brown ME, Walker MC, Nakashige TG, Iavarone AT, Chang MCY. 2011. Discovery and characterization of heme enzymes from unsequenced bacteria: application to microbial lignin degradation. *J Am Chem Soc* 133:18006–18009. <https://doi.org/10.1021/ja203972q>.
  13. Massalha H, Korenblum E, Tholl D, Aharoni A. 2017. Small molecules below-ground: the role of specialized metabolites in the rhizosphere. *Plant J* 90:788–807. <https://doi.org/10.1111/tjp.13543>.
  14. Pétriacq P, Williams A, Cotton A, McFarlane AE, Rolfe SA, Ton J. 2017. Metabolite profiling of non-sterile rhizosphere soil. *Plant J* 92:147–162. <https://doi.org/10.1111/tjp.13639>.
  15. Moussaieff A, Rogachev I, Brodsky L, Malitsky S, Toal TW, Belcher H, Yativ M, Brady SM, Benfey PN, Aharoni A. 2013. High-resolution metabolic mapping of cell types in plant roots. *Proc Natl Acad Sci U S A* 110: E1232–E1241. <https://doi.org/10.1073/pnas.1302019110>.
  16. Schoefer L, Mohan R, Schwierz A, Braune A, Blaut M. 2003. Anaerobic degradation of flavonoids by *Clostridium orbiscindens*. *Appl Environ Microbiol* 69:5849–5854. <https://doi.org/10.1128/aem.69.10.5849-5854.2003>.
  17. Atkinson S, Williams P. 2009. Quorum sensing and social networking in the microbial world. *J R Soc Interface* 6:959–978. <https://doi.org/10.1098/rsif.2009.0203>.
  18. Harwood CS, Gibson J. 1988. Anaerobic and aerobic metabolism of diverse aromatic compounds by the photosynthetic bacterium *Rhodospseudomonas palustris*. *Appl Environ Microbiol* 54:712–717. <https://doi.org/10.1128/AEM.54.3.712-717.1988>.
  19. Davis JR, Sello JK. 2010. Regulation of genes in *Streptomyces* bacteria required for catabolism of lignin-derived aromatic compounds. *Appl Microbiol Biotechnol* 86:921–929. <https://doi.org/10.1007/s00253-009-2358-0>.
  20. Trautwein K, Wilkes H, Rabus R. 2012. Proteogenomic evidence for β -oxidation of plant-derived 3-phenylpropanoids in “*Aromatoleum aromaticum*” EbN1. *Proteomics* 12:1402–1413. <https://doi.org/10.1002/pmic.201100279>.
  21. Parke D, Ornston LN. 2003. Hydroxycinnamate (*hca*) catabolic genes from *Acinetobacter* sp. strain ADP1 are repressed by HcaR and are induced by hydroxycinnamoyl-coenzyme A thioesters. *Appl Environ Microbiol* 69:5398–5409. <https://doi.org/10.1128/aem.69.9.5398-5409.2003>.
  22. Rabus R, Widdel F. 1995. Anaerobic degradation of ethylbenzene and other aromatic hydrocarbons by new denitrifying bacteria. *Arch Microbiol* 163:96–103. <https://doi.org/10.1007/BF00381782>.
  23. Rabus R, Trautwein K, Wöhlbrand L. 2014. Towards habitat-oriented systems biology of “*Aromatoleum aromaticum*” EbN1. Chemical sensing, catabolic network modulation and growth control in anaerobic aromatic compound degradation. *Appl Microbiol Biotechnol* 98:3371–3388. <https://doi.org/10.1007/s00253-013-5466-9>.
  24. Rabus R, Wöhlbrand L, Thies D, Meyer M, Reinhold-Hurek B, Kämpfer P. 2019. *Aromatoleum* gen. nov., a novel genus accommodating the phylogenetic lineage including *Azoarcus evansii* and related species, and proposal of *Aromatoleum aromatoleum* sp. nov., *Aromatoleum petrolei* sp. nov., *Aromatoleum bremense* sp. nov., *Aromatoleum toluolicum* sp. nov. and *Aromatoleum diolicum* sp. nov. *Int J Syst Evol Microbiol* 69:982–997. <https://doi.org/10.1099/ijsem.0.003244>.
  25. Wöhlbrand L, Kallerhoff B, Lange D, Hufnagel P, Thiermann J, Reinhardt R, Rabus R. 2007. Functional proteomic view of metabolic regulation in “*Aromatoleum aromaticum*” strain EbN1. *Proteomics* 7:2222–2239. <https://doi.org/10.1002/pmic.200600987>.
  26. Wöhlbrand L, Wilkes H, Halder T, Rabus R. 2008. Anaerobic degradation of *p*-ethylphenol by “*Aromatoleum aromaticum*” strain EbN1: pathway, regulation, and involved proteins. *J Bacteriol* 190:5699–5709. <https://doi.org/10.1128/JB.00409-08>.
  27. Rabus R, Kube M, Heider J, Beck A, Heitmann K, Widdel F, Reinhardt R. 2005. The genome sequence of an anaerobic aromatic-degrading denitrifying bacterium, strain EbN1. *Arch Microbiol* 183:27–36. <https://doi.org/10.1007/s00203-004-0742-9>.
  28. Vagts J, Scheve S, Kant M, Wöhlbrand L, Rabus R. 2018. Towards the response threshold for *p*-hydroxyacetophenone in the denitrifying bacterium “*Aromatoleum aromaticum*” EbN1. *Appl Environ Microbiol* 84: e01018–18. <https://doi.org/10.1128/AEM.01018-18>.
  29. Vagts J, Weiten A, Scheve S, Kalvelage K, Swirski S, Wöhlbrand L, Neidhardt J, Winkhofer M, Rabus R. 2020. Nanomolar responsiveness of an anaerobic degradation specialist to alkylphenol pollutants. *J Bacteriol* 202:e00595–19. <https://doi.org/10.1128/JB.00595-19>.
  30. Büsing I, Kant M, Dörries M, Wöhlbrand L, Rabus R. 2015. The predicted σ^{54} -dependent regulator EtpR is essential for expression of genes for anaerobic *p*-ethylphenol and *p*-hydroxyacetophenone degradation in “*Aromatoleum aromaticum*” EbN1. *BMC Microbiol* 15:251. <https://doi.org/10.1186/s12866-015-0571-9>.
  31. Eaton RW. 1997. *p*-Cymene catabolic pathway in *Pseudomonas putida* F1: cloning and characterization of DNA encoding conversion of *p*-cymene to *p*-cumate. *J Bacteriol* 179:3171–3180. <https://doi.org/10.1128/jb.179.10.3171-3180.1997>.
  32. Hirakawa H, Schaefer AL, Greenberg EP, Harwood CS. 2012. Anaerobic *p*-coumarate degradation by *Rhodospseudomonas palustris* and identification of CouR, a MarR repressor protein that binds *p*-coumaroyl-coenzyme A. *J Bacteriol* 194:1960–1967. <https://doi.org/10.1128/JB.06817-11>.
  33. Eglund PG, Harwood CS. 1999. BadR, a new MarR family member, regulates anaerobic benzoate degradation by *Rhodospseudomonas palustris* in concert with AadR, an Fnr family member. *J Bacteriol* 181:2102–2109. <https://doi.org/10.1128/JB.181.7.2102-2109.1999>.
  34. Pietri R, Zerbs S, Corgliano DM, Allaire M, Collart FR, Miller LM. 2012. Biophysical and structural characterization of a sequence-diverse set of solute-binding proteins for aromatic compounds. *J Biol Chem* 287:23748–23756. <https://doi.org/10.1074/jbc.M112.352385>.
  35. Lu P, Vogel C, Wang R, Yao X, Marcotte EM. 2007. Absolute protein expression profiling estimates the relative contributions of transcriptional and translational regulation. *Nat Biotechnol* 25:117–124. <https://doi.org/10.1038/nbt1270>.
  36. Dietzel I, Kolb V, Boos W. 1978. Pole cap formation in *Escherichia coli* following induction of the maltose-binding protein. *Arch Microbiol* 118:207–218. <https://doi.org/10.1007/BF00415731>.
  37. Koman A, Harayama S, Hazelbauer GL. 1979. Relation of chemotactic response to the amount of receptor: evidence for different efficiencies of signal transduction. *J Bacteriol* 138:739–747. <https://doi.org/10.1128/JB.138.3.739-747.1979>.
  38. Silhavy TJ, Szmecman S, Boos W, Schwartz M. 1975. On the significance of the retention of ligand by protein. *Proc Natl Acad Sci U S A* 72:2120–2124. <https://doi.org/10.1073/pnas.72.6.2120>.
  39. Willis RC, Furlong CE. 1974. Purification and properties of a ribose-binding protein from *Escherichia coli*. *J Biol Chem* 249:6926–6929. [https://doi.org/10.1016/S0021-9258\(19\)42146-7](https://doi.org/10.1016/S0021-9258(19)42146-7).
  40. Berg HC, Purcell EM. 1977. Physics of chemoreception. *Biophys J* 20:193–219. [https://doi.org/10.1016/S0006-3495\(77\)85544-6](https://doi.org/10.1016/S0006-3495(77)85544-6).
  41. Ames GF-L, Liu CE, Joshi AK, Nikaido K. 1996. Liganded and unliganded receptors interact with equal affinity with the membrane complex of periplasmic permeases, a subfamily of traffic ATPases. *J Biol Chem* 271:14264–14270. <https://doi.org/10.1074/jbc.271.24.14264>.
  42. Ramos JL, Martínez-Bueno M, Molina-Henares AJ, Terán W, Watanabe K, Zhang X, Gallegos MT, Brennan R, Tobes R. 2005. The TetR family of transcriptional repressors. *Microbiol Mol Biol Rev* 69:326–356. <https://doi.org/10.1128/MMBR.69.2.326-356.2005>.
  43. Kendall SL, Burgess P, Balhana R, Withers M, ten Bokum A, Lott JS, Gao C, Uhia-Castro I, Stoker NG. 2010. Cholesterol utilization in mycobacteria is controlled by two TetR-type transcriptional regulators: *kstR* and *kstR2*. *Microbiology* 156:1362–1371. <https://doi.org/10.1099/mic.0.034538-0>.
  44. Trautwein K, Grundmann O, Wöhlbrand L, Eberlein C, Boll M, Rabus R. 2012. Benzoate mediates repression of *C*₄-dicarboxylate utilization in “*Aromatoleum aromaticum*” EbN1. *J Bacteriol* 194:518–528. <https://doi.org/10.1128/JB.05072-11>.
  45. Heider J, Boll M, Breese K, Breinig S, Ebenau-Jehle C, Feil U, Gad'on N, Laempe D, Leuthner B, Mohamed ME-S, Schneider S, Burchhardt G, Fuchs G. 1998. Differential induction of enzymes involved in anaerobic metabolism of aromatic compounds in the denitrifying bacterium *Thauera aromatica*. *Arch Microbiol* 170:120–131. <https://doi.org/10.1007/s002030050623>.
  46. Barragán MJL, Carmona M, Zamarro MT, Thiele B, Boll M, Fuchs G, García JL, Díaz E. 2004. The *bzd* gene cluster, coding for anaerobic benzoate catabolism, in *Azoarcus* sp strain CIB. *J Bacteriol* 186:5762–5774. <https://doi.org/10.1128/JB.186.17.5762-5774.2004>.
  47. Zimmermann T, Sorg T, Siehler SY, Gerischer U. 2009. Role of *Acinetobacter baylyi* Crc in catabolite repression of enzymes for aromatic

- compound catabolism. *J Bacteriol* 191:2834–2842. <https://doi.org/10.1128/JB.00817-08>.
48. Galán B, Kolb A, Sanz JM, García JL, Prieto MA. 2003. Molecular determinants of the *hpa* regulatory system of *Escherichia coli*: the HpaR repressor. *Nucleic Acids Res* 31:6598–6609. <https://doi.org/10.1093/nar/gkg851>.
  49. Roper DJ, Fawcett T, Cooper RA. 1993. The *Escherichia coli* C homoproteocatechuate degradative operon: *hpc* gene order, direction of transcription and control of expression. *Mol Gen Genet* 237:241–250.
  50. Barragán MJL, Blázquez B, Zamarro MT, Mancheño JM, García JL, Díaz E, Carmona M. 2005. BzdR, a repressor that controls the anaerobic catabolism of benzoate in *Azoarcus* sp. CIB, is the first member of a new subfamily of transcriptional regulators. *J Biol Chem* 280:10683–10694. <https://doi.org/10.1074/jbc.M412259200>.
  51. Durante-Rodríguez G, Valderrama JA, Mancheño JM, Rivas G, Alfonso C, Arias-Palomo E, Llorca O, García JL, Díaz E, Carmona M. 2010. Biochemical characterization of the transcriptional regulator BzdR from *Azoarcus* sp. CIB. *J Biol Chem* 285:35694–35705. <https://doi.org/10.1074/jbc.M110.143503>.
  52. Lahme S, Trautwein K, Strijkstra A, Dörries M, Wöhlbrand L, Rabus R. 2014. Benzoate mediates the simultaneous repression of anaerobic 4-methylbenzoate and succinate utilization in *Magnetospirillum* sp. strain pMbN1. *BMC Microbiol* 14:269. <https://doi.org/10.1186/s12866-014-0269-4>.
  53. Davies SJ, Golby P, Omrani D, Broad SA, Harrington VL, Guest JR, Kelly DJ, Andrews SC. 1999. Inactivation and regulation of the aerobic C_4 -dicarboxylate transport (*dctA*) gene of *Escherichia coli*. *J Bacteriol* 181:5624–5635. <https://doi.org/10.1128/JB.181.18.5624-5635.1999>.
  54. Busch A, Lacal J, Silva-Jimenez H, Krell T, Ramos JL. 2010. Catabolite repression of the TodS/TodT two-component system and effector-dependent transphosphorylation of TodT as the basis for toluene dioxygenase catabolic pathway control. *J Bacteriol* 192:4246–4250. <https://doi.org/10.1128/JB.00379-10>.
  55. Rojo F. 2010. Carbon catabolite repression in *Pseudomonas*: optimizing metabolic versatility and interactions with the environment. *FEMS Microbiol Rev* 34:658–684. <https://doi.org/10.1111/j.1574-6976.2010.00218.x>.
  56. Wöhlbrand L, Rabus R. 2009. Development of a genetic system for the denitrifying bacterium "*Aromatoleum aromaticum*" strain EbN1. *J Mol Microbiol Biotechnol* 17:41–52. <https://doi.org/10.1159/000159194>.
  57. Sanger F, Nicklen S, Coulson AR. 1977. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* 74:5463–5467. <https://doi.org/10.1073/pnas.74.12.5463>.
  58. Koßmehl S, Wöhlbrand L, Drüppel K, Feenders C, Blasius B, Rabus R. 2013. Subcellular protein localization (cell envelope) in *Phaeobacter inhibens* DSM 17395. *Proteomics* 13:2743–2760. <https://doi.org/10.1002/pmic.201300112>.
  59. Oelmüller U, Krüger N, Steinbüchel A, Friedrich CG. 1990. Isolation of prokaryotic RNA and detection of specific mRNA with biotinylated probes. *J Microbiol Methods* 11:73–84. [https://doi.org/10.1016/0167-7012\(90\)90050-G](https://doi.org/10.1016/0167-7012(90)90050-G).
  60. Ramakers C, Ruijter JM, Lekanne Deprez RH, Moorman AFM. 2003. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett* 339:62–66. [https://doi.org/10.1016/S0304-3940\(02\)01423-4](https://doi.org/10.1016/S0304-3940(02)01423-4).
  61. Kawaguchi A, Yoshimura T, Okuda S. 1981. A new method for the preparation of acyl-CoA thioesters. *J Biochem* 89:337–339. <https://doi.org/10.1093/oxfordjournals.jbchem.a133207>.
  62. Bradford MM. 1976. A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3).
  63. Zech H, Hensler M, Koßmehl S, Drüppel K, Wöhlbrand L, Trautwein K, Hulsch R, Maschmann U, Colby T, Schmidt J, Reinhardt R, Schmidt-Hohagen K, Schomburg D, Rabus R. 2013. Adaptation of *Phaeobacter inhibens* DSM 17395 to growth with complex nutrients. *Proteomics* 13:2851–2868. <https://doi.org/10.1002/pmic.201200513>.