Published in final edited form as:

J Allergy Clin Immunol. 2021 May ; 147(5): 1838–1854.e4. doi:10.1016/j.jaci.2020.10.045.

Suppression of IgE-mediated anaphylaxis and food allergy with monovalent anti-FcεRIα monoclonal antibodies

Marat V. Khodoun, DVM, PhD¹, Suzanne C. Morris, PhD¹, Wen-Hai Shao, PhD¹, Crystal Potter, BA¹, Elizabeth Angerman, BS², Artem Kiselev, PhD³, Alexander E. Yarawsky, PhD², Andrew B. Herr, PhD^{2,4,5}, Katja Klausz, PhD⁶, Anna Otte, PhD⁶, Matthias Peipp, PhD^{#6}, Fred D. Finkelman, MD^{#1,2,†}

¹Division of Allergy, Immunology and Rheumatology, Department of Internal Medicine, University of Cincinnati College of Medicine, Cincinnati, OH 45267;

²Division of Immunobiology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229;

³Institute of Cytology, Russian Academy of Sciences, Saint Petersburg, 194064, Russia

⁴Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH 45229, USA;

⁵Division of Infectious Diseases, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229;

⁶Division of Stem Cell Transplantation and Immunotherapy, University Hospital Schleswig-Holstein, University of Kiel, 24105 Kiel, Germany.

[#] These authors contributed equally to this work.

Abstract

Background: Mast cell and basophil activation by antigen crosslinking of FcεRI-bound IgE is central to allergy pathogenesis. We previously demonstrated global suppression of this process by rapid desensitization with anti-FcεRIα mAbs.

Objective: Determine whether use of monovalent (mv) anti-FcεRIα mAbs increases desensitization safety without loss of efficacy.

Methods: mv anti-human (hu) FcεRIα mAbs were produced with mouse-derived immunoglobulin V regions and huIgG1 or huIgG4 C regions, and used to suppress murine IgE-mediated anaphylaxis and food allergy. mAbs were administered as a single dose or as serially increasing doses to mice that express hu instead of mouse FcεRIα, mice that additionally have an allergy-promoting IL-4Rα mutation, and hu cord blood-reconstituted immunodeficient, hu cytokine-secreting, mice that have large numbers of activated hu mast cells. Anaphylaxis

[†]Corresponding author: finkelfd@ucmail.uc.edu, Postal address: Apartment 79, 26 Sderot Nitza, Netanya, Israel, 4226259, Telephone: 513 427-3241, FAX: None available.

Declaration of Interest Statement: SCM, MVK, CP, ABH, and FDF have rights in a patent for rapid desensitization with anti-FcεRIα mAbs that was awarded to the University of Cincinnati and could benefit from the licensing of this patent. FDF is a principal in a start-up company that is trying to commercialize rapid desensitization with anti-FcεRIα monoclonal antibodies. The other authors have no relevant conflicts of interest.

susceptibility was sometimes increased by treatment with IL-4 or a β -adrenergic receptor antagonist.

Results: mv anti-huFc ϵ RI α mAbs have considerably less ability than divalent mAbs to induce anaphylaxis and deplete mast cell and basophil IgE, but still strongly suppress IgE-mediated disease. The mv mAbs can be safely administered as a single large dose to mice with typical susceptibility to anaphylaxis, while a rapid desensitization approach safely suppresses disease in mice with increased susceptibility. Our huIgG4 variant of mv anti-huFc ϵ RI α mAb is safer than our huIgG1 variant, apparently because reduced interactions with Fc γ Rs decrease ability to indirectly crosslink Fc ϵ RI.

Conclusion: mv anti-Fc ϵ RI α mAbs more safely suppress IgE-mediated anaphylaxis and food allergy than divalent variants of the same mAbs. These mv mAbs may be useful for suppression of huIgE-mediated disease.

Capsule summary:

Treatment of humanized mice with monovalent monoclonal anti-Fc ϵ RI α antibody safely decreased mast cell IgE and responsiveness to IgE crosslinking, which together suppressed IgE-mediated anaphylaxis. Antibody safety was increased by removing aggregates and limiting Fc γ receptor binding.

Our studies in humanized mice provide evidence that treatment with a monovalent monoclonal anti-Fc ϵ RI α antibody that interacts poorly with Fc γ receptors can safely, rapidly, and effectively suppress IgE mediated anaphylaxis and food allergy.

Keywords

Mouse; humanized; mast cell; desensitization; Syk; crosslinking; IL-4; IgG1; IgG4

Introduction:

The IgE-Fc ϵ RI-mast cell/basophil axis provides a mechanism for the rapid release of vasoactive mediators, cytokines, and enzymes that quickly induce an inflammatory response¹. Although this response protects against helminth and insect parasites^{2–5}, it is also important in allergy pathogenesis, particularly those allergic disorders in which disease develops within minutes of allergen exposure^{6, 7}. The decreased importance of multicellular parasites and the increased prevalence of allergic disorders in developed countries has shifted the balance between the beneficial and adverse effects of this response and promoted the development of approaches to suppress the IgE-Fc ϵ RI-mast cell/basophil axis and its contributions to allergy.

One such approach is based on in vitro and in vivo evidence that IgE/Fc ϵ RI crosslinking does not invariably lead to mast cell/basophil activation and degranulation. Limited crosslinking, instead, has been associated with desensitization of these cells, characterized by the development of an anergic state in which even strong IgE/Fc ϵ RI crosslinking causes little or no activation and degranulation^{8–10}. This process provides the basis for the therapeutic procedure known as rapid, or rush desensitization, in which patients who require

a drug to which they have developed IgE-mediated allergy are treated with serially increasing doses of that drug, starting with a dose too small to induce disease and finishing with a full therapeutic dose^{11–14}.

During the past few years, we have used conventional, then humanized mouse models to study the feasibility of using a rapid desensitization approach that replaces a single allergen with a monoclonal antibody (mAb) to FcεRIα (the IgE-binding chain of the FcεRI heterotetramer¹⁵), with the goal of suppressing all IgE-mediated allergy rather than only disease caused by the single allergen^{12, 16}. Our studies have demonstrated the feasibility and efficacy of this approach, which, in addition to suppressing all IgE-mediated anaphylaxis was safer and longer lasting in our models than allergen-mediated desensitization¹². However, we also found that an extended period of treatment with very slow increases in anti-FcεRIα mAb dose was required to desensitize mice that were particularly susceptible to anaphylaxis¹⁶.

This limitation has led us to modify our approach by using an engineered monovalent (mv) anti-FcεRIα mAb for desensitization that has only one Fab, which is associated with complete Fc domains. We initially expected that because a mv antibody would bind to, but not crosslink its target, it would be unable to either activate or desensitize basophils and mast cells, but could slow FcεRIα crosslinking by a subsequently injected divalent (dv) mAb that had the same V regions and bound to the same FcεRIα epitope. Indeed, this approach works well. However, as also shown herein, injection of the dv mAb is not necessary; although the mv mAb has much less ability than the dv mAb to activate mast cells and cause disease, it can still desensitize mast cells by limited, indirect FcεRI crosslinking that is predominantly mediated through interactions of its Fc with FcγRs.

Methods:

Mice.

BALB/c mice were purchased from Taconic (Hudson, NY) or bred in the Cincinnati Children's Hospital Medical Center (CCHMC) animal facility. BALB/c background huFcεRIα mice, which lack mouse FcεRIα and express transgenic huFcεRIα; and huFcεRIα/F709 mice, which additionally express IL-4Rα with a mutation (F709) that allows only positive signaling by IL-4 and IL-13, were bred as previously described^{16–18}. These mice, unlike otherwise identical mice that lack the IL-4Rα F709 mutation, develop egg allergy in response to epicutaneous (e.c), followed by oral gavage (o.g.) with egg¹⁶. Human IL-3/SCF/GM-CSF transgenic, immune-deficient NOD/LtSz-SCID IL-2RG^{–/–} (NSGS) or NOD/Rag1^{–/–}/IL-2RG^{–/–} (NRGS) mice were bred at a CCHMC core and engrafted with T cell-depleted human cord blood cells as previously described^{16, 19, 20}. 8–12 week old mice of both sexes were used. All mouse studies were approved by the Cincinnati Children's Hospital Research Foundation Institutional Animal Care and Use Committee (IACUC).

Reagents.

Ovalbumin (OVA) and bovine serum albumin (BSA) were purchased from Sigma (St Louis, MO). TNP-OVA and TNP-BSA were prepared by haptening OVA or BSA, respectively, with 2,4,6-trinitrophenyl- ϵ -aminocaproyl-O-succinamide (Biosearch Technologies, Novato, CA). Propranolol was purchased from West Ward Pharmaceuticals, Eatontown, NJ). Egg antigen (Ag) was prepared as previously described¹⁶. Briefly, egg white (EW) and egg yolk, removed under sterile conditions from fresh organic chicken eggs, were pooled separately. The EW was dialyzed against distilled water, analyzed for protein concentration using a BCA Protein Assay Kit (Pierce, Waltham, MA), then pressure concentrated with a stirred ultrafiltration cell (Millipore, Temecula, CA) with a 30-kDa cut-off Diaflo membrane. The yolks were diluted 1:1 with 0.15M NaCl and centrifuged to remove egg yolk granules. The supernatant (egg yolk plasma (EYP)) was added to the EW while the EW was being concentrated to create a homogenous mixture with a final concentration of 333 mg of EW protein/ml. A long-acting formulation of IL-4 was produced by mixing the anti-IL-4 BVD4-1D11 mAb with recombinant murine IL-4 (Peprotech, Cranbury, NJ) at a 5:1 w/w ratio.

Antibodies.

Generation of a hybridoma that secretes IE7 mouse IgG1 anti-huFc ϵ RI α has been described¹⁶, as has generation of a hybridoma that secretes AER-37 (also called CRA-1) mouse IgG2b anti-human Fc ϵ RI α ²¹. IE7 and AER-37 bind the same or closely positioned epitopes, as shown by mutual crossblocking¹⁶. IE7 and AER-37 bind huFc ϵ RI α with similar affinities; however, AER-37 has considerably higher on- and off-rates than IE7¹⁶. A chimeric huIgG1 version of AER-37 was produced in our lab by subcloning synthetic coding DNA fragments (GeneArt, Thermo Fisher Scientific, Waltham, MA) into a pEE12.4 vector (Lonza, Basel, Switzerland). Transfected cells were seeded at low density into 96-well plates to obtain clones derived from single colonies. Clones were identified visually and then selected based on the amount of AER-37 detected in the culture medium by ELISA. CHO-K1 cells that stably expressed recombinant huIgG1 AER-37 were grown in glutamine-free medium in roller bottles. Cell culture medium was collected every 7 d and replaced with fresh medium. Conditioned medium was concentrated and dialyzed into pH 7.0 50 mM Tris / 0.15M NaCl, adsorbed to a column of GammaBind Plus Sepharose (GE Life Sciences, Marlborough, MA), and eluted with Gentle Ag/Ab Elution Buffer (Thermo Fisher Scientific, Waltham, MA). The eluate was dialyzed into 0.15M NaCl, concentrated using an Amicon stirred cell, and its concentration of huIgG1 AER-37 was determined by ELISA. Mouse IgE anti-TNP mAb (clone IGELa2, from ATCC), human IgE anti-TNP (clone NS0 1F3.5C5), rat IgG2b anti-mouse Fc γ RIII/RIIb (clone 24G2), and control rat IgG2b (clone J1.2) mAbs were produced as ascites²² and in some cases were purified by DEAE ion exchange or protein G affinity chromatography (Thermo Fisher Scientific, Waltham, MA) according to the manufacturer's instructions.

Generation of monovalent (mv) IE7 and AER-37 chimerized anti-huFc ϵ RI α mAb.

We generated a series of mv hIgG anti-huFc ϵ RI α recombinant constructs consisting of one full human κ light chain, one full human heavy chain and one truncated heavy chain

resulting in formation of mv mAbs that had a single Fab linked through a hinge region to a complete Fc. Preferential assembling of the truncated and full heavy chains was determined by four Fc amino acid exchanges: E356K, D399K, K392D, and K409D (Supp. Fig. 1A)²³. Recombinant mv antibodies were generated by transient expression in CHO-S cells, using a Maxcyte STX flow electroporation system (Maxcyte, Inc., Gaithersburg, MD) according to the manufacturer's instructions. A three step purification strategy was used that consisted of: (1) human CH1-specific affinity chromatography; (2) size exclusion chromatography; and (3) human κ light chain-specific affinity chromatography. mv mAb structure and molecular mass were validated by protein gel electrophoresis and western blotting (Supp. Fig. 1B–D) and size exclusion chromatography (Supp. Fig. 1E). These mv Fab/Fc mAbs were used instead of mv mAb Fab fragments because the former have a much longer in vivo half-life than the latter as a result of efficient FcRn recycling²⁴.

Detection of systemic anaphylaxis.

Anaphylaxis was detected as a change in mouse activity and the development of hypothermia. Mouse temperature was measured every 5 to 15 min with a rectal probe²⁵. Anaphylaxis-associated activity was scored on a 5 point scale, where a score of 0 indicates normal mobility; 1 = decreased spontaneous movement, 2 = spontaneous movement only after a pinch, 3 = slow or absent spontaneous movement after a pinch, 4 = inability to maintain upright posture, and 5 = death. Diarrhea was defined as profound liquid stool discharge, and was evaluated 30 min post-Ag challenge, unless otherwise indicated.

Measurement of serum levels of mouse mast cell protease 1 (MMCP1) and mouse mast cell protease 7 (mcpt7).

Serum levels of proteases were measured in blood drawn 4 hrs (unless other specified) after challenge. MMCP1 was evaluated with a commercial ELISA kit (Thermo Fisher Scientific, Waltham, MA), according to the manufacturer's instructions. mcpt7 was evaluated by an ELISA developed in our lab. ELISA plate wells were coated overnight with 0.2 mg/ml of a goat anti-mouse mcpt7 polyclonal Ab (clone AF1937, R&D Systems, Minneapolis, MN). Serial dilutions of serum samples or recombinant standard (R&D Systems, Minneapolis, MN) were loaded into wells and bound mcpt7 was detected with 0.2 mg/ml of a biotin-labeled goat anti-mouse mcpt7 polyclonal Ab (BAF1937, R&D Systems, Minneapolis, MN), followed by streptavidin peroxidase conjugate (Thermo Fisher Scientific, Waltham, MA) and SuperSignal™ ELISA Femto Substrate (Pierce, Waltham, MA) according to the manufacturer's instructions. Although both assays show the greatest effects of desensitization when serum is obtained within 30 min of challenge, they still show marked effects when serum is obtained up to 4 hrs after challenge. Obtaining serum at the latter time point was frequently necessary to simultaneously test for cytokine responses and because it is difficult to obtain serum when mice are recovering from shock.

Passive sensitization and induction of anaphylaxis.

Unless otherwise indicated, mice were sensitized by i.v. injection of 10 μ g of huIgE anti-TNP mAb and challenged i.v. the next d with 10 μ g of TNP-OVA.

Active sensitization of huFcεRIα/F709 mice.

Egg allergy was induced in huFcεRIα/F709 mice, as described¹⁶.

Immunization of mice with hulG4 IE7 and detection of mouse antibodies to hulG4 IE7.

huFcεRIα/F709 Tg mice were immunized intraperitoneally (i.p.) with 10 µg of dv hIgG4 IE7 in 0.1 ml 0.15M NaCl, adsorbed to an equal volume of Inject Alum (Thermo Fisher Scientific, Waltham, MA) twice at a one month interval. 1 wk after the second immunization mice were bled and sera were tested by ELISA for level of mouse IgG anti-human IgG4 Ab, using wells coated with recombinant dv hulG4 IE7. Bound Ab was detected with biotin-labeled anti-mouse IgGκ mAb (Clone RMK-12, Biolegend, San Diego, CA) followed by a streptavidin-peroxidase conjugate (Thermo Fisher Scientific, Waltham, MA) and SuperSignal™ ELISA Femto Substrate (Pierce, Waltham, MA) according to the manufacturer's instructions

Rapid desensitization.

Unless otherwise indicated, rapid desensitization was accomplished by i.p. injection of doubling doses of an anti-FcεRIα mAb every 2 hrs, starting at 50 µg and ending with 400 µg.

Cell preparation.

Peritoneal cells enriched in mast cells and blood, bone marrow, and splenic nucleated cells were prepared as described¹⁶.

Immunofluorescence and flow cytometry.

1×10^6 cells in 0.1 ml of Hank's balanced salt solution with 10% newborn calf serum and 0.2% sodium azide (HNA) were incubated with 1 µg of 2.4G2 or human IgG for 15 minutes on ice, then stained with 1 µg each of fluorochrome-labeled mAbs specific for mouse IgE (clone RME-1, Biolegend), hu IgE (clone MHE-18, Biolegend), or the anti-huFcεRIα mAbs AER-37 or 15.1, and/or mAbs to human c-kit (clone 104D2, Biolegend) or mouse c-kit (clone 2B8, Biolegend), and human IL-3 receptor (clone 6H6, Biolegend), or mouse IL-3 receptor (clone 5B11, Biolegend), mouse B220 (clone RA3-6B2, Biolegend) and/or human CD3 (clone OKT3, Biolegend), or mouse CD3 (clone 145-2C11) or mouse pan-NK cells marker CD49b (clone DX5, Biolegend); Syk (clone 4D10.2, Biolegend); phospho-Syk (clone mochi1ct, ThermoFisher); phospho-SHIP (clone P1C1-A5, Biolegend) Akt (C67E7, from Cell Signaling Tech.), phospho-AKT (D9E, Cell Signaling Tech.), phospho-ERK (clone 4B11B69, Biolegend). Cells were washed twice, fixed with 2% paraformaldehyde, and analyzed with a Becton-Dickinson LSR II Flow Cytometer (BD Biosciences). Mast cells were identified as high side scatter, c-kit positive, IL-3 receptor positive, CD3 negative and B220 negative cells. Basophils were identified as forward/side scatter intermediate, IL-3 receptor positive, CD49b positive, CD3 negative cells¹⁶.

IVCCA for IL-4.

IL-4 secretion was quantitated with the in vivo cytokine capture assay (IVCCA), as described²⁶.

Measurement of mast cell signaling.

After surface staining to identify cell types, peritoneal wash cells were fixed with 4% formaldehyde solution, permeabilized with permeabilization/wash buffer (R&D System, Minneapolis, MN) according to the manufacturer's instruction, stained with Abs specific for the intra-cellular targets, washed and analyzed by flow cytometry.

Detection of mAb aggregates and deaggregation procedure.

For deaggregation, mAbs were loaded onto a HiLoad 16/600 Superdex 200 pg (GE Healthcare, Marlborough, MA) column equilibrated in PBS. Fractions containing the pure monomer were pooled so that the final concentration of the pooled monomer remained high enough to perform functional in vivo experiments without needing to concentrate the material. Evaluation of the mAb aggregation was performed by sedimentation velocity analytical ultracentrifugation (AUC) testing with a Beckman Coulter XL-I (Beckman, Brea, CA) using absorbance optics at 280 nm. Samples were loaded into 1.2 cm, two-sector epon-charcoal centerpieces with sapphire windows. Evaluations were performed at 48,000 rpm in an An-60 Ti rotor at 20°C. Data analysis was done in SEDFIT version 16.1 using the continuous c(s) distribution model²⁷. SEDNTERP was used to calculate the partial specific volume, buffer density, and buffer viscosity of each protein in PBS buffer²⁸.

Statistics.

GraphPad Prism version 8.4.3 was used for statistical calculations. A 2-tailed Mann-Whitney U test, a Kruskal-Wallis test, a Tukey honest significant difference test (THSDT), and/or a 1-way ANOVA were used when appropriate to test for statistical significance. The tests used for each experiment are reported in figure legends. A p value of <.05 was considered significant.

Results:

mv anti-huFcεRIα mAb protects against dv anti-huFcεRIα mAb-induced shock.

Our initial experiment evaluated whether a mv anti-huFcεRIα mAb protects against anaphylaxis induction by a dv mAb that has the same V regions. Separate groups of huFcεRIα mice were treated with increasing doses of mv huIgG1 IE7 followed by 100 µg of dv huIgG1 IE7. We observed dose-dependent protection against hypothermia (Fig. 1A, left panel) and mast cell degranulation and basophil activation (detected, respectively, as decreased serum levels of mouse mast cell protease 1 (MMCP1) and interleukin-4, Fig. 1A, right panel). Sequential treatment with mv, followed by dv huIgG1 IE7 completely protected IgE anti-TNP mAb-sensitized mice from shock (hypothermia) induction by TNP-OVA (Fig. 1B). Surprisingly, single injections of mv huIgG1 IE7 and dv IE7 had similar abilities to remove IgE from mast cells at the doses used and times studied, even though IE7 does not compete with IgE for binding to FcεRI¹⁶ and although mv huIgG1 IE7 had much less ability than dv IE7 to activate mast cells and basophils (determined by MMCP1 and IL-4 levels, Fig. 1C). Protection against IgE-mediated anaphylaxis and mast cell activation could be maintained for at least 18 d in mice that had originally been treated with sequential mv huIgG1 IE7 and dv IE7 by repeated injections of the dv mAb (Fig. 1D).

mv hulgG1 IE7 safely protects against IgE-mediated anaphylaxis.

The ability of mv huIgG1 IE7 to remove IgE from mast cells and to modestly activate mast cells and basophils raised the possibility that this mv mAb might protect mice against IgE-mediated anaphylaxis directly, without a need to subsequently administer dv IE7. To test this possibility, IgE anti-TNP mAb-sensitized huFcεRIα mice were injected with mv huIgG1 IE7 or dv IE7, followed for the development of hypothermia, and 1 d later challenged with TNP-OVA. Although, as expected, dv, but not mv IE7 directly induced hypothermia, the mv and dv forms of this anti-huFcεRIα mAb both completely suppressed IgE-mediated anaphylaxis and strongly suppressed the mast cell (MMCP1) and basophil (IL-4) responses to Ag challenge (Fig. 2A). Further comparison of the in vivo effects of mv hulgG1 IE7 vs. dv IE7 showed that neither consistently depleted peritoneal or bone marrow mast cells, both depleted blood, but not bone marrow basophils, both removed most mast cell and basophil IgE (although the dv mAb did this more completely in this experiment), the dv mAb remained associated with FcεRI more than the mv mAb (when evaluated ex vivo by ability to block the binding of the anti-huFcεRIα mAb AER-37, which cross-blocks IE7¹⁶), and the dv mAb was more effective than the mv mAb at decreasing the expression of mast cell surface FcεRIα that is not associated with IgE (determined by the ex vivo binding of 15.1, an anti-huFcεRIα mAb that competes with IgE for binding and is not crossblocked by IE7^{16, 29}) (Fig 2B). Kinetic experiments revealed that a single injection of mv hulgG1 IE7 protected IgE anti-TNP mAb-sensitized mice against anaphylactic shock (hypothermia) in 2 hrs or less, with protection lasting for at least 48 hrs (Fig. 2C, upper left panel). Protection coincided with mast cell loss of IgE (Fig. 2C, lower left panel), but persisted even when mast cells had regained considerable surface IgE and baseline levels of free FcεRIα (Fig. 2C, lower middle panel). mv mAb treatment had no significant effect on peritoneal mast cell number (Fig. 2C, upper middle panel). Thus; loss of mast cells or mast cell IgE cannot fully explain the suppressive effects of our mv mAb on anaphylaxis; an IgE-independent decrease in mast cell responsiveness (desensitization, or anergy) is also important. This is particularly apparent when mast cell surface IgE (Fig. 2C, lower left panel) is compared with the MMCP1 response 48 hours after mv mAb injection (Fig. 2C, lower right panel). Surprisingly, although anaphylaxis was strongly suppressed 2 hrs after mv mAb administration, the MMCP1 response to Ag challenge was not significantly suppressed at that time (compare Fig. 2C upper left and lower right panels). This is probably explained by the development of hypothermia 5–10 minutes after Ag challenge, while we measured MMCP1 levels 4 hours after challenge. Consistent with this explanation, we find that desensitization has a considerably greater suppressive effect on the release of MMCP1 by mucosal mast cells and the release of mcpt7 by connective tissue mast cells³⁰ 2 minutes after Ag challenge than 4 hours later (Supplemental Fig. 2). Thus, desensitization with a single dose of mv anti-huFcεRIα mAb suppresses IgE-mediated anaphylaxis by delaying mast cell degranulation as well as by decreasing it.

Mast cell SYK and SHIP phosphorylation are suppressed by mv hulgG1 IE7.

In vitro mast cell desensitization has been explained, at least partially, by decreases in the total quantity and activation by phosphorylation of the tyrosine kinase Syk, which has an early and critical role in IgE-mediated mast cell activation and degranulation³¹. Consequently, we evaluated the effects of in vivo mv anti-huFcεRIα mAb treatment on total

mast cell Syk as well as the activation (phosphorylation) of Syk and 2 other relevant signaling molecules, ERK and AKT, as well as on the inhibitory phosphatase SHIP³². mv mAb treatment significantly decreased the phospho-Syk and phospho-SHIP responses to Ag challenge, but had no significant effects on phospho-AKT or ERK and had no significant effect on total Syk levels (Fig. 3). These results are consistent with mv anti-huFcεRIα mAb at least partially desensitizing mast cells by inhibiting Syk activation.

Aggregation and FcγRs contribute to mast cell activation by mv anti-FcεRIα mAbs.

As far as is known, signaling through FcεRI requires aggregation or crosslinking of this receptor^{33–35}, although it is difficult to prove that signaling is never induced by ligand binding without crosslinking³⁶. A crosslinking requirement explains why mv anti-huFcεRIα mAb is much less potent than the dv mAb at activating mast cells, but does not explain how some mast cell and basophil activation and degranulation is induced by the mv mAb. One simple possibility is spontaneous in vitro mv mAb aggregation, which should allow crosslinking by making the mv mAb divalent or oligovalent for huFcεRIα. Consistent with this, dimers and larger aggregates were detectable by analytical ultracentrifugation and analytical gel filtration and could be removed by preparative gel filtration (Supplemental Fig. 3). These aggregates contributed to the toxicity of mv huIgG1 IE7, as shown when highly susceptible, egg-allergic huFcεRIα/F709 mice were injected i.v. with 100 μg of deaggregated or non-deaggregated mv huIgG1 IE7 (Fig. 4A, left panel). Although injection of the non-deaggregated mv mAb caused an average temperature drop of ~8°C with the death of 7 of 8 injected mice, injection of the same quantity of the deaggregated mAb caused a reversible temperature drop of ~4°C. However, the deaggregated mv mAb retained the ability to desensitize mice, as shown by complete inhibition of hypothermia and diarrhea when the treated mice were challenged with egg the next day by oral gavage (o.g.) (Fig. 4A, center panel). The deaggregated mv mAb also caused considerable loss of mast cell surface IgE (Fig. 4A, right panel). Thus, mv anti-huFcεRIα mAb aggregation appears to explain some, but not all, of this mAb's toxicity when injected into highly sensitized mice, but does not appear to contribute to mast cell desensitization.

In addition to aggregation, mv mAbs have been shown to crosslink their ligands indirectly by interacting through their Fc domains with FcγRs³⁷. To determine the importance of this mechanism in mv mAb toxicity and desensitization, we sensitized huFcεRIα/F709 mice with IgE anti-TNP mAb, and made them particularly susceptible to anaphylaxis by treating them with a long-acting form of IL-4 (IL-4C) and propranolol^{38, 39} (Supplemental Fig. 4). To block mv mAb interactions with FcγRs, some mice were also pre-treated with 2.4G2, a rat IgG2b mAb that directly blocks mouse FcγRIIB and FcγRIII and indirectly inhibits FcγRI and FcγRIV by binding them through its Fc domains³⁹. 2.4G2 treatment blocked the toxicity of deaggregated mv, but not dv anti-huFcεRIα mAb and considerably inhibited the ability of the mv, but not the dv mAb to suppress IgE-mediated anaphylaxis (Fig. 4B). This suggests that indirect, FcγR-mediated FcεRI crosslinking by mv huIgG1 IE7 is important for both the toxic and therapeutic effects of this mAb.

As an additional way to evaluate the importance of mv anti-huFcεRIα mAb - Fc interactions with FcγRs for mAb toxic and therapeutic effects, we engineered mv IE7 mAb that had Ab

C regions derived from huIgG1, as before, or from huIgG4, which binds less strongly than huIgG1 to both mouse and hu FcγRs^{40, 41}. Consistent with the results of our anti-FcγR blocking experiment, mv huIgG4 IE7 was much less toxic than mv huIgG1 IE7 when injected i.v into IL-4C and propranolol pre-treated huFceRIα mice (Fig. 4C, left panel), but also had less ability to suppress IgE-mediated anaphylaxis (Fig. 4C, right panel). Intravenous injection of a higher dose of mv huIgG4 IE7 into propranolol- + IL-4C-pre-treated huFceRIα mice could induce hypothermia, particularly if the mv mAb were not deaggregated prior to injection (Fig. 4D). A dose-response study showed that no single dose of mv huIgG4 IE7 could be injected i.v. into highly susceptible IL-4C pre-treated huFceRIα/F709 mice that had no toxicity, but completely protected against IgE-mediated anaphylaxis (Fig. 4E, left 2 panels). In the latter experiment, increasing the dose of mv mAb injected had no consistent effect on peritoneal mast cell number, but decreased mast cell surface IgE (Fig. 4E, right 2 panels). Pre-treatment of mice with 2.4G2 has little effect on the small hypothermic response induced by injection of 400 μg of mv huIgG4 IE7 (Figure 4F, left panel), but blocked the ability of this mv mAb to inhibit IgE-mediated anaphylaxis (Fig. 4F, right panel). Taken together, our results show that FceRI crosslinking by mv anti-huFceRIα mAb contributes to toxicity whether mediated through mv mAb aggregation or by interaction with FcγRs, whereas FcγR interactions, but not aggregation, contribute to mast cell desensitization.

Rapid desensitization and prolonged treatment with mv huIgG4 IE7 safely protect against IgE-mediated anaphylaxis.

Although injection of mice with mv huIgG4 IE7 was much safer than injection with dv IE7, the mv mAb still induced toxicity when injected into highly susceptible mice in a single dose that could totally suppress IgE-mediated anaphylaxis in these mice. This led us to test whether using the mv mAb with our previously validated rapid desensitization procedure¹⁶ could fully desensitize mice without toxicity and whether rapid desensitization with a mv mAb had any advantage over the same procedure with a dv mAb. An initial experiment showed that rapid desensitization with just 4 doses of the mv mAb permitted injection of 400 μg of the same mv mAb without detectable toxicity, although injection of 400 μg of this mAb into mock desensitized mice caused an ~7°C temperature drop (Fig. 5A, upper 2 panels) and that safe desensitization with mv IE7 could be accomplished much more quickly than safe desensitization with dv IE7 (compare Fig. 5A with the results in our previous paper¹⁶). Both the rapid desensitization and the single dose approaches prevented lethal IgE-mediated anaphylaxis (Fig. 5A, lower left and center panels), although rapid desensitization was more effective than a single mv mAb dose at removing mast cell IgE (Fig. 5A, lower right panel). Rapid desensitization with mv huIgG4 IE7 also safely prevented hypothermia and considerably decreased mast cell degranulation (MMCP1), IgE, and the incidence of diarrhea in our food allergy model (Fig. 5B). An additional experiment demonstrated that repeated injection of 400 μg of mv huIgG4 IE7 every 2 d for 7 d after rapid desensitization safely increased the protective effect of rapid desensitization with the same mv mAb in very susceptible mice and increased removal of mast cell surface IgE (Fig. 5C).

mv hulG4 IE7 desensitizes human mast cells in vivo.

To determine whether mv huIgG4 IE7 could safely desensitize hu mast cells in vivo, reNSGS mice were sensitized with huIgE anti-TNP mAb by injecting them i.p., then i.v. with 100 µg doses of this mAb or control mAb, and challenging them i.v. 1 day later with TNP-OVA (Fig. 6A). This short procedure was relatively ineffective at removing IgE from mast cells (Fig. 6A, left panel), and 1 of the 5 treated mice died following the mv mAb treatment while the remaining 4 developed considerable hypothermia (Fig. 6A, center panel). Two of the 4 surviving treated mice showed complete protection against Ag challenge, while one became hypothermic and one died 40 min after Ag challenge; in contrast, all of the control mAb-treated mice died 5–10 min after Ag challenge (Fig. 6A, right panel). To improve safety and efficacy, IgE-sensitized reNSGS mice were first rapidly desensitized with 25–400 µg of mv huIgG4 IE7, followed 3 d later by a second 400 µg dose of the same mv mAb and challenged one day after that with TNP-OVA. These mice developed no toxicity and were completely protected against IgE-mediated anaphylaxis, while 3 of 5 control mAb-treated mice developed lethal anaphylaxis in response to TNP-OVA challenge and the other 2 developed severe hypothermia and loss of mobility (Fig. 6B).

Anti-hulG4 Abs does not increase mv hulG4 IE7 mAb toxicity.

One concern about the therapeutic use of mv anti-huFcεRIα mAbs is that some patients might develop anti-therapeutic mAb Abs, which, theoretically, might aggregate the mv mAbs into polyvalent complexes that could induce mast cell degranulation and anaphylaxis. We initially tried to test this possibility by repeatedly injecting mice i.p. with mv huIgG4 IE7; however, although this mv mAb is foreign to mice, it failed to induce anti-mAb Ab development. Because of this, we immunized mice i.p. twice with the same mv mAb adjuvanted with alum; this led to an at least 500-fold increase in anti-mAb titer (Fig. 7A, upper left panel). These mice and mice inoculated i.p., as a control, with mouse IgG/alum, were then sensitized with IgE anti-TNP mAb, desensitized with mv huIgG4 IE7 or left undesensitized, and challenged with TNP-OVA. Desensitization failed to induce toxicity in mice that had pre-existing anti-mAb Abs (Fig 7A, upper right panel), although these mice were less protected against Ag-induced IgE-mediated anaphylaxis and developed higher MMCP1 levels than desensitized mice that lacked pre-existing anti-mAb Abs (Fig. 7A, lower panels). Because mv anti-huFcεRIα mAb-treated patients might develop anti-mAb Abs after desensitization and before receiving a large, single dose of the mv mAb to sustain mast cell suppression, a second experiment was performed in which mice that had first been induced to develop anti-mv mAb Abs were sensitized with IgE anti-TNP mAb, rapidly desensitized with mv huIgG4 IE7, injected with 400 µg of the same mv mAb 3 d later, and challenged with TNP-OVA 1 d after that. No adverse effects were induced by either the rapid desensitization (as with the first experiment) or the subsequent injection of 400 µg of the mv mAb and the treatment with this mAb still removed some surface IgE from mast cells and partially protected against IgE-mediated anaphylaxis (Fig. 7B). Thus, although anti-mv mAb Abs decrease the efficacy of mv anti-huFcεRIα mAbs, presumably by decreasing their effective concentration, they do not increase the toxicity of mv anti-huFcεRIα mAb therapy.

In vivo potency of mv anti-FcεRIα mAb is determined by its rate of binding to huFcεRIα.

The potency of a therapeutic mAb is influenced by the strength and speed of its interactions with its ligand⁴². To evaluate the effect of the binding kinetics of anti-huFcεRIα mAbs on in vivo toxicity and therapeutic efficacy, we produced two mv anti-huFcεRIα mAbs that had identical huIgG4 C regions but had V regions derived from either of 2 anti-huFcεRIα mAbs: IE7 and AER-37. These mAbs bind to identical or closely positioned huFcεRIα epitopes with similar affinity; however AER-37 has a considerably greater on-rate and off-rate than IE7¹⁶. In vivo, mv huIgG4 AER-37 was ~4 times as toxic as mv huIgG4 IE7 (Fig. 8A, upper panels and 8B, left panel), but was >4 times as potent at blocking IgE-mediated anaphylaxis (Fig. 8A, lower panels and 8B, right panel). Neither mv mAb decreased the percent of peritoneal mast cells (Fig. 8C, left panel, and mv huIgG4 AER-37 was 2–4-fold more potent than mv huIgG4 IE7 at removing mast cell surface IgE (Fig. 8C, right panel). The two mv mAbs had similar in vivo half-lives, even when mv huIgG4 AER-37 was injected into mice at one fourth the dose of mv huIgG4 IE7, to adjust for their differences in in vivo potency (Fig. 8D). Taken together, these results indicate that on-rate is more predictive than affinity or off-rate of the in vivo potency of mv anti-huFcεRIα mAbs and suggest that a therapeutic mAb with AER-37 V regions would have the advantage over one with IE7 V regions of being effective in lower quantity.

Discussion:

The data presented here demonstrate that a mv anti-huFcεRIα mAb composed of one Fab linked to a complete Fc has advantages over an intact dv anti-huFcεRIα mAb that has the same variable (V) regions for suppression of IgE-mediated anaphylaxis and food allergy. Specifically: (1) mv anti-FcεRIα mAb is much less potent than dv anti-FcεRIα mAb at inducing anaphylaxis; (2) pretreatment with mv anti-FcεRIα mAb blocks the ability of a dv anti-FcεRIα mAb that has the same V regions to induce anaphylaxis; (3) mv anti-FcεRIα mAb protects mice against IgE-mediated anaphylaxis and food allergy; (4) mv anti-FcεRIα mAb's already limited ability to induce anaphylaxis can be decreased further by administration through a rapid desensitization approach, with safe desensitization requiring considerably less time than the same procedure with a dv mAb; (5) the ability of mv anti-FcεRIα mAb to suppress IgE-mediated anaphylaxis can be safely extended by repeated administration; and (6) preformed antibodies to mv anti-FcεRIα mAb decrease efficacy but do not pose a safety risk. These results suggest that mv anti-FcεRIα mAb would be a safer therapy for IgE-related disease than either dv anti-FcεRIα mAb or desensitization with the disease-causing allergen, without loss of either efficacy or ability to suppress disease.

In addition to these translational observations, our results support the hypothesis that prolonged, limited crosslinking of FcεRI leads to mast cell desensitization rather than to activation. In this regard, the effects of a mv anti-FcεRIα mAb somewhat resemble those of a very low affinity dv anti-IgE mAb, which also has been described to deactivate, rather than activate mast cells⁴³, presumably by usually dissociating from one FcεRI-linked IgE molecule before it can bind to a second and crosslink FcεRI. However, targeting FcεRI rather than IgE has the theoretical advantage of not being influenced by the large range of IgE concentrations that are found in different allergic individuals⁴⁴.

In addition to their translational importance, our results increase understanding of the mechanisms involved in FcεRI-mediated mast cell activation and desensitization by a mv anti-FcεRIα mAb. Specifically, they demonstrate that: (1) treatment with mv anti-FcεRIα mAbs decreases both mast cell IgE expression and mast cell responsiveness to crosslinking of the remaining IgE; (2) the induction of decreased mast cell responsiveness by mv anti-FcεRIα mAb is associated with decreased levels of phospho-Syk, the activated form of a tyrosine kinase that is essential for mast cell activation^{31, 45}; (3) the limited ability of mv anti-FcεRIα mAb to induce anaphylaxis largely depends on aggregation and interactions with FcγRs; (4) mv anti-FcεRIα mAb treatment totally prevents the degranulation of some mast cells, but only delays the degranulation of others (as shown by serum MMCP1 and mcpt7 levels); and (5) mv anti-FcεRIα mAb potency correlates better with its on-rate than with its affinity or off-rate.

These translational and mechanistic observations both increase understanding and raise questions about how a mv anti-FcεRIα mAb suppresses IgE-mediated allergy and the optimal design of a therapeutic mv mAb. Most of our observations are consistent with the hypothesis that some FcεRI crosslinking is required for our mv mAb's therapeutic and residual untoward effects: (1) The removal of mAb aggregates prior to their administration decreases the ability of our mv mAbs to induce anaphylaxis, but has no observable effect on ability to suppress IgE-mediated anaphylaxis. This may be explained kinetically; FcεRIα-mediated induction of anaphylaxis occurs in a few minutes²⁵, while its induction of decreased mast cell responsiveness does not occur immediately and increases over hours to days. Because IgG aggregates are rapidly removed from circulation^{46–48}, they would be expected to have the ability to promote anaphylaxis to a much greater extent than they could promote unresponsiveness. This has a practical implication: administration through a route that would lead to relatively slow absorption (e.g.; subcutaneous administration) should minimize the adverse effects of aggregated mv mAbs. (2) Indirect crosslinking of FcεRI by mv mAbs is induced through mv mAb interactions with FcγRs. This is shown both by decreases in mv mAb adverse effects and efficacy when our mAbs are engineered with Fc domains that have decreased binding to FcγRs (i.e.; huIgG4 instead of huIgG1⁴¹) and by the effects of blocking FcγRs with the anti-FcγR mAb 2.4G2. Although these results are robust, they have limitations: huIgG4 retains some ability to interact with FcγRs⁴⁹ and we cannot exclude the possibility that 2.4G2 suppresses both anaphylaxis and induction of hyporesponsiveness through its interaction with the inhibitory receptor FcγRIIB^{39, 50}. Studies with mv anti-FcεRIα mAbs or mv fusion proteins that do not bind at all to FcγRs should be helpful, but even these may have undetectable, but sufficient levels of aggregated mAb to induce some level of FcεRIα crosslinking; it is also possible that mv anti-FcεRIα mAbs associate with each other after binding to mast cells. Thus, although crosslinking of FcεRIα is undoubtedly the main mechanism behind mv mAb activation and desensitization, it is difficult to rule out the possibility that binding without crosslinking has some effect. These same considerations are relevant to the issues of how mv anti-FcεRIα mAbs cause mast cell loss of IgE, even though the mAbs used do not directly interfere with IgE binding to FcεRI¹⁶, and how deaggregated IgE increases mast cell FcεRI expression^{36, 51, 52}.

Additional, related issues raised by our observations are the importance of Syk in mast cell desensitization by mv anti-FcεRIα mAbs, the variable extent of mast cell desensitization,

and the importance of mAb affinity for FcεRIα in mast cell activation and desensitization. Syk is rapidly activated through tyrosine phosphorylation by mast cell or basophil FcεRI crosslinking and is essential for FcεRI-mediated induction of mast cell degranulation^{31, 45}. Unpublished in vivo work by our group (S. Morris et al., submitted for publication) shows that a Syk phosphorylation inhibitor blocks anti-FcεRIα mAb-induced mast cell desensitization as well as mast cell activation, although the opposite observation has been reported with in vitro studies⁸. Thus, either the small amount of FcεRIα crosslinking that is induced by our mv mAbs is sufficient to phosphorylate Syk, or this event can be induced to a sufficient extent by mv mAb binding without FcεRI crosslinking.

Our kinetic study of MMCP1 and mcpt7 responses to FcεRI crosslinking in IgE-sensitized, mv anti-FcεRIα mAb-treated mice demonstrated that although mv mAb treatment decreased levels of both the mucosal and connective tissue mast cell-associated proteases at all time points studied, percent suppression was greater shortly after Ag challenge than it was later. Thus, while mv mAb desensitization totally prevented degranulation in some mast cells, it only delayed it in others. This demonstrates that IgE-mediated desensitization is not an all or none processes, and may relate to different degrees of inhibition of activation of Syk or other signaling molecules in different mast cells. In this regard, mast cell desensitization resembles anergy induced in B lymphocytes by exposing them to their cognate Ag in the absence of a second stimulatory signal⁵³, a process that also involves Syk⁵⁴⁻⁵⁶. However, because the induction of anaphylaxis requires rapid, relatively massive mast cell degranulation^{6, 25, 57} and anaphylaxis can be totally suppressed in our studies by mv anti-FcεRIα mAb treatment, delaying and spacing out mast cell degranulation appears to be sufficient to protect against anaphylaxis.

Our comparison of mv huIgG4 IE7 with mv huIgG4 AER-37 indicates that the latter mv mAb is ~4 times as potent as the former at both directly inducing anaphylaxis and at desensitizing mice, even though IE7 has a slightly higher affinity than AER-37¹⁶. This slight difference in affinity, as measured by surface plasmon resonance (SPR), reflects a considerably greater on-rate for AER-37 that is more than balanced by an even greater off-rate. Thus, to the extent to which SPR results reflect in vivo mv mAb binding and dissociation with a cell-associated ligand, the rate of binding of a mv mAb to FcεRI appears to be considerably more important than the duration of binding. This suggests that mv anti-FcεRIα mAb effects that contribute to anaphylaxis induction and desensitization occur very rapidly, before the rate of dissociation becomes consequential, at least within the range encompassed by our two mAbs. Taken together with our studies of the importance of FcγR binding, these observations suggest that an optimal therapeutic mv anti-FcεRIα mAb should have a relatively high on-rate, low or absent FcγR binding, and resist aggregation. Interestingly, ELISA measurements that correlate with mAb binding characteristics, unlike SPR measurements, are consistent with a 4-fold greater potency for AER-37 as compared to IE7¹⁶; thus, characterization of mAb binding by ELISA may be more useful than characterization of by SPR for predicting the in vivo potency of rapidly acting therapeutic mAbs.

An additional important consideration is the extent to which our humanized mouse model observations apply to humans. We have attempted to make our studies as applicable as

γ_c	the cytokine receptor common γ chain, also known as IL-2 receptor γ chain
d	day
dv	divalent
hr	hour
hu	human
mAb	monoclonal antibody
min	minute
mv	monovalent
reNSGS mice	NSGS mice that have been reconstituted with T cell-depleted human cord blood cells
NSGS mice	NOD background SCID mice that lack γ_c and express transgenic human GM-CSF, IL-3, and stem cell factor (kit ligand)
o.g.	oral gavage
R	receptor
THSDT	Tukey's honest significant difference test

References:

1. Finkelman FD, Rothenberg ME, Brandt EB, Morris SC, Strait RT. Molecular mechanisms of anaphylaxis: lessons from studies with murine models. *J Allergy Clin Immunol* 2005; 115:449–57. [PubMed: 15753886]
2. Gurish MF, Bryce PJ, Tao H, Kisselgof AB, Thornton EM, Miller HR, et al. IgE enhances parasite clearance and regulates mast cell responses in mice infected with *Trichinella spiralis*. *J Immunol* 2004; 172:1139–45. [PubMed: 14707089]
3. Finkelman FD, Urban JF Jr. The other side of the coin: the protective role of the TH2 cytokines. *J Allergy Clin Immunol* 2001; 107:772–80. [PubMed: 11344341]
4. Grecis RK, Else KJ, Huntley JF, Nishikawa SI. The in vivo role of stem cell factor (c-kit ligand) on mastocytosis and host protective immunity to the intestinal nematode *Trichinella spiralis* in mice. *Parasite Immunol* 1993; 15:55–9. [PubMed: 7679484]
5. Matsuda H, Watanabe N, Kiso Y, Hirota S, Ushio H, Kannan Y, et al. Necessity of IgE antibodies and mast cells for manifestation of resistance against larval *Haemaphysalis longicornis* ticks in mice. *J Immunol* 1990; 144:259–62. [PubMed: 2295794]
6. Altman LC. Basic immune mechanisms in immediate hypersensitivity. *Med Clin North Am* 1981; 65:941–57. [PubMed: 7026917]
7. Dombrowicz D, Flamand V, Brigman KK, Koller BH, Kinet JP. Abolition of anaphylaxis by targeted disruption of the high affinity immunoglobulin E receptor alpha chain gene. *Cell* 1993; 75:969–76. [PubMed: 8252632]
8. MacGlashan D Jr. Subthreshold desensitization of human basophils re-capitulates the loss of Syk and FcεRI expression characterized by other methods of desensitization. *Clin Exp Allergy* 2012; 42:1060–70. [PubMed: 22702505]

9. Sancho-Serra Mdel C, Simarro M, Castells M. Rapid IgE desensitization is antigen specific and impairs early and late mast cell responses targeting FcεRI internalization. *Eur J Immunol* 2011; 41:1004–13. [PubMed: 21360700]
10. MacGlashan D Jr., Lavens-Phillips S, Katsushi M. IgE-mediated desensitization in human basophils and mast cells. *Front Biosci* 1998; 3:d746–56. [PubMed: 9680509]
11. Cernadas JR, Brockow K, Romano A, Aberer W, Torres MJ, Bircher A, et al. General considerations on rapid desensitization for drug hypersensitivity - a consensus statement. *Allergy* 2010; 65:1357–66. [PubMed: 20716314]
12. Khodoun MV, Kucuk ZY, Strait RT, Krishnamurthy D, Janek K, Lewkowich I, et al. Rapid polyclonal desensitization with antibodies to IgE and FcεRIα. *J Allergy Clin Immunol* 2013; 131:1555–64. [PubMed: 23632296]
13. O'Donovan WJ, Klorfajn I. Sensitivity to penicillin; anaphylaxis and desensitisation. *Lancet* 1946; 2:444–6. [PubMed: 20998913]
14. Sullivan TJ. Antigen-specific desensitization of patients allergic to penicillin. *J Allergy Clin Immunol* 1982; 69:500–8. [PubMed: 6176609]
15. Kinet JP. The high-affinity IgE receptor (FcεRI): from physiology to pathology. *Annu Rev Immunol* 1999; 17:931–72. [PubMed: 10358778]
16. Khodoun MV, Morris SC, Angerman E, Potter C, Schuman R, Wunderlich M, et al. Rapid desensitization of humanized mice with anti-human FcεRIα monoclonal antibodies. *J Allergy Clin Immunol* 2020; 145:907–21 e3. [PubMed: 31836406]
17. Dombrowicz D, Brini AT, Flamand V, Hicks E, Snouwaert JN, Kinet JP, et al. Anaphylaxis mediated through a humanized high affinity IgE receptor. *J Immunol* 1996; 157:1645–51. [PubMed: 8759751]
18. Mathias CB, Hobson SA, Garcia-Lloret M, Lawson G, Poddighe D, Freyschmidt EJ, et al. IgE-mediated systemic anaphylaxis and impaired tolerance to food antigens in mice with enhanced IL-4 receptor signaling. *J Allergy Clin Immunol* 2011; 127:795–805 e1–6. [PubMed: 21167580]
19. Wunderlich M, Chou FS, Link KA, Mizukawa B, Perry RL, Carroll M, et al. AML xenograft efficiency is significantly improved in NOD/SCID-IL2RG mice constitutively expressing human SCF, GM-CSF and IL-3. *Leukemia* 2010; 24:1785–8. [PubMed: 20686503]
20. Wunderlich M, Brooks RA, Panchal R, Rhyasen GW, Danet-Desnoyers G, Mulloy JC. OKT3 prevents xenogeneic GVHD and allows reliable xenograft initiation from unfractionated human hematopoietic tissues. *Blood* 2014; 123:e134–44. [PubMed: 24778156]
21. Iikura M, Yamaguchi M, Hirai K, Miyamasu M, Yamada H, Nakajima T, et al. Regulation of surface FcεRI expression on human eosinophils by IL-4 and IgE. *Int Arch Allergy Immunol* 2001; 124:470–7. [PubMed: 11340330]
22. Finkelman FD, Katona IM, Urban JF Jr., Snapper CM, Ohara J, Paul WE. Suppression of in vivo polyclonal IgE responses by monoclonal antibody to the lymphokine B-cell stimulatory factor 1. *Proc Natl Acad Sci U S A* 1986; 83:9675–8. [PubMed: 3491987]
23. Gunasekaran K, Pentony M, Shen M, Garrett L, Forte C, Woodward A, et al. Enhancing antibody Fc heterodimer formation through electrostatic steering effects: applications to bispecific molecules and monovalent IgG. *J Biol Chem* 2010; 285:19637–46. [PubMed: 20400508]
24. Junghans RP, Anderson CL. The protection receptor for IgG catabolism is the β2-microglobulin-containing neonatal intestinal transport receptor. *Proc Natl Acad Sci U S A* 1996; 93:5512–6. [PubMed: 8643606]
25. Strait RT, Morris SC, Yang M, Qu XW, Finkelman FD. Pathways of anaphylaxis in the mouse. *J Allergy Clin Immunol* 2002; 109:658–68. [PubMed: 11941316]
26. Finkelman FD, Morris SC, Orekhova T, Sehy D. The in vivo cytokine capture assay (IVCCA) for measurement of in vivo cytokine production in the mouse. *Current Protocols in Immunology* 2003;6.2.8.1–6.2.8.10. [PubMed: 18432901]
27. Schuck P Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and lamm equation modeling. *Biophys J* 2000; 78:1606–19. [PubMed: 10692345]

28. LT M, SB D, RT M, PS L. Computer-aided interpretation of sedimentation data for proteins. In: HS E, RA J, HJ C, editors. *Analytical Ultracentrifugation in Biochemistry and Polymer Science*. London: Royal Society of Chemistry; 1992. p. 90–125.
29. Maurer D, Fiebiger E, Reininger B, Wolff-Winiski B, Jouvin MH, Kilgus O, et al. Expression of functional high affinity immunoglobulin E receptors (FcεRI) on monocytes of atopic individuals. *J Exp Med* 1994; 179:745–50. [PubMed: 8294882]
30. Stevens RL, Friend DS, McNeil HP, Schiller V, Ghildyal N, Austen KF. Strain-specific and tissue-specific expression of mouse mast cell secretory granule proteases. *Proc Natl Acad Sci* 1994; 91:128–32. [PubMed: 8278352]
31. Matsubara S, Li G, Takeda K, Loader JE, Pine P, Masuda ES, et al. Inhibition of spleen tyrosine kinase prevents mast cell activation and airway hyperresponsiveness. *Am J Respir Crit Care Med* 2006; 173:56–63. [PubMed: 16192454]
32. Ong CJ, Ming-Lum A, Nodwell M, Ghanipour A, Yang L, Williams DE, et al. Small molecule agonists of SHIP1 inhibit the phosphoinositide 3-kinase pathway in hematopoietic cells. *Blood* 2007.
33. Benhamou M, Ryba NJ, Kihara H, Nishikata H, Siraganian RP. Protein-tyrosine kinase p72syk in high affinity IgE receptor signaling. Identification as a component of pp72 and association with the receptor γ chain after receptor aggregation. *J Biol Chem* 1993; 268:23318–24. [PubMed: 7693687]
34. Yamashita T, Mao SY, Metzger H. Aggregation of the high-affinity IgE receptor and enhanced activity of p53/56lyn protein-tyrosine kinase. *Proc Natl Acad Sci U S A* 1994; 91:11251–5. [PubMed: 7526394]
35. Holowka D, Baird B. Antigen-mediated IgE receptor aggregation and signaling: a window on cell surface structure and dynamics. *Annu Rev Biophys Biomol Struct* 1996; 25:79–112. [PubMed: 8800465]
36. Kawakami T, Galli SJ. Regulation of mast-cell and basophil function and survival by IgE. *Nat Rev Immunol* 2002; 2:773–86. [PubMed: 12360215]
37. Goroff DK, Finkelman FD. Activation of B cells in vivo by a Fab/Fc fragment of a monoclonal anti- IgD antibody requires an interaction between the antibody fragment and a cellular IgG Fc receptor. *J Immunol* 1988; 140:2919–24. [PubMed: 2966202]
38. Strait RT, Morris SC, Smiley K, Urban JF Jr., Finkelman FD. IL-4 exacerbates anaphylaxis. *J Immunol* 2003; 170:3835–42. [PubMed: 12646651]
39. Clay CD, Strait RT, Mahler A, Khodoun MV, Finkelman FD. Anti-FcγRIIB mAb suppresses murine IgG-dependent anaphylaxis by Fc domain targeting of FcγRIII. *J Allergy Clin Immunol* 2018; 141:1373–81 e5. [PubMed: 28624610]
40. Bruhns P, Iannascoli B, England P, Mancardi DA, Fernandez N, Jorieux S, et al. Specificity and affinity of human Fcγ receptors and their polymorphic variants for human IgG subclasses. *Blood* 2009; 113:3716–25. [PubMed: 19018092]
41. Dekkers G, Bentlage AEH, Stegmann TC, Howie HL, Lissenberg-Thunnissen S, Zimring J, et al. Affinity of human IgG subclasses to mouse Fcγ receptors. *MAbs* 2017; 9:767–73. [PubMed: 28463043]
42. Lippow SM, Wittrup KD, Tidor B. Computational design of antibody-affinity improvement beyond in vivo maturation. *Nat Biotechnol* 2007; 25:1171–6. [PubMed: 17891135]
43. Zhang K, Elias M, Zhang H, Liu J, Kepley C, Bai Y, et al. Inhibition of allergic reactivity through targeting FcεRI-bound IgE with humanized low-affinity antibodies. *J Immunol* 2019; 203:2777–90. [PubMed: 31636239]
44. Incorvaia C, Mauro M, Russello M, Formigoni C, Riario-Sforza GG, Ridolo E. Omalizumab, an anti-immunoglobulin E antibody: state of the art. *Drug Des Devel Ther* 2014; 8:197–207.
45. Scharenberg AM, Kinet JP. Initial events in FcεRI signal transduction. *J Allergy Clin Immunol* 1994; 94:1142–6. [PubMed: 7798551]
46. Emlen W, Mannik M. Clearance of circulating DNA-anti-DNA immune complexes in mice. *J Exp Med* 1982; 155:1210–5. [PubMed: 7061954]
47. Grace SA, Brennan FM. Clearance and localization of immunoglobulin oligomers in mice with chronic circulating endogenous complexes. *Immunology* 1982; 47:221–6. [PubMed: 7118163]

48. Kurlander RJ, Ellison DM, Hall J. The blockade of Fc receptor-mediated clearance of immune complexes in vivo by a monoclonal antibody (2.4G2) directed against Fc receptors on murine leukocytes. *J Immunol* 1984; 133:855–62. [PubMed: 6736648]
49. Bruhns P, Jonsson F. Mouse and human FcR effector functions. *Immunol Rev* 2015; 268:25–51. [PubMed: 26497511]
50. Zhu D, Kepley CL, Zhang M, Zhang K, Saxon A. A novel human immunoglobulin Fc γ Fc ϵ bifunctional fusion protein inhibits Fc ϵ RI-mediated degranulation. *Nat Med* 2002; 8:518–21. [PubMed: 11984598]
51. Lantz CS, Yamaguchi M, Oettgen HC, Katona IM, Miyajima I, Kinet JP, et al. IgE regulates mouse basophil Fc ϵ RI expression in vivo. *J Immunol* 1997; 158:2517–21. [PubMed: 9058781]
52. Matsuda K, Piliponsky AM, Iikura M, Nakae S, Wang EW, Dutta SM, et al. Monomeric IgE enhances human mast cell chemokine production: IL-4 augments and dexamethasone suppresses the response. *J Allergy Clin Immunol* 2005; 116:1357–63. [PubMed: 16337471]
53. Morris SC, Moroldo M, Giannini EH, Orekhova T, Finkelman FD. In vivo survival of autoreactive B cells: characterization of long-lived B cells. *J Immunol* 2000; 164:3035–46. [PubMed: 10706692]
54. Eray M, Liwszyc GE, Paasinen-Sohns A, Stahls A, Kaartinen M, Andersson LC. p72syk protein tyrosine kinase: an early transducer of sIgG-triggered apoptotic signalling in human follicular lymphoma cells. *Int Immunol* 1998; 10:1573–81. [PubMed: 9796924]
55. Vilen BJ, Burke KM, Sleater M, Cambier JC. Transmodulation of BCR signaling by transduction-incompetent antigen receptors: implications for impaired signaling in anergic B cells. *J Immunol* 2002; 168:4344–51. [PubMed: 11970976]
56. Cambier JC, Getahun A. B cell activation versus anergy; the antigen receptor as a molecular switch. *Immunol Lett* 2009; 128:6–7. [PubMed: 19808046]
57. Martin TR, Galli SJ, Katona IM, Drazen J. Role of mast cells in anaphylaxis. *J Clin Invest* 1989; 83:1375–83. [PubMed: 2784802]
58. Shah M. Commentary: New perspectives on protein aggregation during biopharmaceutical development. *Int Jour Pharmaceut* 2018; 552:1–6.
59. Ohtake S, Kita Y, Arakawa T. Interactions of formulation excipients with proteins in solution and in the dried state. *Adv Drug Delivery Rev* 2011; 63:1053–73.
60. Vafa O, Gilliland GL, Brezski RJ, Strake B, Wilkinson T, Lacy ER, et al. An engineered Fc variant of an IgG eliminates all immune effector functions via structural perturbations. *Methods* 2013; 65:114–26. [PubMed: 23872058]

Clinical implication:

Mouse model studies provide evidence that a monovalent monoclonal anti-FcεRIα antibody with low Fcγ receptor binding should safely and rapidly suppress IgE-mediated disease.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. mv anti-Fc ϵ RI α mAb protects mice against dv anti-Fc ϵ RI α mAb.

A. Left panel: huFc ϵ RI α mice (3–4/group) were injected i.v. with 10 μ g of biotin-anti-IL-4 mAb and the doses of mv huIgG1 IE7 shown, followed 1 d later by i.v. injection of 100 μ g of dv huIgG1 IE7. Maximum temperature drops after each injection are shown. Fractions indicate number of mice that showed reduced activity after injection $\div$ number of mice injected. **Right panel:** IL-4 and MMCP1 responses determined in serum 4 hrs after injection of dv IE7. **B.** One d after treatment with dv IE7, naïve huFc ϵ RI α mice and mice that had originally been injected with 50 or 100 μ g of mv huIgG1 IE7 (5/group) were injected i.v. with 10 μ g of mouse IgE anti-TNP mAb and challenged i.v. the next d with 10 μ g of TNP-

OVA. Rectal temperatures during the hr after challenge are shown. **C.** huFcεRIα mice (3–4/group) were injected i.v. and i.p. with IgE anti-TNP mAb (5 µg each route), followed 1 d later by saline, dv mouse IE7 or mv huIgG1 IE7 (50 µg i.v. + 50 µg i.p.) and bled 3 hrs later for determination of serum IL-4 and MMCP1. Peritoneal lavage was performed immediately afterwards and mast peritoneal cell IgE was determined. **D.** huFcεRIα mice (4/group) were injected i.v. with 10 µg of mouse IgE anti-TNP mAb. One d later, mice were injected i.v. with saline or 25 or 50 µg of mv huIgG1 IE7. Starting 1 d after that and continuing 2x/wk for 18 d, mice that had received mv IE7 were injected i.p. with dv IE7, while mice originally injected with saline continued to receive saline. All mice were injected i.v. on d 18 with 10 µg of mouse IgE anti-TNP mAb and on d 19 with TNP-OVA (50 µg i.v.). Rectal temperatures of these mice and naïve mice on d 19 were followed for 1 hr after this last injection. MMCP1 levels on d 19 were determined for sera obtained 4 hrs after TNP-OVA injection. Statistical analysis: For all figures, ns = not significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$; all tests 2-tailed unless otherwise noted. B: Mann-Whitney U test; C: 1-way ANOVA and THSDT; D: 1-way ANOVA and THSDT.

Figure 2. mv huIgG1 IE7 safely protects against IgE-mediated anaphylaxis.

A. huFcεR1α mice (4/group) were injected i.v. with 10 μg of mouse IgE anti-TNP mAb. One d later, mice were injected i.v. with 100 μg of dv mouse IE7, 100 μg of mv huIgG1 IE7, or saline and challenged i.v. the next d with saline or 10 μg of TNP-OVA. B. huFcεR1α mice (3/group) were injected i.v. with 10 μg of IgE anti-TNP mAb plus saline, 100 μg of dv mouse IE7, or 100 μg of mv huIgG1 IE7. Cells obtained 1 d later were stained for mast cell and basophil markers and mouse IgE, as well as with mAb AER-37 and mAb 15.1. Numbers show percent inhibition as compared to cells from saline-injected mice. C. huFcεR1α mice (3–4/group) were injected on d 0 with mouse IgE anti-TNP mAb (5 μg i.v.; 5 μg i.p.). All

mice received an additional 10 µg of mouse IgE anti-TNP mAb i.p. on d 2. Groups of mice were injected with mv huIgG1 IE7 (50 µg i.p., 50 µg i.v.) on d 1, d 2, or day 3. Two hrs after the d 3 injection of mv huIgG1 IE7, all mice were injected i.v. with 20 µg of TNP-OVA. Rectal temperatures were followed for 1 hr after the last injection. Mice were bled 4 hrs after the last injection and blood levels of MMCP1 and IE7 were determined. Peritoneal cells were obtained immediately after the mice were bled and analyzed by flow cytometry for percent of mast cells, mast cell surface IgE, and mast cell FcεRI that was not bound by IgE (free FcεRIα). Statistical analysis: All panels: 1-way ANOVA and THSDT.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 3. Mast cell Syk phosphorylation is inhibited by mv huIgG1 IE7. huFc ϵ RI α mice (6/group) were treated as diagrammed; the 3 groups were treated with hu IgG control mAb and left unchallenged; with huIgG control mAb and challenged with TNP-OVA; or with non-deaggregated mv huIgG2 σ IE7 and challenged with TNP-OVA. Peritoneal cells obtained 5 min after i.p. TNP-OVA challenge were stained for mast cell-specific markers, permeabilized, and stained for Syk, phospho-Syk, phospho-SHIP, phospho-AKT, and phospho-ERK and analyzed for fluorescence intensity by flow cytometry. Statistical analysis: Kruskal-Wallis test followed by 2-tailed Mann-Whitney test.

Figure 4. Aggregation and Fc γ Rs contribute to mast cell activation by mv anti-Fc ϵ RI α mAbs. A. Egg allergy was induced in huFceRI α /F709 mice (5/group). Left panel: Mice were injected i.v. with 100 μ g of control huIgG, non-deaggregated mv huIgG1 IE7, or deaggregated mv huIgG1 IE7 and rectal temperatures were followed for 1 hr. Middle and right panel: 10 hrs after control Ab or deaggregated mv huIgG1 IE7 injection, mice were challenged o.g. with 100 mg of EW + EYP and followed for 1 hr for development of diarrhea and hypothermia. Mice were bled 3.5 hrs after Ag challenge and serum MMCP1 levels were determined. Surface IgE expression was determined for peritoneal mast cells obtained 4 hrs after Ag challenge. B. Six huFceRI α mice were sensitized with IgE anti-TNP,

IL-4C, and propranolol and treated with 2.4G2 to block Fc γ Rs. Two mice each were injected i.v. with control IgG mAb, mv huIgG1 IE7, or dv huIgG1 IE7, followed 1 d later by i.v. injection of TNP-OVA. Rectal temperatures during the hr following i.v. injections are shown. C. huFc ϵ RI α mice (4/group) were sensitized by treatment with IgE anti-TNP mAb, IL-4C, and propranolol, then injected i.v. with 100 μ g of control IgG, mv huIgG1 IE7 or mv huIgG4 IE7, and, the next d, injected i.v. with 20 μ g of TNP-OVA. Rectal temperatures were followed for 1 hr after each injection. D. huFc ϵ RI α mice (4/group) were sensitized by treatment with IgE anti-TNP mAb, IL-4C, and propranolol, then injected i.v. with 200 μ g of control IgG, non-deaggregated huIgG4 IE7, or deaggregated huIgG4 IE7 and followed for 1 hr for development of hypothermia. E. huFc ϵ RI α /F709 mice (4/group) that had been inoculated e.c. with EW + EYP, followed by repeated o.g. with the same Ags, but had never developed hypothermia or diarrhea in response to the o.g., were sensitized with IgE anti-TNP mAb plus IL-4C, then injected with the doses of deaggregated mv huIgG4 IE7 shown, followed 1 d later by i.v. challenge with TNP-OVA. Rectal temperatures were followed for 1 hr after each i.v. injection. Mast cell percentage and surface IgE staining were determined by flow cytometry for peritoneal cells obtained 1 hr after TNP-OVA injection. F. A protocol similar to Fig. 4B was followed, as shown on the time line, but with mv huIgG4 IE7 instead of mv huIgG1 IE7. 4 mice/group. Statistical analysis: A: 1-way ANOVA and THSDT (left panel), Mann-Whitney U test, or Fisher's exact test; B: 1-way ANOVA and THSDT; C: 1-way ANOVA and THSDT; D: 1-way ANOVA and THSDT or Fisher's exact test.

Figure 5. Rapid desensitization with mv huIgG4 IE7.

A. huFcεRIα/F709 mice (4/group) were sensitized with mouse IgE anti-TNP mAb and IL-4C. One day later, mice were serially injected i.p. every 2 hrs with 50, 100 and 200 µg of control huIgG or mv huIgG4 IE7. Two hrs after the last i.p. injection, mice were injected i.v. with 400 µg of control huIgG or mv huIgG4 IE7. One d later all mice were challenged i.v. with 20 µg of TNP-OVA. Mice were followed after each injection for development of hypothermia and changes in mobility (anaphylaxis score). Peritoneal cells were obtained 1 hr after Ag challenge and analyzed for mast cell surface IgE by flow cytometry. B. Egg allergy was induced in huFcεRIα/F709 mice (4–5/group). Mice were rapidly desensitized or

mock desensitized by serial i.p. injection every 2 hrs with 25, 50, 100, 200, and 400 μ g of control huIgG4 or mv huIgG4 IE7. Mice were challenged o.g. with 100 mg of EW + EYP and followed for 1 hr for development of diarrhea and hypothermia. Peritoneal cells and sera were obtained 2 hrs after the o.g. challenge and analyzed for mast cell surface IgE and MMCP1 levels, respectively. C. huFc ϵ RI α /F709 mice (5/group) were sensitized with mouse IgE anti-TNP mAb and IL-4C, injected with GK1.5 anti-CD4 mAb, then mock desensitized with control huIgG, rapidly desensitized for 1 d with mv huIgG4 IE7, or rapidly desensitized with mv huIgG4 IE7, followed by injection of 400 μ g of mv huIgG4 IE7 every other d for 7 d, as shown in the flow diagram. All mice were challenged i.v. with TNP-OVA one d after the last dose of control IgG or mv huIgG4 IE7 and followed for 1 hr for hypothermia, changes in mobility (anaphylaxis score) and death. Peritoneal cells obtained 1 hr after Ag challenge with analyzed for percent of mast cells and mast cell surface IgE. Statistical analysis: A: 1-way ANOVA and THSDT; B: Mann-Whitney U test; C: 1-way ANOVA and THSDT.

Figure 6. In vivo desensitization of human mast cells with mv huIgG4 IE7.

A. reNSGS mice (4–5/group) were sensitized with huIgE anti-TNP mAb and the next d injected i.p. with 100 µg of control huIgG or deaggregated mv huIgG4 IE7, followed 3 hrs later by i.v. injection of 100 µg of the same mAb. Mice were challenged i.v. 1 d later with 20 µg of TNP-OVA. Mice were followed for 1 hr for hypothermia and death after the i.v. injection. Spleens and peritoneal cells were obtained 1 hr after Ag challenge or immediately after death and analyzed for mast cell surface IgE. B. reNSGS mice (6/group) were sensitized with huIgE anti-TNP mAb on d 0 and 3, rapidly desensitized or mock desensitized i.p. with five doubling doses of deaggregated mv huIgG4 IE7 or huIgG control every 2 hrs, starting with 25 µg of mAb; injected i.p. on d 3 with 400 µg of deaggregated mv huIgG4 IE7 or huIgG control, and challenged i.v. on d 4 with 20 µg of TNP-OVA. Mice were followed for 1 hr for hypothermia and death after the i.v. injection. Statistical analysis: 1-way ANOVA and THSDT or Fisher's exact test. B: Mann-Whitney U test.

Figure 7. The presence of anti-huIgG4 Abs does not increase the adverse effects of injecting mv hulG4 IE7.

A. huFcεRIα/F709 mice (3–4/group) were immunized twice i.p. with dv huIgG4 IE7 + alum to induce anti-huIgG4 Ab (upper left panel) or with mouse IgG + alum (control). Mice with high anti-huIgG4 Ab titers, control mice, and naïve mice were sensitized with IgE anti-TNP mAb. One d later all except the naïve mice were desensitized by serial i.p. injection every 2 hrs with 25, 50, 100, 200, and 400 μg of mv huIgG4 IE7. Mice were followed for 1 hr after each injection for development of hypothermia. All mice were challenged i.v. with TNP-OVA 1 d after desensitization and followed for 1 hr for development of hypothermia and

changes in mobility. Serum was obtained 1 hr after Ag challenge and analyzed for MMCP1 level. B. huFcεRIα/F709 mice (4/group) were immunized twice s.c. with dv huIgG4 IE7 + alum or mouse IgG + alum. All immunized mice were then sensitized with mouse IgE anti-TNP mAb. One d later, one group each of huIgG4-immunized and mouse IgG-immunized mice was rapidly desensitized i.p. with mv huIgG4 IE7, while one group of mouse IgG-immunized mice received no further treatment. Four d later, mice that had been rapidly desensitized received an additional 400 µg of mv huIgG4 IE7 (half i.p.; half i.v.). All mice were challenged i.v. with TNP-OVA one d later. Mice were followed for the development of hypothermia and death after each mAb and Ag administration. Peritoneal cells obtained 1 hr after Ag challenge were analyzed for percent mast cells and mast cell surface IgE. Statistical analysis: A: Mann-Whitney U test (left panel) or 1-way ANOVA and THSDT; B. 1-way ANOVA and THSDT or Fisher's exact test.

Figure 8. mv anti-FcεRIα mAb on-rate determines in vivo potency.

huFcεRIα/F709 mice (4/group) were treated with IL-4C and sensitized with 10 µg of IgE anti-TNP mAb. One d later, mice were injected i.v. with 200 µg of huIgG control mAb or the doses of mv huIgG4 IE7 or mv huIgG4 AER-37 shown and followed for development of hypothermia (A, upper panels) and changes in mobility (B, left panel). All mice were challenged i.v. with 20 µg of TNP-OVA and followed for death and hypothermia (A, lower panels) and changes in mobility (B, right panel). Peritoneal cells obtained 1 hr after challenge were evaluated for percent of mast cells and mast cell IgE fluorescence (C). D. huFcεRIα mice (4/group) were injected s.c. with 400 µg of mv huIgG4 IE7 or 100 µg of mv

huIgG4 AER-37. Different groups of mice were bled 2, 24, 48, 72, or 120 hr later. mv mAb levels were determined by ELISA. Percent decreases in serum mAb levels from 24 to 120 hrs did not differ significantly for mv huIgG4 IE7 and mv huIgG4 AER-37 (2-tailed Mann-Whitney U test). Statistical analysis: A: Kruskal-Wallis test and Mann-Whitney U test or Fisher's exact test; B. Mann-Whitney U test with compensation for multiple comparisons.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript