Rethinking Geschwind Syndrome Beyond Temporal Lobe Epilepsy

Burçin ÇOLAK¹ , Rıfat Serav İLHAN¹ , Berker DUMAN^{2,3}

¹Department of Psychiatry, Faculty of Medicine, Ankara University, Ankara, Turkey

²Division of Consultation-Liaison Psychiatry, Department of Psychiatry, Faculty of Medicine, Ankara University, Ankara, Turkey

³Neuroscience and Neurotechnology Center of Excellence (NÖROM), Ankara, Turkey

Geschwind Syndrome (GS) is a controversial clinical diagnosis defined as a cluster of inter-ictal behavioral manifestations as hypergraphia, hyperreligiosity, hyposexuality, mental rigidity, verbal and non-verbal viscosity (1). Behavioral manifestations of this syndrome are traditionally thought to be stemmed from temporal lobe epileptic seizures (TLE) via hyper-reactivity in the limbic networks (2). According to N. Geschwind, limbic damage that occurred during seizures cause the syndrome (3). The syndrome also has been known as temporolimbic personality; reflecting behavioral manifestations stemmed from recurrent seizures (4). However, the causality and the association between TLE and GS is still an enthusiastic old debate going on (5). In most cases, the behavioral presentation is interictal without a specific relationship to individual seizures. Besides this, GS-like manifestations also have been reported in other neuropsychiatric conditions (6, 7). For instance, GS has been described in patients with the right temporal variant of frontotemporal lobar degeneration (FTLD), right temporal stroke, right hippocampal atrophy, and various neurodegenerative diseases (8-12). There are also salient overlaps with the phenomenological manifestations of GS and neurodevelopmental disorders such as schizophrenia, schizoaffective disorder, and bipolar disorder without TLE or any neurological disease (13-16).

Electroencephalography (EEG) anomalies with or without any epileptic seizures are also another important aspect of neurodevelopmental disorders. For instance, 60% of patients with autism spectrum disorders (ASD) show abnormal epileptiform activities without any seizures that may contribute to the symptom presentation of ASD (17, 18). EEG and event-related potential (ERP) studies also indicate epileptiform activities without manifest seizures in schizophrenia and bipolar disorders (19). These changes may also be associated with emotional, cognitive, and behavioral aspects of these neuropsychiatric conditions. The rationale for the treatment of these neuropsychiatric conditions with anti-epileptics indicates the importance of subthreshold epileptiform abnormalities in neuropsychiatric conditions. As indicated above, symptoms like hyper-religiosity, hypergraphia, hyposexuality or interpersonal viscosity may indicate shared pathophysiology associated with right temporolimbic structures such as atrophy, sclerosis, or neurodevelopmental deviations in neurosynaptic formations.

Previous research on GS and TLE are still inconclusive probably due to assuming a probable false causality between these two clinical phenomena (20). Perhaps the main problem is in the presentation of the condition. Rather than psychiatric symptoms, mainly the first hospital admission is due to the TLE seizures which attract the clinicians' attention. Hence, behavioral manifestations of GS could easily be misattributed to the TLE. On the other hand, most of the TLE patients are free of these behavioral manifestations which could interfere with previous research findings.

We suggest that GS should be redefined as a neurodevelopmental condition that probably starts in childhood and is associated with temporo-limbic networks with or without any epileptic seizures. Indeed, we hypothesized that although the dysfunctional temporo-limbic networks were the assumed substrate for GS, TLE neither excludes nor supports the condition.

Also, GS may have significant impacts on social groups. Behavioral manifestations such as hypergraphia, hyperreligiosity, hyperviscosity in interpersonal relational patterns may create important and novel memes that motivate the populous in terms of group behavior. Some individuals with GS are also prone to mystic experiences that in turn increases the influence on the group (21). Additionally, GS has been previously associated with various political and ideological charismatic leaders (22, 23). GS could be viewed as a deviation from typical neurodevelopmental pathways with manifest or latent EEG abnormalities to striking neuropsychiatric symptoms such as auditory hallucinations, or charismatic personality characteristics misinterpreted by the groups as a sign of the sacred.

Cite this article as: Çolak B, İlhan RS, Duman B. Rethinking Geschwind Syndrome Beyond Temporal Lobe Epilepsy. Arch Neuropsychiatry 2021;58:169-170.

To our opinion, GS should be investigated as a unique neurodevelopmental condition independent from TLE with its potential phylogenetic outcomes. We hypothesized that the core feature of GS is its behavioral manifestations and TLE and/or EEG abnormalities should only be included as a specifier. Associations between patients with delusional disorders, paranoid, schizoid, and schizotypal personality disorders, and autism spectrum disorders and GS should also be investigated independently of epileptic seizures (24). This can lead researchers and clinicians to find and apply efficient pharmacological and psychotherapeutic interventions.

REFERENCES

- Waxman SG, Geschwind N. The interictal behavior syndrome of temporal lobe epilepsy. *Arch Gen Psych* 1975;32:1580–1586. [\[Crossref\]](#)
- Bear DM, Fedio P. Quantitative analysis of interictal behavior in temporal lobe epilepsy. *Arch Neurol* 1977;34:454–467. [\[Crossref\]](#)
- Waxman SG, Geschwind N. Hypergraphia in temporal lobe epilepsy. *Epilepsy Behav* 2005;6:282–291. [\[Crossref\]](#)
- Aycicegi-Dinn A, Dinn WM, Caldwell-Harris CL. The temporolimbic personality: A cross-national study. *Eur J Psychiatry* 2008;22:211–224. [\[Crossref\]](#)
- Benson DF. The Geschwind syndrome. *Adv Neurol* 1991;55:411–421. <https://pubmed.ncbi.nlm.nih.gov/2003418/>
- Mungas D. Interictal behavior abnormality in temporal lobe epilepsy: a specific syndrome or nonspecific psychopathology?. *Arch Gen Psychiatry* 1982;39:108–111. [\[Crossref\]](#)
- Dodrill CB, Batzel LW. Interictal behavioral features of patients with epilepsy. *Epilepsia* 1986;27:S64–S76. [\[Crossref\]](#)
- Veronelli L, Makaretz SJ, Quimby M, Dickerson BC, Collins JA. Geschwind Syndrome in frontotemporal lobar degeneration: neuroanatomical and neuropsychological features over 9 years. *Cortex* 2017;94:27–38. [\[Crossref\]](#)
- Hoffmann M. Isolated right temporal lobe stroke patients present with Geschwind Gastaut syndrome, frontal network syndrome and delusional misidentification syndromes. *Behav Neurol* 2008;20:83–89. [\[Crossref\]](#)
- Wuerfel J, Krishnamoorthy ES, Brown RJ, Lemieux L, Koepp M, Van Elst LT, Trimble MR. Religiosity is associated with hippocampal but not amygdala volumes in patients with refractory epilepsy. *J Neurol Neurosurg Psychiatry* 2004;75:640–642. [\[Crossref\]](#)
- van Elst LT, Krishnamoorthy ES, Bäumer D, Selai C, Von Gunten A, Gene-Cos N, Ebert D, Trimble MR. Psychopathological profile in patients with severe bilateral hippocampal atrophy and temporal lobe epilepsy: evidence in support of the Geschwind syndrome?. *Epilepsy Behav* 2003;4:291–297. [\[Crossref\]](#)
- Postiglione A, Milan G, Pappata S, De Falco C, Lamenza F, Schiattarella V, Gallotta G, Sorrentino P, Striano S. Fronto-temporal dementia presenting as Geschwind's syndrome. *Neurocase* 2008;14:264–270. [\[Crossref\]](#)
- Marques JG, Teixeira J, Carnot MJ. Geschwind's syndrome in a patient with schizophrenia. *Psychiatry Res* 2015;228:976. [\[Crossref\]](#)
- Marques JG. A neuropsychiatric case report: Klüver-Bucy and Gastaut-Geschwind syndromes in one patient with schizoaffective disorder presenting with brain 99-mTc HMPAO single-photon emission computed tomography showing bilateral discrete decrease of temporal lobes perfusion. *CNS Spectrums* 2018;23:358–360. [\[Crossref\]](#)
- O'Connell K, Keaveney J, Paul R. A novel study of comorbidity between schizoaffective disorder and Geschwind syndrome. *Case Rep Psychiatry* 2013;486064. [\[Crossref\]](#)
- Mudgal V, Paliwal A, Pal VS, Prajapati D. Gastaut-Geschwind Syndrome in a patient of bipolar disorder: A case report. *Ind J Psychol Med* 2021;43:270–272. [\[Crossref\]](#)
- Boutros NN, Lajiness-O'Neill R, Zillgitt A, Richard AE, Bowyer SM. EEG changes associated with autistic spectrum disorders. *Neuropsychiatr Electrophysiol* 2015;1:1–20. [\[Crossref\]](#)
- Gilby KL, O'Brien TJ. Epilepsy, autism, and neurodevelopment: Kindling a shared vulnerability?. *Epilepsy Behav* 2013;26:370–374. [\[Crossref\]](#)
- Degabriele R, Lagopoulos J. A review of EEG and ERP studies in bipolar disorder. *Acta Neuropsychiatrica* 2009;21:58–66. [\[Crossref\]](#)
- Bear DM. Behavioural changes in temporal lobe epilepsy: conflict, confusion, challenge. In: Trimble MR, Bolwig TG, editors. *Aspects of Epilepsy and Psychiatry*. Chichester, England: Wiley; 1986. p.19–30.
- Devinsky O, Lai G. Spirituality and religion in epilepsy. *Epilepsy Behav* 2008;12:636–643. [\[Crossref\]](#)
- Muhammed L. A retrospective diagnosis of epilepsy in three historical figures: St Paul, Joan of Arc and Socrates. *J Med Biography* 2013;21:208–211. [\[Crossref\]](#)
- Saver JL, Rabin J. The neural substrates of religious experience. *J Neuropsychiatr Clin Neurosci* 1997;9:498–510. [\[Crossref\]](#)
- Trimble MR, Mendez MF, Cummings JL. Neuropsychiatric symptoms from the temporolimbic lobes. *J Neuropsychiatr Clin Neurosci* 1997;9:429–438. [\[Crossref\]](#)