Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Adaptation, Implementation, and Impact of Cooking Matters® for Diabetes on Dietary Outcomes, Food Security, and Quality of Life

Author(s): J. Shrodes¹, C. Spees², J. Radabaugh³, A. Braun², J. Joseph¹, A. Williams¹; ¹The Ohio State University Wexner Medical Center, ²The Ohio State University, ³The Ohio State University Medical Dietetics

Learning Outcome: Describe the impact of Cooking Matters® for Diabetes (CMFD) on dietary outcomes, food security, and health-related quality of life (HRQOL) compared to standard of care (SOC) in people with diabetes

Background: Cooking Matters®, an evidence-based program, facilitates behavior change yet has not been evaluated for incorporation into diabetes self-management education and support (DSMES).

Methods: Forty-eight adults with diabetes (type 1 or 2) were randomized to CMFD intervention (n=26) or waitlist control groups receiving SOC (n=22). The 6-week intervention included weekly (2-hour) group diabetes education with interactive cooking demonstrations and free groceries. Validated measurements of dietary patterns (Diet History Questionnaire III), diet quality (Healthy Eating Index (HEI) 2015), diet-related self-care (Summary of Diabetes Self-Care Activities general and specific diet scores), food insecurity (USDA module), and HRQOL (SF-12) were evaluated at baseline, post-intervention, and 3 months post-intervention. Differences were compared via linear mixed-effects models.

Results: Compared to controls, CMFD participants had higher mental HRQOL scores post-intervention (+6.707, p=0.025), higher general diet scores post-intervention (+1.51, p=0.015) and at 3 months (+1.23, p=0.05), and higher HEI vegetable component scores at 3 months (+0.917, p=0.023). Longitudinally, general (+1.061, p=0.001) and specific (+0.810, p=0.002) diet scores improved post-intervention, while food insecurity decreased at 3 months (-0.637, p=0.021) among all CMFD participants after going through intervention (n=48). Decreases in daily energy (-348kcal, p=0.013), carbohydrate (-51g, p=0.005), and total sugar (-23.802, p=0.02) intakes were also observed post-intervention.

Conclusion: This study supports the incorporation of evidence-based cooking interventions into DSMES as an effective tool in promoting positive behavior change and diet-related outcomes.

Funding source: Diabetes Care and Education Dietetic Practice Group Karen Goldstein Memorial Grant from the Academy of Nutrition and Dietetics Foundation

An Analysis of MyPlate FoodGroup Intake in Children with Autism Spectrum Disorder

Author(s): N. Withrow, E. Hsueh; University of Northern Colorado

Learning Outcome: Understand the differences in food variety intake in children with ASD compared to typically developing children.

Background: Autism Spectrum Disorder (ASD) is a complex developmental disorder impacting 1 in 59 children in the United States and is characterized by social, emotional, communication, and behavioral challenges. Children with ASD commonly exhibit food selectivity due to sensory processing difficulties, aberrant mealtime behaviors, and motor impairment. Previous studies have shown that food selectivity can impact healthy growth and development with implications for nutrient deficiencies.

Method: The objective of this study was to understand the differences of intake and variety of MyPlate food groups between children with an ASD (n=10) and typically developing children (TD)(n=27). A secondary analysis was conducted using the data of a pilot study to develop a comprehensive eating screening inventory named the Sensory Processing, Aberrant Mealtime Behaviors, Motor, Inventory for Eating (SAMIE). Three-day diet records were analyzed for average daily intake of grains, fruits, vegetable, dairy, and protein using a diet analysis software, Nutritionist Pro. Food variety was assessed by categorizing unique food items in each food group.

Results: No differences were found in calories and average daily intake of food groups between ASD and TD, however ASD showed significantly less variety in vegetables (p=0.04) and protein (p=0.03). Mean differences showed that ASD had less variety in all food groups compared to TD.

Conclusions: This study supports previous findings that children with ASD exhibit more food selectivity and have limited food repertoire compared to typically developing children.

Funding source: University of northern Colorado research, faculty grant

Content Analysis of Type 1 Diabetes Nutrition Images on Instagram

Author(s): D. Kelleher, L. Weatherspoon; Michigan State University

Learning Outcome: Describe the main themes present in Type 1 Diabetes nutrition related Instagram posts.

Background: Social media platforms are used daily by over 90% of young adults aged 18-29 years with Instagram being highly popular with this age group. Users are active participants in the conversation sharing health information including those with chronic health conditions such as type 1 diabetes. Little is known about the specific nutrition information being shared on social media including who is posting and if the information is consistent with current diabetes recommendations. This research sets to understand the major nutrition themes being shared on Instagram for two popular type 1 diabetes hashtags: #type1diabetes and #diabadass.

Methods: A sample of 300 nutrition related Instagram images marked with #type1diabetes and #diabadass from April-July 2020 were coded for Instagram user types, nutrition themes, meal occasions, diet and disordered eating related captions.

Results: A total of 298 Instagram images were analyzed posted by 256 unique Instagram users. The Instagram users were mostly females (80%), persons with diabetes (87.5%). Other main user types included personal businesses (7.7%) and health care providers (7%). The main nutrition theme for posts included meals (52.6%), desserts (6%), and normalizing food (5.7%). Diets were promoted in nearly half of all posts (48.7%) with ketogenic diet being the most popular diet (48.2%). Body image was a topic of post or caption in 8.7% of all posts.

Conclusion: Nutrition information is being shared on social media by persons with diabetes. This form of support involves diet specific information including the ketogenic diet which is not in line with current MNT recommendations.

Funding source: None

COVID-19 Impact on Stress and Self-Efficacy among RDNs and CDMs Practicing in Long-Term Care, Post-Acute Care, and Assisted Living Settings

Author(s): M. Johnson, J. Barosso; East Tennessee State University

Learning Outcome: Describe stressors experienced by RDNs and CDMs related to the COVID-19 pandemic during practice in long-term care and assisted living facilities.

While literature exploring the effects of COVID-19 on medical practitioners has emerged over time, evidence regarding the impact on nutrition professionals remains limited. This study aimed to identify pandemic-related stressors among RDNs and CDMs working in long-term care, rehab, and assisted living settings. With ethics review board approval, a 47-question, Qualtrics survey was distributed to 86 RDNs and 14 CDMs across 20 states. 78.7% (n=37) of respondents were RDNs; 21.2% (n=10) CDMs. The majority described working in long-term care with post-acute rehabilitation (51.6%), long-term care (28%), and assisted living (17%), serving 4616 residents. Mean facility census was 94.2. Despite census declines, responsibilities did not decrease. RDNs completed some (51.4%) or all (37.8%) work remotely. Lack of direct access to residents and families, staffing shortages, and infection control-related barriers (weights, feeding practices, providing nutrition interventions) decreased self-efficacy, due to perceived inability to provide high quality care. Further, 78.7% of respondents indicated they experienced increased personal stress during the pandemic. Emerging themes were resident-related, reporting sadness about resident mortality, loneliness, and decreased psychological health, and personal, including loss of childcare, depression, isolation in their jobs and personal lives, and fear of contracting the virus, and infecting family members. On a positive note, participants felt supported by employers, with access to training and consistent policy updates. They were quick to identify the strengths of foodservice employees and their hard work, and enjoyed the opportunity to increase the perceived value of our professions, with creative solutions they plan to carry forward after the pandemic.

Funding source: None