Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

CORRESPONDENCE

Perspective in Remote
Cardiopulmonary Bypass Control

TO THE EDITOR: Technology has made great progress today—it would be possible to replace the Formula 1 race car driver with remote control. We have idealized the relationship that the perfusionist has with extracorporeal circulation as the one the pilot has with the formula 1 car. The goal of cardiopulmonary bypass (CPB) is to guarantee accurate metabolic support during cardiac surgery procedures. The macro areas that have been identified related to CPB are safety, practice, understanding, interpretation, and management during extracorporeal circulation.¹ In cardiac surgery practice, telesurgery is a surgical system that utilizes wireless networking and robotic technology to connect surgeons and patients who are distantly located from one another. The system overcomes today's shortage of surgeons, geographical inaccessibility of immediate and high-quality surgical care, significant financial burden, potential complications, and long-distance travel.²

We read with high interest the article by Hermesen and colleagues,³ “Multimodal cardiopulmonary bypass skills assessment within a high-fidelity simulation environment.” They present a high-fidelity simulator that uses a perfused porcine heart, cannulae, and tubing that has been demonstrated to be a useful training adjunct. They hypothesized that multimodal assessment of CPB skills within this high-fidelity simulated environment could discern expert from trainee performance.³

In this context, we present for support the learning curve, the remote control of CPB on cardiovascular perfusion training and practice. The remote CPB control could be the future challenge for the support of the young perfusionist to perform CPB with the senior perfusionist, even though they are not physically in the same location during complex CPB management. It is a form of telepresence. A robot perfusion system generally consists of one or more arms (controlled by the perfusionist), a master controller (console), and a sensory system giving feedback to the user. Remote perfusion combines elements of robotics, cutting-edge communication technology such as high-speed data connections, and elements of management information systems. In perspective, remote CPB control is essentially advanced telecommuting for perfusionists, where the physical distance between the senior perfusionist and young operators is more and less relevant. That would allow the experience of specialized perfusionists to support their colleagues worldwide in complex situations, without the need for the senior perfusionist to travel to the hospital.

Ignazio Condello, PhD

Department of Cardiac Surgery
Anthea Hospital
GVM Care and Research
Via Camillo Rosalba 35/37
Bari 70124, Italy
email: ignicondello@hotmail.it

Giovanni Battista Camarda, MS

Department of Cardiac Surgery
Maria Eleonora Hospital
GVM Care and Research
Palermo, Italy

REFERENCES

1. Condello I. The flow of time in cardiopulmonary bypass. *Artif Organs*. 2020;44:435-436.
2. Choi PJ, Oskoui RJ, Tubbs RS. Telesurgery: past, present, and future. *Cureus*. 2018;10:e2716.
3. Hermesen JL, Mohamadipannah H, Yang S, et al. Multimodal cardiopulmonary bypass skills assessment within a high-fidelity simulation environment. *Ann Thorac Surg*. 2021;112:652-660.

Overcoming Reluctance to Use
Available Tools: Silver Linings of
COVID

REPLY TO THE EDITOR: We read with appreciation the comments by Condello and Camarda¹ regarding our recent publication.² Although I would not necessarily have drawn a line between our work and theirs, this “cross-pollination” of ideas highlights one of the unforeseen benefits intrinsic to academic pursuits.

Their description of “remote control,” or as I read it, “remote oversight and mentorship” of young or less experienced perfusionists working in underserved areas is worthy of consideration. That seems most akin to the system of electronic intensive care units (eICUs) where a remotely located intensivist has access to the complete complement of monitoring and laboratory data, the ability to examine the patient through a camera, and the ability to communicate, direct, and coordinate care with the bedside nurses and therapists.³ There is no reason to think that a similar type of arrangement could not be utilized to augment the care of a patient remote from the senior perfusionist.

A year ago I may have been more likely to focus on the hurdles inherent in such a system and emphasized the human factors inherent to our work and teams. The importance of these facts has not changed but the lens through which I view them has. The COVID-19 pandemic is a scourge, but has shown us, in real time, our abilities to adapt how we communicate and connect with our team members in functional,

meaningful ways. My new normal for the past 9 months is video-conference rounding on our inpatient service, including the sickest patients in the intensive care unit. This adaptation born of necessity has worked, and I expect will persist after we have all been vaccinated!

Kudos to Condello and Camarda¹ to expand the reaches of telemedicine to include the very technical and hands-on arena of perfusion. If my grandmother's Zoom funeral can bring me to the same tears I would have shed in person, there is little reason to think one perfusionist cannot help another take care of a patient from a distance.

Joshua L. Hermesen, MD

Section of Congenital Heart Surgery
Division of Cardiothoracic Surgery
American Family Children's Hospital
University of Wisconsin-Madison
600 Highland Ave
CSC, H4/352
Madison, WI 53792
email: hermsen@surgery.wisc.edu

REFERENCES

1. Condello I, Camarda GB. Perspective in remote cardiopulmonary bypass control. *Ann Thorac Surg.* 2021;112:1378.
2. Hermesen JL, Mohamadipanah H, Yang S, et al. Multimodal cardiopulmonary bypass skills assessment within a high-fidelity simulation environment. *Ann Thorac Surg.* 2021;112:652-660.
3. Gupta S, Dewam S, Kaushal A, et al. eICU reduced mortality in STEMI patients in resource-limited areas. *Glob Heart.* 2014;9:425-427.

A Different Perspective on Hospital Readmissions

TO THE EDITOR: I read with great interest the article in *The Annals of Thoracic Surgery* by Hirji and colleagues¹ on the clinical impact on nonindex readmissions after cardiac surgery. I do, however, question the assumptions they offered in their introduction.

To start, their opening statement, "Hospital readmissions . . . have been shown to compromise quality of care," is misleading and is not confirmed in the references they offer in support of the statement. In fact, the reverse is true: The quality of care in the initial hospitalization may contribute in part to the need for a readmission (although the degree to which that is true is open to debate). Most readmissions actually improve the quality of care by addressing and correcting chronic conditions that may be exacerbated after surgery in our older adult population with multiple comorbidities. (When presenting a readmission paper at the 2014 meeting of the American Association for Thoracic Surgery, a commenter ask me rhetorically why readmissions were considered "a priori bad" when indeed they are "good" for our patients.²)

The final statement of that first paragraph (extolling the promising results of the Hospital Readmissions Reduction Program) also needs to be viewed in the entirety of the outcomes of this regulatory program to date. Indeed, although readmission rates have declined in both targeted and nontargeted conditions, and coordinated care now has an additional incentive, negative ramifications are similarly well documented. These include disproportionate penalties to hospitals serving the most indigent populations (with higher readmission rates linked to lower socioeconomic status, as Hirji and colleagues¹ pointed out) and the danger of delay or avoiding necessary readmissions.^{3,4} It has also encouraged the practice of classifying a return to the hospital under "Observation Status" rather than a "Hospital (Re) Admission" to avoid financial penalty (unfortunately shifting that penalty to the patients, whose financial responsibility increases with the former vs the latter).⁵

I offer these points not to detract from an otherwise excellent paper, but rather to deemphasize the stigma and underlying desire to avoid readmissions at all costs, and instead to underline the call made by Hirji and colleagues¹ for more integrated systems of care to ensure that those patients who do require readmission receive the best possible care when they return.

Robert Lancey, MD, MBA

Cardiothoracic Surgery Services
Sentara Rockingham Memorial Hospital
2020 Health Campus Dr
Harrisonburg, VA 22801
email: ralancey@sentara.com

REFERENCES

1. Hirji SA, Percy ED, Zogg CK, et al. Thirty-day nonindex readmissions and clinical outcomes after cardiac surgery. *Ann Thorac Surg.* 2020;110:484-491.
2. Lancey R, Kurlansky P, Argenziano M, et al. Uniform standards do not apply to readmission following coronary artery bypass surgery: a multi-institutional study. *J Thorac Cardiovasc Surg.* 2015;149:850-857.
3. McIlvennan CK, Eapen ZJ, Allen LA. Hospital Readmissions Reduction Program. *Circulation.* 2015;131:1796-1803.
4. Fonarow GC, Konstam MA, Yancy CW. The Hospital Readmissions Reduction Program is associated with fewer readmissions, more deaths: time to reconsider. *J Am Coll Cardiol.* 2017;70:1931-1934.
5. Zuckerman RB, Sheingold SH, Orav EJ, et al. Readmissions, observation, and the Hospital Readmissions Reduction Program. *N Engl J Med.* 2016;374:1543-1551.

Integrated Cardiac Surgery Systems of Care and Care Fragmentation: A Continuum Beyond Discharge

REPLY TO THE EDITOR: We read with great interest the letter to the editor by Dr Lancey, titled "A different perspective on hospital readmissions,"¹ in response to our recent publication² in *The Annals of Thoracic Surgery*