Published in final edited form as:

Infect Control Hosp Epidemiol. 2020 May ; 41(5): 571–578. doi:10.1017/ice.2019.373.

Considerable variability in antibiotic use among United States children's hospitals in 2017–2018

Hannah G. Griffith¹, Keerti Dantuluri, MD², Cary Thurm, PhD³, Derek J. Williams, MD, MPH², Ritu Banerjee, MD, PhD², Leigh M. Howard, MD, MPH^{*,2}, Carlos G. Grijalva, MD, MPH^{*,1}

¹Department of Health Policy, Vanderbilt University Medical Center, Nashville, Tennessee, USA

²Department of Pediatrics, Vanderbilt University Medical Center, Nashville, Tennessee, USA

³Children's Hospital Association, Overland Park, Kansas City, Kansas, USA

Abstract

Objective—To characterize the prevalence of and seasonal and regional variation in inpatient antibiotic use among hospitalized US children in 2017–2018.

Design—This is a cross-sectional examination of hospitalized children. Assessments were conducted on a single day in spring (May 3, 2017), summer (August 2, 2017), fall (October 25, 2017), and winter (January 31, 2018). The main outcome of interest was receipt of an antibiotic on study day.

Setting—51 freestanding US children's hospitals that participate in the Pediatric Health Information System (PHIS)

Patients—This study included all patients less than 18 years old who were admitted to a participating PHIS hospital, excluding patients who were admitted solely for research purposes.

Results—Of 52,769 total hospitalized children, 19,174 (36.3%) received antibiotics on the study day, and 6,575 (12.5%) received broad-spectrum antibiotics. The overall prevalence of antibiotic use varied across hospitals from 22.3% to 51.9%. Antibiotic use prevalence was 29.2% among medical patients and 47.7% among surgical patients. Although there was no significant seasonal variation in antibiotic use prevalence, regional prevalence varied, ranging from 32.7% in the Midwest to 40.2% in the West ($p < 0.001$). Among units, pediatric intensive care unit patients had the highest prevalence of both overall and broad-spectrum antibiotic use at 58.3% and 26.6%, respectively ($p < 0.001$).

Conclusions—On any given day in a national network of children's hospitals, over a third of hospitalized children received an antibiotic, and one in eight received a broad-spectrum antibiotic. Variation across hospitals, setting and regions identifies potential opportunities for enhanced antibiotic stewardship activities.

Address correspondence to: Hannah G. Griffith, Department of Health Policy, Vanderbilt University Medical Center, 1500 21st Avenue S., Suite 2600, Nashville, TN 37212, hannah.griffith.1@vumc.org, (615) 343-5499.

*These authors contributed equally to this work.

INTRODUCTION

Appropriate antibiotic therapy is the cornerstone of management of patients with bacterial infections, and antibiotics are the most commonly prescribed class of medications among children [1]. However, antibiotics are widely overused, and recent studies have demonstrated that a substantial proportion of antibiotic prescriptions may be inappropriate, defined as use without an appropriate indication or prescription of the incorrect drug, dose, or treatment duration [2–5].

Antimicrobial resistance is one of the most serious threats to public health worldwide. Antibiotic use is an important driver of the development and spread of antibiotic-resistant bacteria. In addition, antibiotic use is associated with adverse events including disruption of healthy enteric bacterial flora, *Clostridioides difficile* infections, hypersensitivity reactions, nephrotoxicity, myelosuppression, hepatitis, and increased healthcare costs [6–8]. Inappropriate antibiotic use needlessly exposes patients to these potential risks. The Centers for Disease Control and Prevention (CDC) has recommended that all acute care hospitals develop and implement Antibiotic Stewardship Programs (ASP) to improve appropriate use of antibiotics [6]. Implementation of antibiotic stewardship programs at pediatric hospitals has been associated with an average decline in monthly antibiotic use of almost 6% between 2007 and 2012 [9]. To enhance appropriate use of antibiotics and to minimize the potential for preventable harm, it is necessary first to understand the patterns of antibiotic use.

A cross-sectional study of hospitalized children showed that more than half of them received at least one dose of an antibiotic during their hospital stay in 2010 [10]. Another cross-sectional assessment of antibiotic use in US hospitals encompassing both children and adults in 2011 showed that up to half of all hospitalized patients received at least one antibiotic during their stay [11]. However, children represented only a small proportion of that study population, and the seasonal variability in antibiotic use was not assessed. Few other studies have characterized the epidemiology of antibiotic use in the inpatient setting in children. In the last decade, most acute care hospitals have implemented antibiotic stewardship programs to reduce overuse and misuse of antibiotics, and provider awareness of the need for judicious antibiotic use has increased [9]. We examined inpatient antibiotic use prevalence and characterized the seasonal and regional variation across all four seasons in one recent year among hospitalized US children.

PATIENTS AND METHODS

Overview

We conducted a cross-sectional study of inpatient antibiotic use among children admitted to hospitals that participate in the Children's Hospital Association's Pediatric Health Information System (PHIS). Antibiotic use prevalence was measured at four specific points in time. We compared prevalence of antibiotic use among participating hospitals and by services (medical versus surgical) and units where children were hospitalized. We also examined the potential association between antibiotic use and clinical complexity at individual hospitals.

Data Source

PHIS is a clinical and resource utilization database for over 50 free-standing US children's hospitals. PHIS data contain detailed information for each patient hospitalization, including demographics, diagnoses, medications, procedures, and laboratory tests. Data are maintained by the Children's Hospital Association, participating hospitals, and Truven Health Analytics. Children are assigned unique study identifiers to allow for linking across encounters within a hospital while preserving the privacy of their health information [12]. This study of existing, deidentified patient data was reviewed and considered exempt from further review by the Vanderbilt University Medical Center Institutional Review Board.

Study Population and Selection Criteria

Our study population included all patients less than 18 years old who were admitted to a participating PHIS hospital. We excluded patients who were admitted to the hospital solely for research purposes. We selected a priori four individual study dates that occurred in the middle of each of the four seasons. Selected study days were Wednesdays: May 3, 2017 (spring), August 2, 2017 (summer), October 25, 2017 (fall), and January 31, 2018 (winter). We limited our assessments to hospitals for which antibiotic use data were available for all four study days.

We classified each encounter according to clinical service line (medical versus surgical) determined by all patient-refined diagnosis-related groups (APR-DRGs, version 32, 3M) service line classifications, and we also characterized the setting of care according to clinical units using billing data: neonatal intensive care unit (NICU), pediatric intensive care unit (PICU), and all other units/wards. We further classified hospitalizations according to the census region where the hospitals were located: Midwest, Northeast, South, and West.

Additionally, we characterized children according to the presence of complex chronic conditions. These were defined using a validated method for characterizing coded diagnoses-based pediatric complex chronic conditions, represented by 11 categories: neuromuscular, cardiovascular, respiratory, renal, gastrointestinal, hematologic or immunologic, metabolic, malignancy, genetic or other congenital defect conditions, premature, and neonatal [13]. Furthermore, to ascertain the clinical complexity of the patient population of each individual hospital, we used the case-mix index (CMI), a widely used surrogate for severity of illness and risk of mortality. In PHIS, CMI is based on APR-DRGs categories and severity levels and is calculated as the ratio of the average charge for encounters in a particular APR-DRG category/severity level combination to the average charge for all encounters included in the PHIS database [14].

Antibiotic Classifications

We included all antibiotics administered via oral, intravenous, or inhalational route and excluded those administered via other routes, such as topical, otic, and ophthalmic (see Supplementary Table 1 for full list of antibiotics included). We defined broad-spectrum antibiotics as previously published [15], and included additional antibiotics with increasing availability in recent years. The antibiotics considered broad-spectrum for this study included vancomycin, daptomycin, ceftolozane/tazobactam, cefepime, ceftaroline,

ceftazidime/avibactam, piperacillin/tazobactam, ticarcillin/clavulanate, carbapenems, fluoroquinolones, and linezolid.

Statistical Analysis

We described sociodemographic and clinical features of the study population, including age, gender, race, payor source, hospitalization day (number of days admitted prior to study day), mechanical ventilation status, presence of a complex chronic condition, and primary reason for hospitalization, as identified using APR-DRGs (version 32). We calculated the overall prevalence of antibiotic use in the study population by dividing the total number of patients who received at least one antibiotic on the study day by the total number of patients hospitalized for non-research purposes in a study hospital on that day. Ninety-five percent confidence intervals were calculated for these proportions. Prevalence of broad-spectrum antibiotic use was calculated similarly. Prevalence of overall and broad-spectrum antibiotic use was also stratified and compared among seasons (spring, summer, fall, winter), clinical service lines (medical/surgical), unit of hospitalization (NICU, PICU, all other units/wards), and census region. In secondary analyses, we assessed antibiotic use in the subset of hospitals with cardiac ICU patients within the PICU as well as the subset of hospitals with hematology-oncology units within the other units/wards encounters. We also assessed the correlation between the prevalence of overall and broad-spectrum antibiotic use and hospital median CMI. As a sensitivity analysis, we repeated our assessments after exclusion of healthy newborns given their unique indication for hospitalization. The most commonly used antibiotics were also identified. If a child received more than one antibiotic on the study date, each antibiotic used was considered separately. The most common antibiotic combinations were summarized and ranked. Categorical variables are expressed as proportions and were compared using χ^2 . Analyses were conducted using Stata 15.1 (StataCorp, College Station, TX) or SAS version 9.4 (SAS Institute, Cary, North Carolina).

RESULTS

Study population

Overall, we observed 52,769 encounters among 48,692 distinct patients across four seasons, with 1,850 patients appearing in more than one season. Patient and encounter characteristics are summarized in Table 1 and Figure 1. Almost a third (30.6%) of the study population were infants younger than 1 month of age, and children younger than 5 years accounted for 63.2% of all hospitalized children. Males comprised a slight majority of patients. About 62% of hospitalizations were attributed to a medical service line, and 63.3% had complex medical conditions.

Prevalence of antibiotic use

Overall, 36.3% of hospitalized children received antibiotics on the day of the study, 12.5% received at least one broad-spectrum antibiotic (Table 1), and 14.6% received two or more antibiotics. Of the antibiotics administered, 71% were delivered intravenously, 27.7% were given orally, and 1.3% were inhaled. Children younger than one month of age comprised 30.6% of the study population but only 19.2% and 14.8% of those receiving any or broad-spectrum antibiotics, respectively. Children with reported white race represented 47.4% of

the study population but accounted for only 14.8% of the patients receiving broad-spectrum antibiotics, while Asian children represented 3.2% of the study population but accounted for 23.5% of those receiving broad-spectrum antibiotics. Patients with a complex chronic condition accounted for 63.3% of hospitalized children and represented 72.2% of children receiving any antibiotic and 84.9% of children who received broad-spectrum antibiotics. One-third (36.6%) of encounters were assigned an infection diagnostic code at discharge, indicating an infection was present upon admission or was diagnosed at some point during the hospitalization. This proportion rose to 57.8% among those receiving any antibiotic and to 71.4% among those receiving a broad-spectrum antibiotic. A quarter (26.7%) of all medical encounters were assigned an infection diagnostic code, which included 57.7% of medical patients receiving any antibiotic and 65.5% of those receiving a broad-spectrum antibiotic. Over half (52.4%) of surgical encounters included an infection diagnostic code, which included 57.9% of surgical patients receiving any antibiotic and 76.4% of those receiving a broad-spectrum antibiotic. After exclusion of healthy newborns from the analysis (n=1559 [2.95%] hospitalizations), there was little change in the prevalence of overall and broad-spectrum inpatient antibiotic use (Supplementary Table 2).

Variability in inpatient antibiotic use

Prevalence in inpatient antibiotic use varied little by season (Supplementary Table 3), but significant variation was observed by service line, unit, and geographic region (Figure 2). Overall prevalence of antibiotic use and prevalence of broad-spectrum antibiotic use was higher among surgical patients (47.7% and 17.5%, respectively) than medical patients (29.2% and 9.3%, respectively, $p < 0.001$). Similarly, patients in the PICU had a higher prevalence of any antibiotic (58.3%) or broad-spectrum (26.2%) antibiotic use than patients in the NICU (23.1% and 6.5%, respectively) or other units/wards (36.4% and 11.6%, respectively, $p < 0.001$). In the 13 hospitals with designated hematology-oncology units, 56.3% of patients in these units received an antibiotic, and 31.1% received a broad-spectrum antibiotic. By census region, overall and broad-spectrum antibiotic use was most prevalent in hospitals located in the West region (40.2% and 14.6%, respectively) than in other regions, with the Midwest region exhibiting the lowest prevalence (32.7% and 10.9%, $p < 0.001$ for both overall and broad-spectrum variation in antibiotic use by region).

Prevalence of overall inpatient antibiotic use and broad-spectrum antibiotic use also varied by hospital, from a low of 22.3% to a high of 51.9% for overall and from 3.9% to 23.7% for broad-spectrum antibiotic use (Figure 3, $p < 0.001$). Importantly, there was no statistically significant relationship between median CMI (a surrogate for clinical complexity of cases seen at each hospital) and overall antibiotic use prevalence (Spearman's $\rho = 0.06$, $p = 0.674$) and only a weak (Spearman's $\rho = 0.29$), although statistically significant ($p = 0.036$), relationship between median CMI and broad-spectrum antibiotic use (Figure 4).

Most commonly used antibiotics in children's hospitals

The prevalence of the most commonly used antibiotics among all patients, including healthy newborns, according to clinical service line and inpatient unit are displayed in Figure 5 and are listed in Supplementary Table 4. Ceftriaxone and ampicillin were most commonly used among medical encounters, and cefazolin and vancomycin were most commonly used

in surgical encounters. The most frequently used antibiotics in the NICU were ampicillin and gentamicin. Those most frequently used in the PICU were cefazolin and vancomycin. In other units, cefazolin and ceftriaxone were used most frequently. The most common antibiotic combinations were ampicillin/gentamicin (9.7% of patients who received 2 antibiotics), cefepime/vancomycin (4.4%), and piperacillin/tazobactam/vancomycin (3.9%). There was little variability in the most common combinations of antibiotics by encounter type (Supplementary Table 5).

DISCUSSION

We found that inpatient antibiotic use is common and highly variable among children admitted to freestanding US children's hospitals and observed variability in the prevalence of antibiotic use by hospital service line and inpatient unit, with highest overall and broad-spectrum antibiotic use in surgical services and in the PICU. Although there was minimal variability in antibiotic use among seasons, antibiotic use prevalence varied by census region, with lowest use in the Midwest and highest use in the West. Previous studies have assessed regional differences in antibiotic prescribing in outpatient pediatric settings, but our study characterizes these differences in the inpatient setting. Importantly, although considerable variation in antibiotic use was observed across hospitals, this variability was not fully explained by patients' case-mix differences.

The overall prevalence of antibiotic use among hospitalized children in 2017–18 in our study was 36.3%, higher than the prevalence reported among children (30.7%) in a 2011 study of inpatient use of antibiotics among both children and adults hospitalized in the US [11]. Overrepresentation of adult patients relative to children in the previous study may partially explain this apparent discrepancy. Nevertheless, our assessment focused exclusively on children, assessed antibiotic use in single-day time points across all seasons in a one-year period, and demonstrated that the prevalence of inpatient antibiotic use remained relatively consistent throughout the year. In the 2011 study, the most commonly used antibiotic in the inpatient setting was vancomycin. In contrast, the most commonly used antibiotic in hospitalized children in our study was cefazolin, an antibiotic commonly used for surgical prophylaxis. A recent cross-sectional study reported that nearly a third (32.9%) of European children's hospital inpatients received an antibiotic for prophylactic use (73.4% for medical diseases and 26.6% for surgical diseases) [16]. Another study previously assessed antibiotic use in children admitted to hospitals participating in PHIS in 2008, before widespread implementation of inpatient antibiotic stewardship initiatives, and reported that about 60% of hospitalized children received antibiotics at some point during their hospital stay and also described similar regional variability to that observed in our current study [15]. Differences in study design and measurement approaches likely account for some of the differences between the results of that study and our observed prevalence of 36.6%, as our cross-sectional study focused on a single point-in-time assessment.

The lack of correlation between CMI and the prevalence of overall antibiotic use suggests that other patient-, provider- or hospital-associated factors may be contributing to unwarranted variability in inpatient antibiotic use. A retrospective study of 2010–2014 data found that antibiotic use was higher in PICUs than other wards and that use varied

significantly across PICUs, indicating that the PICU could be a good target for antimicrobial stewardship activities [17]. Additional studies are needed to identify specific drivers of variability and to optimize approaches to antibiotic stewardship activities. This is of particular importance as benchmarking appropriate inpatient antibiotic use and developing methods for objective inter-hospital comparison has become an important priority among hospitals, professional societies, and regulatory bodies [18].

Our study has important strengths including detailed characterization of inpatient services provided to hospitalized children with a broad range of medical and surgical conditions and a large sample size with representation of many large freestanding children's hospitals in the US. Unlike studies that focus on information based on written or electronic prescriptions or pharmacy fills to characterize antibiotic use, a distinct advantage of the current study relates to the source of antibiotic use information, extracted from charges related to antibiotics administered in the inpatient setting, which captures both the providers' preference and actual antibiotic receipt rather than prescription.

Our observations should be interpreted in light of several limitations. The specific indications for antibiotic use, and whether they were used for treatment or prophylaxis, were not available for the study. Although other studies have conducted selected chart reviews to try to obtain this information, we used the primary reason for hospitalization as adjudicated based on discharge diagnoses and procedures through the standardized diagnosis-related group adjudication process. Without more specific diagnostic information, we were unable to assess the appropriateness of antibiotic use as other studies have done using national prescribing guidelines [19]. The cross-sectional nature of our assessment precluded a more detailed characterization of the magnitude and duration of antibiotic exposure among hospitalized children. Our cross-sectional approach may lead to underrepresentation of antibiotic use of very short duration. Additionally, our prevalence figures cannot be directly compared to other measurements, such as days of treatment (DOT) per 1,000 patient days used in other studies. Furthermore, by reporting data only from a single one-year period, it is unclear whether our findings reflect underlying trends in antibiotic use following efforts to improve antibiotic use through increased implementation of ASPs. Future assessments including multiple years would allow characterization of trends in antibiotic use in children's hospitals and may be used to assess the impact of specific policies or interventions. Individual hospital ASP data, such as whether a program exists and when it was established, were not readily available for our study. These factors could explain some of the variability in antibiotic use across hospitals. Our findings are based on data from large freestanding children's hospitals participating in a collaborative network of hospitals, therefore caution is warranted when attempting to extrapolate our findings to other inpatient pediatric settings. Previous research has shown differences in antibiotic use among children's and non-children's hospitals [20].

In conclusion, antibiotic use among children hospitalized in the US is common and varies substantially according to hospital, clinical unit and service type, and census region. Variability seen among individual hospitals does not appear to be explained by the clinical complexity of their patients. Additional studies should identify the main drivers of inpatient antibiotic use variability to inform antibiotic stewardship activities.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support: Hannah G. Griffith's participation in this research was funded by Vanderbilt University.

Potential conflict of interest: Carlos G. Grijalva has received consulting fees from Pfizer, Merck, and Sanofi, and has received research support from Sanofi, the Agency for Healthcare Research and Quality, the National Institutes of Health, the US Food and Drug Administration, and the Centers for Disease Control and Prevention, outside the submitted work. Leigh M. Howard has received research support from Pfizer and the National Institutes of Health, outside the submitted work. Ritu Banerjee has received research support from Biomerieux, Roche, and BioFire, outside the submitted work. Derek J. Williams has received in kind support from Biomerieux, outside the submitted work. All other authors report no potential conflicts.

Abbreviations:

APR-DRG	All patient-refined diagnosis-related groups
ASP	Antibiotic stewardship program
CDC	Centers for Disease Control and Prevention
CMI	Case-mix index
DOT	Days of treatment
NICU	Neonatal intensive care unit
PHIS	Pediatric Health Information System
PICU	Pediatric intensive care unit

REFERENCES

1. Hales CM, et al., Trends in Prescription Medication Use Among Children and Adolescents-United States, 1999–2014. *JAMA*, 2018. 319(19): p. 2009–2020. [PubMed: 29800213]
2. Dellit TH OR, McGowan JE Jr., et al., Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*, 2007. 44(2): p. 159–177. [PubMed: 17173212]
3. Ingram PR SJ, Budgeon CA, Murray R, Point-prevalence study of inappropriate antibiotic use at a tertiary Australian hospital. *Internal medicine journal*, 2012. 42(6): p. 719–721. [PubMed: 22697156]
4. Camins BC KM, Wells JB, et al., mpact of an antimicrobial utilization program on antimicrobial use at a large teaching hospital: a randomized controlled trial. *nfection control and hospital epidemiology :the official journal of the Society of Hospital Epidemiologists of America*, 2009. 30(10): p. 931–938.
5. Levin PD IS, Sprung CL, et al., Antimicrobial use in the ICU: indications and accuracy—an observational trial. *Journal of hospital medicine : an official publication of the Society of Hospital Medicine*, 2012. 7(9): p. 672–678.
6. Fridkin SK BJ, Fagan R, et al., Vital Signs: Improving Antibiotic Use Among Hospitalized Patients. *MMWR. Morbidity and mortality weekly report*, 2014. 63.
7. Josefsdottir KS, et al., Antibiotics impair murine hematopoiesis by depleting the intestinal microbiota. *Blood*, 2017. 129(6): p. 729–739. [PubMed: 27879260]

8. Westphal JF VD, Brogard JM, Hepatic side-effects of antibiotics. *Journal of Antimicrobial Chemotherapy*, 1994. 33(3): p. 387–401.
9. Hersh AL, et al., Antimicrobial Stewardship Programs in Freestanding Children's Hospitals. *Pediatrics*, 2015. 135(1): p. 33. [PubMed: 25489018]
10. Gerber JS, et al., Identifying Targets for Antimicrobial Stewardship in Children's Hospitals. *Infection Control and Hospital Epidemiology*, 2013. 34(12): p. 1252–1258. [PubMed: 24225609]
11. Magill SS, et al., Prevalence of antimicrobial use in US acute care hospitals, May–September 2011. *JAMA*, 2014. 312(14): p. 1438–46. [PubMed: 25291579]
12. Bundy DG, et al., Association of Guideline-Adherent Antibiotic Treatment With Readmission of Children With Sickle Cell Disease Hospitalized With Acute Chest Syndrome Association of Antibiotic Treatment in Sickle Cell Disease With Readmission Rates Association of Antibiotic Treatment in Sickle Cell Disease With Readmission Rates. *JAMA Pediatrics*, 2017. 171(11): p. 1090–1099. [PubMed: 28892533]
13. Feudtner C, et al., Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. *BMC pediatrics*, 2014. 14: p. 199–199. [PubMed: 25102958]
14. Richardson T, et al., Development of Hospitalization Resource Intensity Scores for Kids (H-RISK) and Comparison across Pediatric Populations. *J Hosp Med*, 2018. 13(9): p. 602–608. [PubMed: 29694460]
15. Gerber JS, et al., Variability in antibiotic use at children's hospitals. *Pediatrics*, 2010. 126(6): p. 1067–73. [PubMed: 21078728]
16. Hufnagel M, et al., High Rates of Prescribing Antimicrobials for Prophylaxis in Children and Neonates: Results From the Antibiotic Resistance and Prescribing in European Children Point Prevalence Survey. *J Pediatric Infect Dis Soc*, 2019. 8(2): p. 143–151. [PubMed: 29579259]
17. Brogan TV, et al., Variability in Antibiotic Use Across PICUs. *Pediatr Crit Care Med*, 2018. 19(6): p. 519–527. [PubMed: 29533352]
18. Yu KC, et al., Benchmarking Inpatient Antimicrobial Use: A Comparison of Risk-Adjusted Observed-to-Expected Ratios. *Clinical Infectious Diseases*, 2018. 67(11): p. 1677–1685. [PubMed: 29688279]
19. Fleming-Dutra KE, et al., Prevalence of Inappropriate Antibiotic Prescriptions Among US Ambulatory Care Visits, 2010–2011 Inappropriate Antibiotic Prescriptions Among Ambulatory Care Visits, 2010–2011 Inappropriate Antibiotic Prescriptions Among Ambulatory Care Visits, 2010–2011. *JAMA*, 2016. 315(17): p. 1864–1873. [PubMed: 27139059]
20. Tribble AC, Ross RK, and Gerber JS, Comparison of Antibiotic Prescribing for Pediatric Community-Acquired Pneumonia in Children's and Non-Children's Hospitals Comparison of Antibiotic Prescribing for Community-Acquired Pneumonia by Hospital Type Letters. *JAMA Pediatrics*, 2019. 173(2): p. 190–192. [PubMed: 30534969]

Figure 1.

Flow diagram describing number and distribution of inpatient encounters in 51 PHIS hospitals on a study date, 2017–2018, by clinical unit and service line.

Figure 2.

Distribution of overall and broad-spectrum antibiotic use among children admitted to a PHIS hospital on a study date, 2017–2018, by US census region, clinical unit, and service line. Blue represents overall antibiotic use is represented. Red represents broad-spectrum antibiotic use. Asterisks (*) indicate significant variation in prevalence among groups.

Figure 3.

Prevalence of overall and broad-spectrum antibiotic use by hospital among US children's hospitals on a study date, 2017–2018. Diamonds represent prevalence of antibiotic use. Vertical lines represent 95% confidence intervals. Blue represents overall antibiotic use, and red represents broad-spectrum antibiotic use. Asterisk (*) indicates significant variation in prevalence among hospitals.

Figure 4.

Correlation between prevalence of antibiotic use and median hospital case-mix index (CMI). CMI is calculated using all patient-refined diagnosis-related group (AP-DRG) categories and severity levels. A higher CMI indicates a higher level of clinical severity. Blue circles represent prevalence of any antibiotic. Red circles represent prevalence of broad-spectrum antibiotic use.

Figure 5.

Most frequently used antibiotics among PHIS hospitals, 2017–2018 by service line (A), and by clinical unit (B). Purple dots represent medical use, and orange dots represent surgical use. Blue dots represent use in PICU. Red dots represent use in NICU. Yellow dots represent use in all other units and wards.

Table 1:

Characteristics of children admitted to one of 51 PHIS hospitals on a study date, 2017–2018

Characteristics		Overall, N (%)	With Antibiotics ¹ , n (%)	With Broad-Spectrum ² Antibiotics, n (%)
Number of patients (N)		52769	19174 (36%)	6575 (12%)
Age	0–29 days	16149 (31%)	3685 (19%)	976 (15%)
	30–364 days	8186 (16%)	3007 (16%)	868 (13%)
	1–4 years	9023 (17%)	4216 (22%)	1497 (23%)
	5–11 years	8916 (17%)	4100 (21%)	1544 (23%)
	12–17 years	10495 (20%)	4166 (22%)	1690 (26%)
Gender	Male	28298 (54%)	10389 (54%)	3630 (55%)
Race	White	25030 (47%)	9109 (48%)	976 (15%)
	Black	10142 (19%)	3378 (18%)	868 (13%)
	Hispanic	10087 (19%)	4086 (21%)	1497 (23%)
	Asian	1695 (3%)	624 (3%)	1544 (23%)
	Other	5815 (11%)	1977 (10%)	1690 (26%)
Census region	Midwest	14925 (28%)	4884 (25%)	1627 (25%)
	Northeast	7850 (15%)	2936 (15%)	1046 (16%)
	South	19028 (36%)	6951 (36%)	2296 (35%)
	West	10966 (21%)	4403 (23%)	1606 (24%)
Type of encounter ³	Medical	32436 (61%)	9470 (49%)	3020 (46%)
	Surgical	20333 (39%)	9704 (51%)	3555 (54%)
Infection diagnosis		19311 (37%)	11084 (58%)	4695 (71%)
Complex chronic condition (CCC) ⁴		33400 (63%)	13842 (72%)	5581 (85%)

¹ Antibiotics include any antibacterial antibiotic administered via the oral, intravenous, or inhalational route.

² Broad-spectrum antibiotics include vancomycin, daptomycin, cefepime, ceftaroline, ceftazidime-avibactam, ceftolozane-tazobactam, piperacillin-tazobactam, carbapenems, fluoroquinolones, and linezolid.

³ Each patient encounter day is coded as either medical or surgical based on service line charge data.

⁴ Complex chronic condition was defined per Feudtner et al. as “any medical condition that can be reasonably expected to last at least 12 months (unless death intervenes) and to involve either several different organ systems or 1 organ system severely enough to require specialty pediatric care and probably some point of hospitalization in a tertiary care center.”