Published in final edited form as:

J Chem Inf Model. 2020 March 23; 60(3): 1090–1100. doi:10.1021/acs.jcim.9b01156.

Tautomer Database: A Comprehensive Resource for Tautomerism Analyses

Devendra K. Dhaked, Laura Guasch, Marc C. Nicklaus

Computer-Aided Drug Design Group, Chemical Biology Laboratory, Center for Cancer Research, National Cancer Institute, Frederick, Maryland 21702, United States

Abstract

We report a database of tautomeric structures that contains 2819 tautomeric tuples extracted from 171 publications. Each tautomeric entry has been annotated with experimental conditions reported in the respective publication, plus bibliographic details, structural identifiers (e.g., NCI/CADD identifiers FICTS, FICuS, uuuuu, and Standard InChI), and chemical information (e.g., SMILES, molecular weight). The majority of tautomeric tuples found were pairs; the remaining 10% were triples, quadruples, or quintuples, amounting to a total number of structures of 5977. The types of tautomerism were mainly prototropic tautomerism (79%), followed by ring–chain (13%) and valence tautomerism (8%). The experimental conditions reported in the publications included about 50 pure solvents and 9 solvent mixtures with 26 unique spectroscopic or nonspectroscopic methods. ^1H and ^{13}C NMR were the most frequently used methods. A total of 77 different tautomeric transform rules (SMIRKS) are covered by at least one example tuple in the database. This database is freely available as a spreadsheet at <https://cactus.nci.nih.gov/download/tautomer/>.

Graphical Abstract

Corresponding Author Marc C. Nicklaus – Computer-Aided Drug Design Group, Chemical Biology Laboratory, Center for Cancer Research, National Cancer Institute, Frederick, Maryland 21702, United States; mn1@mail.nih.gov.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jcim.9b01156>.

Spreadsheet S1: List of the publications used in tautomer database generation (XLSX)

Spreadsheet S2: Distribution of solvents or their mixtures, or general experimental environments, by spectroscopic methods (XLSX)

Spreadsheet S3: SMIRKS of tautomeric rule (XLSX) Representative examples of chemical types of tautomerism (Table S1) (PDF)

Spreadsheet S4: Tautomer database_itself (XLSX)

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.jcim.9b01156>

The authors declare no competing financial interest.

INTRODUCTION

Tautomerism is a phenomenon in which a set of molecules can interconvert by movement of a hydrogen or group of atoms and/or molecular rearrangement. The movement of hydrogen atoms along with the migration of pi-bonds is called prototropic tautomerism. The intermolecular arrangements leading to isomerization due to ring opening or cyclization are known as ring-chain tautomerism. (We have recently compiled 11 sets of rules for ring-chain tautomerism in SMIRKS notation.¹) Another type of isomerization, which involves rapid reorganization of single and double bonds without migration of any atom or group, is termed valence tautomerism.

Tautomers usually have different physicochemical properties such as logP, hydrophobicity, pK_a , solubility, electrostatic potential, similarity index, etc. with concomitant computation of such properties and molecular descriptors yielding different results from one tautomer to another.² They may behave differently in docking tools, pose challenges in macromolecular X-ray structures, particularly for small-molecule ligands in protein-ligand complexes, and wreak havoc in compound registration systems and vendor catalogs.³ Therefore, consideration of tautomers has been of high interest to the drug design community for decades.⁴⁻⁷ This leads to the question: How does one select the tautomer(s) of a molecule that allow one to most accurately predict its properties? One of the issues in this context has been the lack of a publicly accessible database providing a significant number of quantitative ratios or qualitative data of tautomeric forms in different solvents.

While a significant body of work exists of individual experimental studies (e.g., spectroscopy on tautomeric molecules), quantum chemical analyses (e.g., energy and structure computations), and—to a limited extent—chemoinformatics studies (e.g., rule-based tautomer prediction), very few systematic collections of experimental results in this field have been undertaken so far. A set of 785 transformations belonging to 11 types of tautomeric reactions with equilibrium constants measured in different solvents and at different temperatures was recently used in an effort to build QSPR models of equilibrium constants of tautomeric molecules.⁸ To the best of our knowledge, there is currently no database publicly available that provides details of several thousands of molecules and their experimentally investigated tautomers under a wide variety of experimental conditions along with a detailed chemoinformatics analysis. We acknowledge, however, the Tautomer Codex database (Tautobase) and its references provided directly to us by Wahl and Sander before its recent publication.⁹ It provides a very useful complement of water-based tautomerism measurements in addition to the diversity of solvents represented in our database. Here, we report on a tautomer database we have created from the literature in an attempt to compile experimental and quantitative tautomeric preferences together with chemical and bibliographic information as well as an analysis along a set of more than 70 tautomeric transforms. We hope this resource, which we made freely available for download at <https://cactus.nci.nih.gov/download/tautomer/> starting October 2017, will allow the scientific community to more easily explore the phenomenon of tautomerism by finding several thousand such molecules in one location. (A preprint version of this paper with a more-extended discussion is available at [10.26434/chemrxiv.10790369.v1](https://doi.org/10.26434/chemrxiv.10790369.v1).)

METHODS AND DATA

Data Set of Database.

The current tautomer database consists of 2819 entries, each comprising an n -tuple of tautomers ($n = 2-5$) studied in a particular set of experimental conditions (pH, solvent, solvent mixture, temperature, experimental technique used). All these tuples together comprise a total of 5977 records. The data were extracted from 171 publications, which included a number of reviews (see full list in the Spreadsheet S1 of the Supporting Information). The initial extraction from these literature sources was done by a contract mechanism (Parthys Reverse Informatics, <http://www.reverseinformatics.com>), whereas a significant workup and curation of the initial data was performed by the authors.

For each entry for all n -tuples in the tautomer database, the corresponding NCI/CADD Chemical Structure Identifiers¹⁰ were calculated using the chemoinformatics toolkit CACTVS¹¹ (in which they have been implemented as standard molecular properties). The nature of these identifiers, which are based on the standard CACTVS molecular hashcodes, is based on turning off or on sensitivity to the following five chemical features: fragments, isotopes, charges, tautomers, and stereochemistry. In this database, we used the FICTS, FICuS, and uuuuu identifiers out of a possible 32 possible set of variants (see the original publication¹⁰ for explanation of the nomenclature). The FICTS identifier, in which all five features are turned on, represents the original input structure as is. It is sensitive to fragments (such as counterions), isotopes, charges, and stereochemistry in the input structure as well as to the specific tautomer drawn. The FICuS identifier is tautomer invariant (but sensitive to all four other features), meaning that different tautomers have the same FICuS hashcode. For the uuuuu identifier, all five features are turned off, implying that molecules differing by fragment, isotope, charge, tautomer, or stereochemistry have the same uuuuu (which can thus be regarded as a sort of parent structure identifier). The FICuS identifier is conceptually similar to the InChIKey, though the latter handles tautomerism less comprehensively than FICuS due to an only limited range of tautomerism transforms implemented in its current version (v. 1.05). Additionally, it is not currently possible for the user to add entirely new types of tautomerism to the InChI[Key] calculation. This and other shortcomings of the current InChI in the handling of tautomerism has led to an IUPAC-sanctioned project of redesigning the handling of tautomerism for an InChI V2,¹² for which this tautomer database forms an experimental backdrop of sorts and whose authors are involved in the IUPAC project.

In order to describe the tautomeric transformation(s) between the members of each of the tautomeric n -tuples in the database, we used a total of 77 rules. This set is closely related to, and essentially a major subset of, the 86 rules described in the context of redesigning of the handling of tautomerism for InChI V2 in the accompanying paper.¹³ All these rules were encoded in SMIRKS line notation.¹⁴ They were all processed in CACTVS, which comes with a default set of 20 prototropic rules covering a wide range of common and some rarer types of tautomerism. Twelve of the 20 rules have a representative in this database. To these, we added a subset of 8 SMIRKS from our recently published 11 types of ring $\rightleftharpoons$ chain rules

(encoded in a total of 38 SMIRKS),¹ plus 57 out of 61 heretofore unpublished rules, which are detailed in the accompanying publication.¹³

We use the following nomenclature (again aligned with the accompanying paper¹³) for the three types of rules discussed in this paper: (1) Prototropic tautomerism rules are called PT_*nn*_*mm*, where *nn* and *mm* are the number of the rule and a possible variant, respectively. The names of most rules end with a 00 indicator, indicating that there is only one variant. (2) Ring-chain tautomerism rules are named RC_*nn*_*mm*, where *nn* and *mm* have the same meaning as described above. (3) Valence tautomerism rules are termed VT_*nn*_*mm* according to the same scheme.

To determine the single transform or sequence of transforms connecting the tuple members with each other, we applied the following procedure: In the first step, we enumerated all possible tautomers from each tautomeric tuple. In the second step, we generated a tautomer network among those enumerated tautomers. In such a network, we typically have several pathways that connect one tautomer to the other by different tautomeric transforms. As the final step, we searched for the shortest pathway, defined by the smallest number of transformation steps within the tautomeric pair. If two different paths had the same number of steps, we used a notation of the type of {PT_03_00/PT_06_00} > PT_09_00. This means the pathway can either use PT_03_00 or PT_06_00 in the first step, followed by PT_09_00 in the second step.

Database Description.

The database is provided to the user as a spreadsheet in Excel format. Each entry consists of three major segments: conditions, tautomer, and publication. Each segment has several fields as listed in Table 1. For each additional (second, third, ...) tautomer of a compound, fields in the second (and third, etc.) instance of the tautomer-specific columns are populated with data, otherwise left empty. The following provides a brief explanation of some key columns in the spreadsheet. Others should be self-explanatory. A legends worksheet is also available in the spreadsheet providing explanations for all columns.

- A.** Size: Number of tautomers reported in the publication as being in equilibrium. In a few publications, only the main tautomer of the compound was described; in such cases, we entered a second entry based on a possible (calculated) tautomer.
- B.** Solvent: Solvent in which the tautomers were observed. This can be a mixture of solvents. If their concentration is indicated, then it is also mentioned in the solvent column.
- C.** Solvent_Proportion: Fraction of solvents or their mixtures; typically measured on a mass, molar, or volume ratio scale (though in some cases, the scale used was not clear from the publication).
- D.** Solvent_Mixture: Indicates whether a single solvent or mixture was used. This column has a “yes” if the “Solvent” column indicates a solvent mixture, otherwise “no”.

- E.** Temperature: Temperature (K) at which the tautomers were observed or the experiment was carried out. In the case of mass spectroscopy experiments, the temperature of the injector was used as the experimental temperature.
- F.** pH: pH of the medium at which the tautomers were observed or experiment was conducted.
- G.** Experimental_Method: This describes the spectroscopic or physical methods that were used in the experimental determination of the tautomers. It may be a single method or a combination of several methods by which the tautomers were established in the experiments. If the experimental details were not available in the review, then those were extracted from the original references.
- H.** Entry_ID1: Unique ID composed from the publication reference (journal name, year, volume, page numbers) along with the tautomer ID in that publication (if given) and the nature of the tautomerism (e.g., “Keto $\rightleftharpoons$ enol”).
- I.** Type_1: The chemical nature of the tautomer, e.g., keto, hydroxy, imine, enamine, etc. An entry with “nul” in this column indicates that it was difficult to assign any specific name from the molecule’s common name or based on similar structures in the database.
- J.** Transf_1_2: The rule(s) (prototropic [PT], ring $\rightleftharpoons$ chain [RC], or valence tautomerism [VT]) which transform(s) tautomer_1 into tautomer_2 (single or multiple steps). A forward slash “/” is used to indicate alternative rules for any step. Curly braces “{ }” are used to group together alternative rules if these appear in multi-step transforms. The greater than sign “>” is used to separate steps in multi-step transformations. An entry with “no_transform” in this column indicates that these pairs are not covered by our rules because these examples are related to zwitterionic and complex protonated structures; hence, we did not develop any rules for them.
- K.** ID_Hash_1: A hashed unique ID generated for each tautomer by the original contractor (Reverse Informatics). Some entries added later by us do not have an ID_Hash.
- L.** FICTS_1: Tautomer-sensitive NCI/CADD structural identifier of tautomer_1.
- M.** HASHISY_1: Tautomer-sensitive CACTVS structural identifier of tautomer_1.
- N.** FICuS_1: Tautomer-insensitive NCI/CADD identifier, which therefore is the same for all tautomers of the same molecule.
- O.** uuuuu_1: NCI/CADD identifier for the parent compound of tautomer_1.
- P.** Quantitative_ratio_1: Quantitative ratio of tautomer_1 compared to other tautomers. This can be a single number, a range, or an upper or lower bound between 0 and 1. Decimal numbers are reported up to the third decimal digit.
- Q.** Qualitative_Prevalence_1: Qualitative prevalence category of tautomer_1 reported in the publication. These keywords describe the prevalence of one tautomer over other tautomer(s) and are mostly extracted from the papers,

assigned based on the quantitative data or the spectra or other informations in the text of paper. “nul” is used if no such keywords were available in the papers or reviews.

- R.** Prevalence_Category_1: In order to make both quantitatively and qualitatively reported prevalences of tautomers comparable, at least in a categorical way, we numerically categorized tautomer_1 into five classes: 0, 1, 2, 3, and 4 based on its quantitative ratio and/or qualitative prevalence as described below.

Numeric classification of qualitative prevalence's keywords:

0: Not observed

1: Less favored, less stable, minor, observed

2: Equally, favored, major, in equilibrium, preferred, similar spectra

3: More favored, more stable, predominant, strongly favored

4: Exclusively observed, only observed, only tautomer, identical tautomer

Numeric classification of quantitative amount of tautomers

0: Quantitive_ratio = 0.0–0.0099

1: Quantitive_ratio = 0.01–0.30

2: Quantitive_ratio = 0.31–0.69

3: Quantitive_ratio = 0.70–0.99

4: Quantitive_ratio = 1

If there were three or more tautomers reported, there would be corresponding columns in the spreadsheet with “3” or “_3”, e.g., Transf_1_3, etc.

DATABASE ANALYSIS

Provenance and Relationship of Tuples.

We did not identify any direct tuples' duplicates in terms of both chemical structure and experimental conditions. Purely chemical duplicates were found for 479 tautomeric tuples in the database, but they differ in conditions such as temperature, solvent, pH, or spectroscopy method.

Size Distribution of Tuples.

The database contains tautomeric tuples ranging in size from 2 to 5 (2530, 250, 28, 11 cases, respectively) (Figure 1).

Solvent.

The database contains tautomeric equilibrium studies performed in solvents (87% of the cases), in the solid state (6%), neat liquid (<1%), gas phase/matrix (5%), and vapor phase (<1%). The majority of experiments were conducted in some kind of solvent or solvent mixture. About 50 different types of solvents were reported in the papers (Figure 2a). The database has 12 solvent mixtures, in which 12 different types of solvents were used (Figure 2b).

Temperature Distribution.

Experimental temperature information was available for 1389 entries in the form of either exact value, range, room (RT), or ambient temperature. About 82% of the studies represented in the database were carried out at a temperature range of 250–350 K (Figure 3). The majority of those (50%) were carried out in the range of 251–300 K. There were only 53 entries below 201 K and 23 entries at a higher temperature (e.g., 523 K).

pH Distribution.

The database has experimental pH details for 100 entries, 63% of which were reported to have used an acidic medium (Figure 4a). For 91 entries of 2-tautomer sets and 9 entries of 3-tautomer sets reported in pH based studies, medium polar to polar solvents (98%) or their mixtures (2%) were used (Figure 4b). These studies used the following spectroscopy methods: ^1H NMR, flash photolysis, Raman, UV, and UV/vis. Of these, UV/vis spectroscopy was used in 79% of the cases with methanol, acetonitrile, and DMSO-water.

Experimental Methods.

In most of the studies (85%), a single spectroscopy or physical method was used, while in the remainder of the studies two to three methods were used, often by way of an additional method used as support of the primary method. In the multiple method studies, spectroscopic methods from ^1H , ^{13}C , ^{14}N , ^{15}N , ^{17}O , and/or ^{31}P NMR spectroscopy were the most common (~75% of the cases). Out of the total 29 unique methods, ^1H NMR (1014), ^{13}C NMR (340), UV (253), IR (172), and UV/vis (139) were the top five methods (Figure 5a). In the multiple method studies (Figure 5b), ^1H NMR and ^{13}C NMR were frequently used together (131). In addition, ^1H NMR was commonly used together with other methods such as ^{31}P , ^{15}N and ^{17}N NMR, and IR. Some of the spectroscopic methods used different types of solvents; for example, ^1H NMR, UV, IR, ^{31}P NMR, and ^{13}C NMR methods were performed in 41, 22, 19, 20, and 14 different solvents and solvent mixtures, respectively. Chloroform and DMSO were the most important solvents in ^1H NMR (279 and 227 cases, respectively) and ^{13}C NMR (182 and 84 cases, respectively) (Spreadsheet S2, Supporting Information). In IR, chloroform (59) and nujol (38) were used extensively. In UV/vis, methanol (89) and acetonitrile (8) were used extensively. In UV, ethanol (76) and water (31) were used extensively.

Analysis by Tautomeric Transform Rules.

As already mentioned, we used as the starting point for the tautomeric rule compilation (a) 20 standard prototropic rules (default CACTVS rules PT_02_00 – PT_21_00) and (b) 11 ring $\rightleftharpoons$ chain (RC_01_00 – RC_11_00) rules (in 38 SMIRKS strings) that have been published by our group recently.¹ In addition, we have compiled¹³ 61 new tautomeric rules derived from various literature sources. These new rules consist of 34 prototropic rules (PT_22_00 – PT_49_00) including two variants with *mm* > 00 and variants of PT_11_00 for long-range hydrogen migration, where *mm* ranges from 01 to 04, 16 ring $\rightleftharpoons$ chain rules (RC_03_03, RC_03_04, RC_04_04, and RC_12_00 – RC_24_00), and 11 valence rules (VT_01_00, VT_01_01 – VT_10_00). (See footnotes of Table 2 for rule naming and numbering nomenclature.)

Table 2 shows the frequency of the applicability of all these rules to the entries in our database showing both the cases where the transformation between the experimental tautomers only required the application of a single rule as well as of cases that needed additional, or allowed alternative, rules in the single- or multi-step transformation between observed tautomers.

The most commonly encountered prototropic, ring $\rightleftharpoons$ chain, and valence rules are shown in Figure 6. The majority of transformations from our database occur in a single step (60%), while the others involve the use of additional rules to complete the transformation. About 35% of the transformations are achieved by the application of PT_06_00 and PT_07_00 in a single step. Some rules (PT_02_00 to PT_05_00, PT_08_00, PT_10_00 to PT_16_00) appeared only in multi-step transformations or as alternative rules to others. Here, 353 cases needed an additional one step (for a total of two steps), and 27 other cases required two or more steps (for a total of three or more steps) to complete the observed tautomeric transformations.

Most frequently, a hydrogen atom migrates in a tautomeric transformation from its initial position in the molecule to an odd numbered (relative) position (such as 3, 5, 7, 9, or 11), designated as “1,3 H-shift,” “1,5 H-shift”, etc. Migration to an even position (such as 2, 4, or 6) is rare. In most cases, hydrogen migrated via 1,3 H-shift (1,120), followed by 1,5 H-shift (707) and 1,7 (91) H-shift, respectively, in the single-step transformations. One notes that this distance traveled by the hydrogen is well correlated with the observed frequency of H-shifts. The 1,3 H-shift can alternatively be achieved via long distance migration using 1,5 H-shift (30) or 1,7 H-shift (118), respectively. Likewise, 1,5 H-shift based transformations can be in competition with 1,7 H-shift and 1,9 H-shift in single-step equilibria. For two-step transformations, we observed the order by frequency of occurrence shown in Table 3. We note that, as in single-step transformations, shorter distance hydrogen migrations are more prevalent than longer ones.

The database contains significantly fewer cases (388) of ring $\rightleftharpoons$ chain tautomerism than of prototropic tautomerism. They generally belong to cyclization to 4-, 5-, and 6-membered ring systems, which can occur either via an endocyclic or exocyclic process where the double bond becomes part of the ring or the side chain, respectively. In 180 cases of endocyclic ring $\rightleftharpoons$ chain transformations, ring closure happened at digonal (sp), trigonal

(sp²), or tetrahedral (sp³) centers. The three rules RC_12_00, RC_18_00, and RC_24_00 do not follow the concept of ring closing and ring opening according to Baldwin's rules. In contrast to other rules, RC_24_00 involves tautomerization between trivalent (chain) and pentavalent (ring) tautomers. There are some rules that involve a 1,2 H-shift (i.e., RC_24_00), 1,4 H-shift (i.e., RC_15_00, RC_20_00 and RC_23_00), and 1,5 H-shift (RC_22_00) during ring closure.

In 193 cases of exocyclic ring⇌chain transformations, the ring closing and opening took place at trigonal or tetrahedral centers. There were some instances of ring⇌chain tautomerism in the thiadiazoline (RC_14_00), boryl/borate (RC_16_00 and RC_17_00), and λ⁵/λ³-phosphane (RC_24_00) systems that did not involve any unsaturated electrophilic center (or endocyclic or exocyclic bonds) during interconversions but rather involved saturated sulfur, boron, and phosphorus centers, respectively.

The ring–chain rules did not occur in combination with any prototropic or ring⇌chain rule; i.e., in all cases, transformation proceeded in a single step. Generally, ring⇌chain tautomerism showed a high prevalence for the chain form over of the ring form. There were 20 cases of ring⇌chain tautomerism where three tautomers are in equilibrium with each other in solution, the two ring tautomers existing as *cis* and *trans* isomers, respectively.

There are 228 cases of valence tautomerism in the database. They all involved ring opening or closing in 4-, 5-, or 6- membered ring systems without migration of any hydrogen atom. The ring-opened tautomers of four rules (VT_02_00, VT_04_00, VT_05_00, and VT_08_00) have a charge-separated moiety in their structures, and this charge disappears in the ring-closed tautomers. In contrast hereto, a charge-separated moiety is present in the ring-closed tautomer of both VT_07_00 and VT_10_00. The tautomeric equilibrium via VT_06_00 involves ring-contraction (6-membered) and ring-expansion (7-membered) in the tautomers. VT_09_00 is the only one rule that involves a valency change during tautomerization: between trivalent phosphinoimine and pentavalent diazaphosphazole tautomers. Among the 11 valence tautomerism rules, our database contains significant counts only for the tetrazole⇌azide tautomerism (VT_02_00), with the tetrazole tautomer being more favored in a polar aprotic solvent and the azide tautomer in the nonpolar solvent.

Type of Tautomerism.

Many of the transforms listed in Table 2 align quite closely with chemotypes the way the organic chemist would usually perceive them. However, others among these transforms, as they are expressed as general SMIRKS patterns,¹³ cover a broader range of compound types. For example, transform PT_06_00 (1,3 heteroatom H-shift) recognizes C, O, N, S, P, Se, and Te in its SMIRKS pattern, thus covering quite diverse types of compounds and tautomerism based on those. Conversely, the interconversion between the hydrazine and the azo species of a compound can be affected at the transform level by a 1,3 H-shift, 1,5 H-shift, and 1,7 H-shift, which are encoded in different transforms. Table 4 shows the distribution of the records along more than 50 chemical types of tautomerism (see molecular examples in Table S1, Supporting Information, which also contains a more extensive discussion of the tautomer types). Table 5 shows commonly identified sets of three tautomers with their occurrences.

Table 6 shows the distribution of some of the common tautomers across the five different prevalence categories described above (0–4).

SUMMARY AND CONCLUSIONS

A significant variety of structures, chemotypes, analytical procedures, and experimental conditions including solvents has been compiled to form the Tautomer Database. We hope that this database of experimental data and its included analysis by chemoinformatics methods (by way of annotation with tautomeric transform rules) may provide a set of data useful for future work in the field of tautomerism. This would include tools such as software and chemical identifiers that could be used to avoid tautomeric duplication in chemical databases and compound registration systems. We also hope it may help in developing approaches to predict the most “medicinally” relevant and “reasonable” tautomer forms. This data set could be a useful training set for machine learning models based on quantum mechanics^{15,16} to rapidly identify the lowest energy tautomer.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We have to send copious thanks to Wolf-Dietrich Ihlenfeldt for his initial work with CACTVS and its treatment of tautomerism, as well as for his support in our generating and testing the new rules. We gratefully acknowledge Thomas Sander and Oya Wahl for providing us with a copy of their Tautomer Codex database, which helped in the generation of a handful of additional rules. This work utilized the computational resources of the NIH HPC Biowulf cluster (<http://hpc.nih.gov>). This work was supported by the Intramural Research Program of the National Institutes of Health, Center for Cancer Research, National Cancer Institute. All authors received funding from the NCI, NIH, Intramural Research Program. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

REFERENCES

- (1). Guasch L; Sitzmann M; Nicklaus MCEnumeration of Ring–Chain Tautomers Based on SMIRKS Rules. *J. Chem. Inf. Model*2014, 54 (9), 2423–2432. [PubMed: 25158156]
- (2). Martin YCLet’s Not Forget Tautomers. *J. Comput.-Aided Mol. Des*2009, 23 (10), 693–704. [PubMed: 19842045]
- (3). Guasch L; Yapamudiyansel W; Peach ML; Kelley JA; Barchi JJ; Nicklaus MCExperimental and Chemoinformatics Study of Tautomerism in a Database of Commercially Available Screening Samples. *J. Chem. Inf. Model*2016, 56 (11), 2149–2161. [PubMed: 27669079]
- (4). Masand VH; Mahajan DT; Gramatica P; Barlow JTautomerism and Multiple Modelling Enhance the Efficacy of QSAR: Antimalarial Activity of Phosphoramidate and Phosphorothioamidate Analogues of Amiprofos Methyl. *Med. Chem. Res*2014, 23 (11), 4825–4835.
- (5). Milletti F; Vulpetti ATautomer Preference in PDB Complexes and Its Impact on Structure-Based Drug Discovery. *J. Chem. Inf. Model*2010, 50 (6), 1062–1074. [PubMed: 20515065]
- (6). Kalliokoski T; Salo HS; Lahtela-Kakkonen M; Poso ATThe Effect of Ligand-Based Tautomer and Protomer Prediction on Structure-Based Virtual Screening. *J. Chem. Inf. Model*2009, 49 (12), 2742–2748. [PubMed: 19928753]
- (7). Oellien F; Cramer J; Beyer C; Ihlenfeldt W-D; Selzer PMThe Impact of Tautomer Forms on Pharmacophore-Based Virtual Screening †. *J. Chem. Inf. Model*2006, 46 (6), 2342–2354. [PubMed: 17125178]

- (8). Gimadiev TR; Madzhidov TI; Nugmanov RI; Baskin II; Antipin IS; Varnek AA Assessment of Tautomer Distribution Using the Condensed Reaction Graph Approach. *J. Comput.-Aided Mol. Des* 2018, 32 (3), 401–414. [PubMed: 29380104]
- (9). Wahl O; Sander TT Autobase: An Open Tautomer Database. *J. Chem. Inf. Model* 2020, DOI: 10.1021/acs.jcim.0c00035.
- (10). Sitzmann M; Filippov IV; Nicklaus MC Internet Resources Integrating Many Small-Molecule Databases 1. *SAR QSAR Environ. Res* 2008, 19 (1–2), 1–9. [PubMed: 18311630]
- (11). Xemistry Chemoinformatics <https://www.xemistry.com/> (accessed 29–01–2020).
- (12). IUPAC projects https://iupac.org/projects/project-details/?project_nr=2012-023-2-800 (accessed 29–01–2020).
- (13). Dhaked DK; Ihlenfeldt W-D; Patel H; Delanneé V; Nicklaus MC Toward a Comprehensive Treatment of Tautomerism in Chemoinformatics Including InChI V2. *J. Chem. Inf. Model* 2020, DOI: 10.1021/acs.jcim.9b01080.
- (14). Daylight Theory Manual <https://www.daylight.com/dayhtml/doc/theory/theory.smirks.html> (accessed 29–01–2020).
- (15). Smith JS; Isayev O; Roitberg AE ANI-1: An Extensible Neural Network Potential with DFT Accuracy at Force Field Computational Cost. *Chem. Sci* 2017, 8 (4), 3192–3203. [PubMed: 28507695]
- (16). Smith JS; Isayev O; Roitberg AE ANI-1, A Data Set of 20 Million Calculated off-Equilibrium Conformations for Organic Molecules. *Sci. Data* 2017, DOI: 10.1038/sdata.2017.193.

Figure 1.
Distribution of sizes of tautomer tuples in the database.

Figure 2. Frequency distribution of most commonly used (a) solvents and (b) solvent mixtures.

Distribution of Experimental Temperature

Figure 3. Temperature range distribution of experimental studies. (The range of 251–300 K includes studies that simply reported “room temperature”).

Figure 4.

(a) Distribution of studies at different pH ranges. (b) Distribution of different solvents used in pH based studies.

Figure 5. Frequency distribution of (a) single experimental methods and (b) multiple experimental methods (only top 15 methods are shown).

Figure 6.
Frequency distribution of commonly observed rules.

Table 1.Summary of Data Fields Used in Tautomer Database^a

Conditions	Tautomer_1	Publication_1
ref	Entry_ID1	Filename_1
Size	Type_1	Publication_DOI_1
Solvent	Transf_1_2	Publication_ID_1
Solvent_Proportion	ID_Hash_1	Authors_1
Solvent_Mixture	FICTS_1	Affiliation_1
Temperature	HASHISY_1	Title_1
pH	FICuS_1	Section_1
Experimental_Method	uuuu_1	Page_Number(s)_1
Solvent_Mixture	Std_InChIKey_1	Notes_1
	Std_InChI_1	Cmpd_Number_1
	SMILES_1	
	Mol_Formula_1	
	Mol_Weight_1	
	IUPAC_Name_1	
	Quantitative_ratio_1	
	Qualitative_Prevalence_1	
	Prevalence_Category_1	

^a Entries with the value “nul” in any column indicate that it was not possible to extract sufficiently specific information from the publication.

Frequency Distribution of Prototropic, Ring $\rightleftharpoons$ Chain, and Valence Tautomerism Rules for Single-Rule Transformations and Those with Combined or Alternative Rules^{a,b,c}

Type	Standard Rules			
	Rule number	Rule name	Single rule	Combined or alternative rule
Prototropic Rules	PT_02_00	1,5 (thio)keto/(thio)enol	0	230
	PT_03_00	simple (aliphatic) imine	0	323
	PT_04_00	special imine	0	127
	PT_05_00	1,3 aromatic heteroatom H-shift	0	184
	PT_06_00	1,3 heteroatom H-shift	708	891
	PT_07_00	1,5 (aromatic) heteroatom H-shift (1)	391	463
	PT_08_00	1,5 (aromatic) heteroatom H-shift (2)	0	88
	PT_09_00	1,7 (aromatic) heteroatom H-shift	89	256
	PT_10_00	1,9 (aromatic) heteroatom H-shift	0	72
	PT_11_00 ^b	1,11 (aromatic) heteroatom H-shift	0	33
Ring-Chain Rules	PT_12_00	1,3 furanones	0	84
	PT_16_00	nitroso/oxime	0	14
	RC_03_00	5_exo_trig	0	50
	RC_03_01	5_exo_trig	0	50
	RC_03_02	5_exo_trig	19	0
	RC_04_01	6_exo_trig	0	49
	RC_04_02	6_exo_trig	0	49
	RC_09_00	5_endo_trig	67	0
	RC_10_00	6_endo_trig	10	15
	RC_10_01	6_endo_trig	29	15
Type	New Rules			
	Rule number	Rule name	Single rule	Combined or alternative rule
Prototropic Rules	PT_22_00	imine/imine	3	0
	PT_23_00	1,5 furanones	12	0
	PT_24_00	1,4 N-oxide/N-hydroxide	8	0
	PT_25_00	1,6 N-oxide/N-hydroxide (1)	4	0

Type	Standard Rules			Combined or alternative rule
	Rule number	Rule name	Single rule	
	PT_26_00	1,6 N-oxide/N-hydroxide (2)	5	0
	PT_27_00	acene	13	0
	PT_27_01	thiophene analogue of acene	15	
	PT_28_00	nitro/aci-nitro via aromatic ring (1): 1,7 H-shift	2	0
	PT_29_00	nitro/aci-nitro via aromatic ring (1): 1,5 H-shift	3	0
	PT_29_01	o-tolualdehyde	2	0
	PT_30_00	nitramide/N-nitronic acid	1	0
	PT_31_00	sulfone-based aliphatic compounds	1	0
	PT_32_00	nitrile/ketenimine: 1,3 H-shift	8	0
	PT_33_00	nitrile/ketenimine: 1,5 H-shift	8	0
	PT_34_00	triad phosphorus-carbon	5	0
	PT_35_00	sulfonyl/sulfinyl: 1,2 H-shift	2	0
	PT_36_00	oxime/nitron: 1,2 H-shift	5	0
	PT_37_00	sulfonyl/S-oxide: 1,4 H-shift	1	0
	PT_38_00	silane-hemiaminal/silanoic amide	2	0
	PT_39_00	nitron/azoxy or Behrend rearrangement	19	0
	PT_40_00	tetrad phosphorus-carbon	1	0
	PT_41_00	pyridine 1-oxide/1-hydroxypyridine	2	0
	PT_42_00	³ / ₄ -pyrro(thio/seleno)lin-2-one	27	0
	PT_43_00	isobenzofuran/phthalan	4	0
	PT_44_00	2-substituted-pyrrole	6	0
	PT_45_00	isopropylidenecycloalkane/isopropylcycloalkene	17	0
	PT_46_00	4-picoline	1	0
	PT_47_00	isoindole/isoindolenine	24	0
	PT_48_00	benzofuranone	4	0
	PT_49_00	N-hydroxyindole	6	0
Ring-Chain Rules	RC_03_03	boronic acid/oxaborole	19	0
	RC_03_04	5_exo_trig	15	0
	RC_04_04	6_exo_trig	25	0
	RC_12_00	5_endo_tet or iminophosphorane/benzoxazaphospholine	39	0

Type	Standard Rules		
	Rule number	Rule name	Single rule Combined or alternative rule
Valence Rules	RC_13_00	6_endo_dig	1 0
	RC_14_00	thiadiazoline rearrangement	9 0
	RC_15_00	5_exo_trig: 1,4 H-shift	3 0
	RC_16_00	boryl/borate	2 0
	RC_17_00	boryl/borate: ion-complex	2 0
	RC_18_00	5_exo_tet or hydroxyphosphorane	4 0
	RC_19_00	nitroolfin/1,2-oxazine N-oxide	6 0
	RC_20_00	5_endo_trig: 1,4 H-shift or aminoethyl nitron/imidazolidin-1-ol	6 0
	RC_21_00	cyclobutane/enamine	3 0
	RC_22_00	5_endo_trig: 1,5 H-shift	12 0
	RC_23_00	6_endo_trig: 1,4 H-shift	1 0
	RC_24_00	λ^5 -/ λ^3 -phosphane	2 0
	VT_01_00	monothio-o-benzoquinone/benzoxathiete	2 0
	VT_01_01	α -dithione/1,2-dithiete	12 0
	VT_02_00	tetrazole/azide	84 0
	VT_03_00	isothiocyanate/triazinethione	8 0
	VT_04_00	tetrazine/azodiazole	21 0
	VT_05_00	1,2,3-triazole/diazoamide	8 0
	VT_06_00	norcaradiene/cycloheptatriene or benzene-oxide/oxepin	18 0
	VT_07_00	phospha-minephosphones	11 0
	VT_08_00	1,2,3,4-tetrazinium/azodiazonium	15 0
	VT_09_00	diazaphosphazole/phosphinoimine	25 0
	VT_10_00	phosphine/phosphonium salt	25 0

^aDifferent classes of tautomerism are defined by prefixing each rule with PT, RC, or VT for prototropic tautomerism, ring=chain tautomerism, and valence tautomerism, respectively. The second placeholder in the rule name between the underscores indicates the rule number in that category (i.e., "02" in PT_02_00), and the last number in the name indicates a variant of that rule (i.e., "01" in VT_01_01, "03" in RC_03_03). A rule ending with "_00" occurs only in one variant for that rule. This naming scheme allows us to add more variants in that rule if it is required in the future.

^bWe also have four variants of PT_11_00 for long-range hydrogen migration, where *mm* ranges from 01 to 04.

^cSMIRKS of these tautomeric rules are given in Spreadsheet S3 of the Supporting Information.

Table 3.Types of Hydrogen Shifts Observed in the One-Step and Two-Step Hydrogen Migrations^a

One-step hydrogen migrations ^a	Count	Two-step hydrogen migrations ^b	Count
1,2	7	1,3 > 1,3	135
1,3	1120	1,3 > 1,7	91
1,4	12	1,5 > 1,5	42
1,5	707	1,5 > 1,11	27
1,6	15	1,3 > 1,5	5
1,7	91	1,5 > 1,7	3
1,3/1,5	30	1,5 > 1,3	2
1,3/1,7	118	1,7 > 1,7	1
1,5/1,9	15	Others	48
1,5/1,7	2	–	–
Others	609	–	–

^a“/” indicates alternative H-shifts possible for the same trans- formation.^b“>” denotes that a first H-shift is followed by a second one to achieve the transformation.

Table 4.

Types of Tautomeric Equilibria in Tautomeric Pairs with Their Occurrences^d

Type of tautomerism	Count
Azo $\rightleftharpoons$ Hydrazone	333
Ring $\rightleftharpoons$ Chain ^a	318
Enol $\rightleftharpoons$ Keto	138
Oxo-enamine $\rightleftharpoons$ Oxo-imine	113
Diketo $\rightleftharpoons$ Keto-enol	108
Enol-imine $\rightleftharpoons$ Oxo-enamine	104
Amine $\rightleftharpoons$ Imine	83
Keto-enethiol $\rightleftharpoons$ Thioketo-enol	82
Azide $\rightleftharpoons$ Tetrazole ^b	82
nul $\rightleftharpoons$ nul ^c	78
Ring $\rightleftharpoons$ Chain ^b (Valence)	77
Enamine $\rightleftharpoons$ Imine	72
Oxo-enamine $\rightleftharpoons$ Phenol-imine	65
Pyridol $\rightleftharpoons$ Pyridone	58
NH $\rightleftharpoons$ NH	57
Phenol-quinone $\rightleftharpoons$ Phenol-quinone	51
Enol-imine $\rightleftharpoons$ Oxo-imine	40
Benzoxazaphospholine $\rightleftharpoons$ Iminophosphorane ^a	39
CH $\rightleftharpoons$ NH	35
Keto-enol $\rightleftharpoons$ Keto-enol	33
NH-imidazole $\rightleftharpoons$ NH-imidazole	31
Lactam $\rightleftharpoons$ Lactim	31
Amine-imine $\rightleftharpoons$ Amine-imine	27
Cyclohexadienone $\rightleftharpoons$ Phenol	27
3H-2-one $\rightleftharpoons$ 5H-2-one	27
Enethiol $\rightleftharpoons$ Thioketo	26
N-hydroxide $\rightleftharpoons$ N-oxide	25

Type of tautomerism	Count
Diazaphosphazole $\rightleftharpoons$ Phosphinoimine ^b	25
Phosphine $\rightleftharpoons$ Phosphonium salt ^b	25
Isoindole $\rightleftharpoons$ Isoindolenine	24
1,4-Dihydro $\rightleftharpoons$ 1,6-Dihydro	19
Nitrone $\rightleftharpoons$ Nitrone	19
Isopropylcycloalkene $\rightleftharpoons$ Isopropylidenecycloalkane	17
Thioamide $\rightleftharpoons$ Thioimidol	16
Keteneimine $\rightleftharpoons$ Nitrile	16
Tropolone $\rightleftharpoons$ Tropolone	12
2H $\rightleftharpoons$ 6H	12
Amide $\rightleftharpoons$ Imidol	12
Amino $\rightleftharpoons$ Imino	12
Arene-imine $\rightleftharpoons$ Azepine ^b	12
Anaquinoid $\rightleftharpoons$ Paraquinonimine	10
NH $\rightleftharpoons$ OH	10
1,2-Dihydro $\rightleftharpoons$ 1,4-Dihydro	9
Carbamoylimino $\rightleftharpoons$ Guanidine ^a	9
Thiol $\rightleftharpoons$ Thione	9
Nitroso-enamine $\rightleftharpoons$ Oxim-imine	7
1,2-Dihydro $\rightleftharpoons$ 2,5-Dihydro	6
Cycloheptatriene $\rightleftharpoons$ Norcaradiene ^b	6
Pyrrole $\rightleftharpoons$ Pyrrolidine	6
1,4-Dihydro $\rightleftharpoons$ 4,6-Dihydro	5
Nitrone $\rightleftharpoons$ Oxime	5
Triazole $\rightleftharpoons$ Triazole	4
2H $\rightleftharpoons$ 4H	4
Enol-enamine $\rightleftharpoons$ Oxo-enamine	4
Amino-thieno $\rightleftharpoons$ Imine-thieno	4
Isobenzofuran $\rightleftharpoons$ Phthalan	4
CH $\rightleftharpoons$ OH	3

Type of tautomerism	Count
1,4-Dihydro $\rightleftharpoons$ 4,5-Dihydro	3
N(1)H $\rightleftharpoons$ N(3)H	3
Amine $\rightleftharpoons$ Zwitterion	3
Selenol $\rightleftharpoons$ Selone	3
Imine $\rightleftharpoons$ Imine	3
Nitro $\rightleftharpoons$ aci-Nitro	3
5,6-Dihydro $\rightleftharpoons$ 5,6-Dihydro	2
5,6-Dihydro-2H $\rightleftharpoons$ 5,6-Dihydro-4H	2
C1-H $\rightleftharpoons$ C3-H	2
Thiol $\rightleftharpoons$ Zwitterion	2
Sulfinyl $\rightleftharpoons$ Sulfinyl	2
Sila-hemiaminal $\rightleftharpoons$ Silanoic-amide	2
λ^3 -Phosphane $\rightleftharpoons$ λ^5 -Phosphane ^a	2
1H $\rightleftharpoons$ 2H	1
2H $\rightleftharpoons$ 2H	1
1,6-Dihydro $\rightleftharpoons$ 3,6-Dihydro	1
4H $\rightleftharpoons$ 6H	1
Nitroso-imine $\rightleftharpoons$ Oxim-imine	1
C3-H $\rightleftharpoons$ N(5)H	1
Oxo-thione $\rightleftharpoons$ nul ^c	1
Pyridol $\rightleftharpoons$ Zwitterion	1
1H $\rightleftharpoons$ 3H	1
N-nitronic acid $\rightleftharpoons$ Nitramide	1
Enol $\rightleftharpoons$ Ylide	1
S-oxide $\rightleftharpoons$ Sulfinyl	1

^aRing—chain tautomerism type (total count for ring—chain tautomerism of two tautomers including Benzoxazaphospholine $\rightleftharpoons$ Iminophosphorane, Carbamoylimino $\rightleftharpoons$ Guanidine, and λ^5 -Phosphane $\rightleftharpoons$ λ^3 -Phosphane pairs is 368).

^bValence tautomerism type.

^c“nul” indicates cases of tautomeric equilibria for which no name for one or the other or both tautomers was given in the references, and we were not able to assign any specific name.

^dExamples for each of these types are given in Table S1 of the Supporting Information.

Table 5.

Types of Tautomeric Equilibria in Three-Tautomer Sets with Their Occurrences

Type of tautomerism	Count
Enethiol $\rightleftharpoons$ Enethiol $\rightleftharpoons$ Thioketo	51
Enol-imine $\rightleftharpoons$ Oxo-enamine $\rightleftharpoons$ Oxo-imine	42
Phenol-quinone $\rightleftharpoons$ Phenol-quinone $\rightleftharpoons$ Phenol-quinone	25
CH $\rightleftharpoons$ NH $\rightleftharpoons$ OH	21
Chain $\rightleftharpoons$ Ring $\rightleftharpoons$ Ring	20
5-Hydroxytriazine $\rightleftharpoons$ Orthoquinonoid $\rightleftharpoons$ Paraquinonoid	11
Thioamide $\rightleftharpoons$ Thioimidol $\rightleftharpoons$ Thioimidol	11
nul $\rightleftharpoons$ nul $\rightleftharpoons$ nul ^a	9
Enol $\rightleftharpoons$ Enol $\rightleftharpoons$ Keto	6
Enol $\rightleftharpoons$ Keto $\rightleftharpoons$ Keto	6
Enamine $\rightleftharpoons$ Imine $\rightleftharpoons$ nul	6
Enamine $\rightleftharpoons$ Enamine $\rightleftharpoons$ Imine	6
1,2-Dihydro $\rightleftharpoons$ 1,4-Dihydro $\rightleftharpoons$ 1,5-Dihydro	5
1,7-Dihydro-7-oxo $\rightleftharpoons$ 4,7-Dihydro-7-oxo $\rightleftharpoons$ 7-Hydroxy	5
Nitroso-enamine $\rightleftharpoons$ Nitroso-imine $\rightleftharpoons$ Oxim-imine	4
Enol $\rightleftharpoons$ Keto $\rightleftharpoons$ Zwitterion	4
Triazole $\rightleftharpoons$ Triazole $\rightleftharpoons$ Triazole	4
Azo $\rightleftharpoons$ Hydrazone $\rightleftharpoons$ Zwitterion	3
Diketo $\rightleftharpoons$ Keto-enol $\rightleftharpoons$ Keto-enol	3
Others	8

^aSee Table 4.

Table 6.

Distribution of Some Common Tautomeric Pairs in Different Prevalence Categories

	Tautomer_1	Prevalence Category					Tautomer_2	Prevalence Category				
		0	1	2	3	4		0	1	2	3	4
Azo		39	73	99	79	43	Hydrazone	54	127	80	38	34
Enol		63	35	20	14	5	Keto	20	34	10	43	30
Oxo-enamine		9	2	37	44	21	Oxo-imine	25	70	7	2	9
Diketo		27	41	9	25	6	Keto-enol	6	37	22	39	4
Enol-imine		6	53	0	29	16	Oxo-enamine	16	29	2	51	6
Keto-enethiol		0	59	19	3	0	Thioketo-enol	0	4	19	58	0
Amine		10	32	5	28	8	Imine	18	33	5	24	3
Enamine		19	13	26	9	5	Imine	5	14	26	8	19
Oxo-enamine		16	15	34	0	0	Phenol-imine	0	0	34	15	16
Pyridol		3	38	8	8	0	Pyridone	2	15	12	27	2
Enethiol		0	6	3	1	16	Thioketo	18	1	4	3	0
Enol-imine		6	53	0	29	16	Oxo-enamine	16	29	2	51	6
Lactam		10	1	4	15	1	Lactim	4	11	5	9	1
5H-2-one		2	4	2	13	6	3H-2-one	6	13	2	4	2
Cyclohexadienone		9	1	5	4	8	Phenol	9	3	5	0	10
Isoindole		1	8	5	10	0	Isoindolenine	0	10	5	8	1
N-hydroxide		1	3	5	9	1	N-oxide	1	10	5	2	1
Keteneimine		5	11	0	0	0	Nitrile	0	6	0	9	1
Ring ^a		28	125	114	64	7	Chain ^a	11	65	171	65	26
Benzoxazaphospholine ^a		0	11	17	10	1	Iminophosphorane ^a	1	10	17	11	0
Diazaphosphazole ^b		2	4	1	17	1	Phosphinoimine ^b	1	17	1	4	2
Phosphine ^b		3	9	5	8	0	phosphonium salt ^b	4	4	5	12	0
Ring ^b		6	22	8	26	15	Chain ^b	18	28	8	17	6
Tetrazole ^b		5	29	8	35	5	Azide ^b	4	70	1	3	4

^aRing-chain tautomerism type.

$b_{\text{Valence tautomerism type.}}$

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript