Published in final edited form as:

Insect Mol Biol. 2019 February ; 28(1): 65–73. doi:10.1111/imb.12528.

Site-specific transgenesis of the *D. melanogaster* Y-chromosome using CRISPR/Cas9

Anna Buchman¹, Omar S. Akbari^{1,2,†}

¹Section of Cell and Developmental Biology, University of California, San Diego, La Jolla, California, 92093, United States of America.

²Tata Institute for Genetics and Society, University of California, San Diego, La Jolla, CA 92093, United States of America.

Abstract

Despite the importance of Y-chromosomes in evolution and sex determination, their heterochromatic, repeat-rich nature makes them difficult to sequence (due, in part, to ambiguities in sequence alignment and assembly) and to genetically manipulate, and therefore they generally remain poorly understood. For example, the *D. melanogaster* Y-chromosome, one of the most extensively studied Y-chromosomes, is widely heterochromatic and composed mainly of highly repetitive sequences, with only a handful of expressed genes scattered throughout its length. Efforts to insert transgenes on this chromosome have thus far relied on either random insertion of transposons (sometimes harboring ‘landing sites’ for subsequent integrations) with limited success or on chromosomal translocations, thereby limiting the types of Y-chromosome related questions that could be explored. Here we describe a versatile approach to site-specifically insert transgenes on the Y-chromosome in *D. melanogaster* via CRISPR/Cas9-mediated Homology Directed Repair (HDR). We demonstrate the ability to insert, and detect expression from, fluorescently marked transgenes at two specific locations on the Y-chromosome, and we utilize these marked Y-chromosomes to detect and quantify rare chromosomal nondisjunction effects. Finally, we discuss how this Y-docking technique could be adapted to other insects to aid in the development of genetic control technologies for the management of insect disease vectors and pests.

Keywords

CRISPR; Cas9; Y-Chromosome; *Drosophila*; transgenesis; HDR

[†]To whom all correspondence should be addressed: Omar S. Akbari, Section of Cell and Developmental Biology, University of California, San Diego, La Jolla, CA 92093, USA, Ph: 858-246-0640, oakbari@ucsd.edu.

Author Contributions

O.S.A and A.B. conceived and designed experiments. A.B. performed molecular and genetic experiments. All authors contributed to the writing, analyzed the data, and approved the final manuscript.

The authors declare no conflict of interest.

Disclosure

The authors declare nothing to disclose.

Introduction

The ability to perform site-specific transgenesis in insects and other organisms has greatly improved researchers' ability to conduct controlled experiments in a precise way. In *D. melanogaster*, for example, prior to the advent of site-specific transgene integration technologies, most transgenesis was carried out using transposon-based vectors that inserted embedded transgenes randomly throughout the genome, leaving them vulnerable to distinct position effects arising from surrounding *cis*-regulatory regions and chromatin structure (Wimmer 2005). The development of site-specific transgenesis techniques such as those that rely on the phage ϕ C31-derived site-specific integrase (Groth et al. 2004) or on recombinase-mediated cassette exchange (RMCE, (Oberstein et al. 2005)), represented a major advance (Wimmer 2005; Venken and Bellen 2005). This is because these techniques enabled more accurate and detailed transgene comparison and identification of insertion loci that are conducive to robust transgene expression levels and are fitness neutral, factors that can be especially important in practical applications of insect transgenesis such as vector control (Irvin et al. 2004; Wimmer 2005). However, the widely adopted site-specific recombinase-based transgenesis techniques (e.g., ϕ C31) are limited by their requirement for pre-existing 'landing sites' within genomes of interest, which must first be generated by random transposon-mediated transgenesis, and unfortunately these sites are lacking in most non-model organisms (Wimmer 2005).

The arrival of CRISPR technologies heralded a new era not only for traditional genome manipulation such as mutagenesis, but also for precise, site-specific transgenesis (Gratz et al. 2015; Bier et al. 2018). By using a simplified two-component system consisting of a *S. pyogenes* Cas9 endonuclease (SpCas9) and a single chimeric guide RNA (sgRNA, (Jinek et al. 2012)), one can generate DNA double-strand breaks (DSB) in a location of one's choosing, provided it contains a 3-bp (NGG) protospacer adjacent motif (PAM) required for CRISPR/Cas9 function (Jinek et al. 2012). Then, as long as a donor template comprising homology-containing stretches (i.e., homology arms) flanking the target transgene is provided, the DSB break can be repaired via the cell's own homology-directed repair (HDR) pathway to integrate the transgene at the precise location of the DSB (Gratz et al. 2015). This technique has been shown to be highly efficient in *D. melanogaster* (Gokcezade, Sienski, and Duchek 2014; Gratz et al. 2014, 2015), and given CRISPR's functionality in many insects (Gantz et al. 2015; Li, Bui, et al. 2017; Hammond et al. 2016; Dong et al. 2018; Sun et al. 2017; Li, Au, et al. 2017; Kohno et al. 2016; Li, Bui, and Akbari 2017; Li, Akbari, and White 2018), should be broadly applicable for insect transgenesis in general (e.g., (Li, Bui, et al. 2017; Hammond et al. 2016; Gantz et al. 2015)).

The Y-chromosome of *D. melanogaster*, and of most other heteromorphic sex chromosome-bearing organisms, has so far presented an elusive target for site-specific transgenesis applications. The Y-chromosome's degenerative, repetitive nature renders it difficult to sequence, assemble, and analyze genomically (Carvalho 2002; Piergentili 2010; Hall et al. 2016); additionally, it is almost entirely heterochromatic (Elgin and Reuter 2013), which likely makes it refractory to transgene integration and/or robust transgene expression (Bernardini et al. 2014). These contributing factors have made it difficult to manipulate the Y-chromosome in a precise fashion, and the only successful attempts at placing transgenes

on the Y-chromosome in insects, for example, have relied on random (and rare) events, such as chromosomal rearrangements or low frequency Y-specific transposon-mediated integrations, and subsequent manipulations (Starz-Gaiano et al. 2001; Janos Szabad, Bellen, and Venken 2012; Bernardini et al. 2014, 2017; Zhang and Stankiewicz 1998; Zhang and Spradling 1994).

Despite the challenges presented by the nature of the Y-chromosome, having the ability to engineer it in *D. melanogaster* and other insects would be of utility in a number of different applications. Firstly, the ability to mark the Y-chromosome with an easily-scorable, specific marker may in itself be useful. For example, it would allow for the facile identification of rare Y-bearing or Y-lacking individuals (e.g., XXY females and XO males arising from nondisjunction, Figure 2A; (Calvin B. Bridges 1913; C. B. Bridges 1916)), which can be of experimental use (e.g., (Brown and Bachtrog 2017; Lemos, Branco, and Hartl 2010; Hearn et al. 1991; Yang, Baxter, and Van Doren 2012). A marked Y-chromosome would also enable interspecific Y-chromosome introgression studies, which can shed light on Y-chromosome evolution and function (Araripe, Tao, and Lemos 2016; Sackton et al. 2011; Bernardini et al. 2017), and can also be useful for sexing embryos (Bernardini et al. 2014; Condon et al. 2007), although it would also mark XXY females and fail to identify XO males, which may be problematic for certain applications.

Additionally, the ability to express specific transgenes directly from the Y-chromosome would be useful in several different contexts. It would enable general studies of Y-chromosome gene function, for example, as well as analyses of chromosome loss and interaction (Janos Szabad, Bellen, and Venken 2012; Soós and Szabad 2014; J. Szabad and Würgler 1987). Perhaps more interestingly, it could aid in the development of genetic control mechanisms for insect vectors and pests, which offer promising solutions to significant health and agricultural problems (Champer, Buchman, and Akbari 2016; Esvelt et al. 2014; Sinkins and Gould 2006). For example, distortion of the sex ratio in favor of males can lead to a gradual population reduction and eventual elimination of a target population (Gould and Schliekelman 2004; Hamilton 1967; Hickey and Craig 1966; Papathanos, Windbichler, and Akbari, n.d.), and natural so-called meiotic driving Y-chromosomes have been described (Wood and Newton 1991; Newton, Wood, and Southern 1976; Sweeny and Barr 1978). A system for sex-ratio distortion can also be engineered by designing transgenes that target the X-chromosome during spermatogenesis, and such an X-shredder element has already been developed in one species of mosquito (Windbichler, Papathanos, and Crisanti 2008; Galizi et al. 2014, 2016). However, to be maximally effective and self-perpetuating, such an element must be linked to the Y-chromosome (Deredec, Burt, and Godfray 2008; Deredec, Godfray, and Burt 2011; Galizi et al. 2016), a feature that the developed systems lacked but that could be achieved by an efficient method of targeted Y-chromosome transgenesis.

To overcome this technological gap, here we describe the development of a CRISPR/Cas9-based technique for site-specific engineering of the *D. melanogaster* Y-chromosome. Specifically, we demonstrate the ability to insert a fluorescently marked transgenic cassette at specific locations on the Y-chromosome, and to utilize the marked Y-chromosome to

identify XXY females and XO males. The interspecies portability and limitations of this technique are also discussed.

Results & Discussion

To generate transgenic elements that could be inserted in specific locations on the *D. melanogaster* Y-chromosome with CRISPR/Cas9-mediated HDR, we engineered a vector comprising a fluorescent marker (tdTomato) driven by the eye-specific 3xP3 promoter (Berghammer, Klingler, and Wimmer 1999) and flanked by the *gypsy* (Holdridge and Dorsett 1991) and CTCF (Kyrchanova et al. 2008) insulators, with unique restriction sites upstream and downstream for cloning specific homology arms (Figure 1). We then selected ten distinct intergenic regions spanning the Y-chromosome for targeting (Figure 1A), reasoning that attempting transgene insertion at multiple different sites within gene-empty regions would increase our chances of a successful detectable integration. We identified a suitable sgRNA target site with no predicted off-target effects in each region, and cloned in homology arms, corresponding to ~800–1,000 base pairs of sequence 5' and 3' of each selected target site, upstream and downstream of the insulator-flanked 3xP3-tdTomato element to generate ten unique Y-chromosome targeting transgenes (AByA-J) (Figure 1A).

Each transgene was then injected, along with the appropriate *in vitro* transcribed sgRNA and Cas9 protein, into a transgenic line expressing a germline source of Cas9 using standard procedures (Gratz et al. 2014), and G₁ progeny were screened for presence of eye-specific tdTomato fluorescence (Figure 1B). Of the ten distinct construct injections, only two (AByF and AByG) out of ten yielded transgenic male individuals, despite the large number (>200 in most cases, 2142 total for all constructs) of G₀ embryos injected for each construct (Figure 1C-D). This observation suggests that either the rate of Y-chromosome transgenesis is not particularly high (especially when compared to rates reported anecdotally and in other studies utilizing CRISPR/Cas9-based HDR in autosomal locations, e.g., (Gratz et al. 2014)), or that a number of tested Y-chromosome target sites were not located in regions conducive to somatic gene expression (although a low transgenesis rate not related to the Y-chromosome cannot be ruled out, and the fact that only injected male G₀ embryos can give rise to transgenics has to be considered). Neither of these explanations would be surprising, given the heterochromatic nature of the *D. melanogaster* Y-chromosome (Carvalho 2002), and simply suggests that several target regions must be tested in order to identify those suitable for HDR and/or that allow robust expression. Following their recovery, we outcrossed transgenic males to white eye (w⁻, w[1118]) females for several generations to confirm paternal inheritance of the tdTomato marker, and saw expected inheritance patterns (i.e., all male progeny and no female progeny of a male inherited the fluorescent marker). We also verified that the transgene was correctly inserted by performing PCR across the transgene-insertion junction on genomic DNA of transgenic males and sequencing the products.

We then set out to determine whether a marked Y-chromosome could be useful in identifying the rare progeny resulting from meiotic sex chromosome nondisjunction events in females (Figure 2A-B), as a clear means for identification of such individuals could be useful both for those that want to avoid them and those that wish to utilize them for

addressing various research questions (e.g., (Brown and Bachtrog 2017; O'Tousa 1982; Gilliland et al. 2014)). To do this, we crossed males from the stronger-expressing of our two transgenic Y-chromosome lines, AByF, to a variety of X-chromosome balancer line virgins. These balancer lines contain genetic elements such as multiple inversions, rearrangements, and deficiencies that are known to increase the probability of nondisjunction by causing abnormalities in recombination as well as unusual chromosomal pairing and segregation patterns (Xiang and Hawley 2006; Gilliland et al. 2014; Ramel 2009; Forbes 1962; Ramel 1962). For four of these crosses (all involving balancer lines with X-chromosome deficiencies), we identified progeny that appeared to be the result of meiotic nondisjunction events in the X-chromosome balancer females - XXY-bearing females (identified by presence of the fluorescent marker) and XO males (identified by the absence of the fluorescent marker) - at various frequencies ranging from 0–27% (Figure 2C). These aberrant progeny were individually outcrossed to a common laboratory stock (w[1118]) to test for fertility. All putative XO males were sterile (n=36) as expected (Griffiths 2005). All XXY females were fertile (n=28), and gave rise to progeny classes that indicated occurrence of secondary nondisjunction in XXY parents, which is in agreement with previous observations (C. B. Bridges 1916; Xiang and Hawley 2006; Carpenter 1973). The presence or absence of the transgenic Y-chromosome in XXY females and XO males, respectively, was verified with PCR as described above. Given the high rates of nondisjunction individuals obtained by crossing the marked Y-chromosome line with certain X-chromosome balancers (e.g., balancer line 579), future experiments requiring large quantities of XXY females and XO males could be conducted using this approach, which could accelerate basic research on deciphering key biological processes and gene networks involved in Y-chromosome functionality and evolution.

In short, here we describe the engineering of two Y-chromosome marked transgenic *D. melanogaster* lines using site-specific, CRISPR/Cas9-mediated insertion. This may be useful for generating, identifying, and tracking progeny arising from rare meiotic nondisjunction events in a more straightforward manner than is typically used (Brown and Bachtrog 2017; Piergentili 2010; Hess and Meyer 1968). This may also be useful for detecting *in vivo* mosaic Y-chromosome loss (Janos Szabad, Bellen, and Venken 2012), although we did not observe evidence for such loss in assayed transgenic XY or XXY individuals. More broadly, however, we provide a proof of principle example of a CRISPR/Cas-based strategy for docking transgenes site-specifically on the Y-chromosome that should in principle be applicable to many other species. Although the often-heterochromatic nature of a Y-chromosome may, as our observations possibly suggest, reduce the number of targeted insertion sites that are amenable to modification and/or robust transgene expression, the ease with which distinct Y-chromosome targeting transgene cassettes can be generated should allow for testing of many possible insertion sites, greatly increasing the chances of successful transgene integration with desired expression. The ability to mark, and express transgenes directly from, the Y-chromosome not only allows for facile identification of Y-chromosome bearing individuals at various life stages and enables researchers to probe the enigmatic function of the Y-chromosome (e.g., (Brown and Bachtrog 2017; Bernardini et al. 2014), but could also help pave the way for engineering genetic insect vector and pest control strategies such as X-chromosome shredding (Hickey and Craig 1966; Hamilton

1967; Gould and Schliekelman 2004; Huang et al. 2007; Galizi et al. 2014; Champer, Buchman, and Akbari 2016; Galizi et al. 2016). This approach in particular depends on the destruction of X-bearing sperm to produce males that only give rise to male progeny (Huang et al. 2007; Champer, Buchman, and Akbari 2016), and requires the ability to meiotically express an X-chromosome targeting element from the Y-chromosome (Beaghton, Beaghton, and Burt 2016; Galizi et al. 2016). The strategy described here could be used to dock such an X-chromosome targeting transgene (perhaps one based on CRISPR/Cas9 technology, (Galizi et al. 2016; Zuo et al. 2017)) on the Y-chromosome of many insects, including the malaria vector *Anopheles gambiae*, to facilitate the engineering of the promising X-chromosome shredding genetic control strategy that can be used to combat malaria and other vectored pathogens (Gould and Schliekelman 2004; Hall et al. 2016; Papathanos, Windbichler, and Akbari, n.d.; Champer, Buchman, and Akbari 2016).

Experimental Procedures

Construct Design and Assembly

To generate vector ABy, the base vector used to generate vectors AByA-AByJ, several components were cloned into the *piggyBac* plasmid pBac[3xP3-DsRed] (Li, Bui, et al. 2017) using Gibson assembly/EA cloning (Gibson et al. 2009). First, a Gypsy insulator fragment amplified with primers ABy.1 and ABy.2 from *Drosophila* genomic DNA, the 3xP3 promoter amplified with primers ABy.3 and ABy.4 from plasmid pBac[3xP3-EGFP afm] (Horn and Wimmer 2000), and a *Drosophila* codon optimized tdTomato marker amplified with primers ABy.5 and ABy.6 from a gene synthesized vector (Genscript, Piscataway, NJ) were cloned into a BstBI/NotI digested pBac[3xP3-DsRed] backbone using EA cloning. The resulting plasmid was digested with AvrII, and the following components were cloned in via EA cloning: an *attP* sequence from plasmid M{3xP3-RFP attP} (Bischof et al. 2007) amplified with primers ABy1.7 and ABy.8, and a CTCF insulator fragment amplified with primers ABy.9 and ABy.10 from *Drosophila* genomic DNA. All primer sequences are listed in Supplementary Table 1. To generate the final vectors containing homology arms specific to each putative Y-chromosome docking site, vector ABy was first digested with PmeI and each 5' homology arm (amplified with primers specific to the Y docking site from *Drosophila* genomic DNA) was individually cloned in using EA cloning. Each resulting intermediate plasmid was then digested with EcoRI, and each corresponding 3' homology arm (amplified with primers specific to the Y docking site from *Drosophila* genomic DNA) was cloned in using EA cloning. The 10 Y-chromosome docking site-specific construct names and the primers used to amplify the 5' and 3' homology arms for each are listed in Supplementary Table 1. (The slight variation in the length of the homology arms was not expected, and not observed, to affect integration efficiency.) Vectors AByF and AByG are available from Addgene (#s 111084 and 111083, respectively). To generate double-stranded DNA breaks for vector incorporation, sgRNAs targeting 10 distinct regions of the Y-chromosome were *in vitro* transcribed with the MEGAscript™ T7 Transcription Kit (ThermoFisher Scientific, cat. # AM1334) using a self-annealing set of primers, with a unique forward primer for each target site and a universal reverse primer. Genomic locations of each sgRNA target sequence, the target sequences themselves, and primers used for *in vitro* transcription are listed in Supplementary Table 1. sgRNA target

sites with no predicted off-target effects were selected using flyCRISPR Target Finder (Gratz et al. 2014).

Fly Culture and Strains

Fly husbandry and crosses were performed under standard conditions at 25°C. Rainbow Transgenics (Camarillo, CA) carried out all of the fly injections. Each construct was pre-mixed with Cas9 protein (PNA Bio Inc., cat. # CP01–20) and *in vitro* transcribed sgRNAs at the following concentrations: 400ng/μL cyclic donor plasmid, 40ng/μL sgRNA, and 300ng/μL Cas9 protein. Constructs were injected into a *vasa*-Cas9 transgenic line marked with 3xP3-GFP (Bloomington Drosophila Stock Center #51324, w[1118]; PBac{y[+mDint2]=vas-Cas9}VK00027). Only transgenic males were recovered, and these were singly outcrossed to w- (w[1118]) virgin females to verify paternal transmission. Male progeny without the *vasa*-Cas9 transgene were selected and further crossed to w- to establish a stock. Stocks for the two transgenic lines recovered (AByF and AByG) were submitted to the Bloomington Drosophila Stock Center.

Molecular characterization of Y-chromosome lines

To confirm correct insertion of transgenes on the Y-chromosome, PCRs were carried out on genomic DNA from transgenic males (Figure 1). Briefly, genomic DNA was extracted from individual flies with the DNeasy Blood & Tissue kit (QIAGEN, cat. # 69504) following the manufacturer's protocol. PCR was carried out using standard procedures to amplify the junction of the transgenes and the surrounding genomic region (as well as the unmodified target genomic region as a negative control and a region of the X-chromosome as a positive control) using primers listed in Supplementary Table 1. The PCR program utilized was as follows: 98°C for 30 seconds; 35 cycles of 98°C for 10 seconds, 56°C for 20 seconds, and 72°C for 30 seconds; then 72°C for 10 minutes. PCR products were purified with the MinElute® PCR Purification Kit (QIAGEN, cat. #28004) according to the manufacturer's protocol and sequenced with Sanger sequencing (SourceBioScience) utilizing the same primers as used for the PCRs. Sequences were analyzed with DNASTar software.

Generation and characterization of XXY females and XO males

To determine whether a marked Y-chromosome could be utilized to identify individuals that result from meiotic nondisjunction events, a single male from the AByF transgenic line (which showed stronger fluorescent marker expression levels than line AByG transgenic males) was crossed to virgins from the following X-chromosome balancer stocks from the Bloomington Drosophila Stock Center (BSC): BSC #s 579, 723, 727, 939, 944, 946, 959, 976, 3347, 6002, 6007, 6019, 6219, 7200, and 27887. Each cross was set in triplicate. Progeny were screened to identify females with fluorescent red eyes (putative XXY individuals) and males without fluorescent red eyes (putative XO individuals). Frequency of putative XXY and XO individuals was calculated by dividing the total number of each individual type found for each type of cross by the total number of females or males, respectively (Figure 2C). Each XO individual was outcrossed to w- (w[1118]) virgins to test for fertility. To molecularly confirm the presence of the transgenic Y-chromosome in putative XXY females and the absence of any Y-chromosome in putative XO males, genomic DNA was extracted as above, and PCRs were performed as above with primers

utilized for the AByF 5' gDNA-transgene junction, the AByF 5' gDNA-transgene junction, the AByF wildtype locus, and the X-chromosome positive control locus (Supplementary Table 1).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported in part by funding from the California Cherry board, an NIH-K22 Career Transition award (5K22AI113060), an NIH Exploratory/Developmental Research Grant Award (1R21AI123937), and a Defense Advanced Research Project Agency (DARPA) Safe Genes Program Grant (HR0011-17-2- 0047) awarded to O.S.A. Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study. We thank Thom Kaufman (Indiana University Bloomington, Indiana) for useful discussions and for providing the BSC X-chromosome balancer lines.

References

- Araripe LO, Tao Y, and Lemos B. 2016. "Interspecific Y Chromosome Variation Is Sufficient to Rescue Hybrid Male Sterility and Is Influenced by the Grandparental Origin of the Chromosomes." *Heredity* 116 (6): 516–22. [PubMed: 26980343]
- Beaghton Andrea, Beaghton Pantelis John, and Burt Austin. 2016. "Gene Drive through a Landscape: Reaction–diffusion Models of Population Suppression and Elimination by a Sex Ratio Distorter." *Theoretical Population Biology* 108 (4): 51–69. [PubMed: 26704073]
- Berghammer Andreas J., Klingler Martin, and Wimmer Ernst A.. 1999. "Genetic Techniques: A Universal Marker for Transgenic Insects." *Nature* 402 (6760): 370–71. [PubMed: 10586872]
- Bernardini Federica, Galizi Roberto, Menichelli Miriam, Papathanos Philippos-Aris, Dritsou Vicky, Marois Eric, Crisanti Andrea, and Windbichler Nikolai. 2014. "Site-Specific Genetic Engineering of the Anopheles Gambiae Y Chromosome." *Proceedings of the National Academy of Sciences of the United States of America* 111 (21): 7600–7605. [PubMed: 24821795]
- Bernardini Federica, Galizi Roberto, Wunderlich Mariana, Taxiarchi Chrysanthi, Kranjc Nace, Kyrou Kyros, Hammond Andrew, et al.2017. "Cross-Species Y Chromosome Function Between Malaria Vectors of the Anopheles Gambiae Species Complex." *Genetics* 207 (2): 729–40. [PubMed: 28860320]
- Bier Ethan, Harrison Melissa M., O'Connor-Giles Kate M., and Wildonger Jill. 2018. "Advances in Engineering the Fly Genome with the CRISPR-Cas System." *Genetics* 208 (1): 1–18. [PubMed: 29301946]
- Bischof Johannes, Maeda Robert K., Hediger Monika, Karch François, and Basler Konrad. 2007. "An Optimized Transgenesis System for Drosophila Using Germ-Line-Specific phiC31 Integrases." *Proceedings of the National Academy of Sciences of the United States of America* 104 (9): 3312–17. [PubMed: 17360644]
- Bridges Calvin B.1913. "Non-Disjunction of the Sex Chromosomes of Drosophila." *The Journal of Experimental Zoology* 15 (4): 587–606.
- Bridges CB1916. "Non-Disjunction as Proof of the Chromosome Theory of Heredity (Concluded)." *Genetics* 1 (2): 107–63. [PubMed: 17245853]
- Brown Emily, and Bachtrog Doris. 2017. "The Drosophila Y Chromosome Affects Heterochromatin Integrity Genome-Wide." *bioRxiv* 10.1101/156000.
- Carpenter AT1973. "A Meiotic Mutant Defective in Distributive Disjunction in Drosophila Melanogaster." *Genetics* 73 (3): 393–428. [PubMed: 4633612]
- Carvalho A. Bernardo. 2002. "Origin and Evolution of the Drosophila Y Chromosome." *Current Opinion in Genetics & Development* 12 (6): 664–68. [PubMed: 12433579]

- Champer Jackson, Buchman Anna, and Akbari Omar S.. 2016. "Cheating Evolution: Engineering Gene Drives to Manipulate the Fate of Wild Populations." *Nature Reviews. Genetics* 17 (3): 146–59.
- Condon Kirsty C., Condon George C., Dafa'alla Tarig H., Fu Guoliang, Phillips Caroline E., Jin Li, Gong Peng, and Alphey Luke. 2007. "Genetic Sexing through the Use of Y-Linked Transgenes." *Insect Biochemistry and Molecular Biology* 37 (11): 1168–76. [PubMed: 17916503]
- Deredec Anne, Burt Austin, and Godfray HCJ. 2008. "The Population Genetics of Using Homing Endonuclease Genes in Vector and Pest Management." *Genetics* 179 (4): 2013–26. [PubMed: 18660532]
- Deredec Anne, Charles HGodfray J, and Burt Austin. 2011. "Requirements for Effective Malaria Control with Homing Endonuclease Genes." *Proceedings of the National Academy of Sciences of the United States of America* 108 (43): E874–80. [PubMed: 21976487]
- Dong Yuemei, Simões Maria L., Marois Eric, and Dimopoulos George. 2018. "CRISPR/Cas9 - Mediated Gene Knockout of Anopheles Gambiae FREP1 Suppresses Malaria Parasite Infection." *PLoS Pathogens* 14 (3): e1006898. [PubMed: 29518156]
- Elgin Sarah C. R., and Reuter Gunter. 2013. "Position-Effect Variegation, Heterochromatin Formation, and Gene Silencing in Drosophila." *Cold Spring Harbor Perspectives in Biology* 5 (8): a017780. [PubMed: 23906716]
- Esvelt Kevin M., Smidler Andrea L., Catteruccia Flaminia, and Church George M.. 2014. "Concerning RNA-Guided Gene Drives for the Alteration of Wild Populations." *eLife* 3 (7): 10.7554/eLife.03401.
- Forbes Clifford. 1962. "The Effect of Heterozygous Inversions on Primary Nondisjunction in Drosophila Melanogaster." *Genetics* 47 (10): 1301. [PubMed: 13958677]
- Galizi Roberto, Doyle Lindsey A., Menichelli Miriam, Bernardini Federica, Deredec Anne, Burt Austin, Stoddard Barry L., Windbichler Nikolai, and Crisanti Andrea. 2014. "A Synthetic Sex Ratio Distortion System for the Control of the Human Malaria Mosquito." *Nature Communications* 5 (6): 3977.
- Galizi Roberto, Hammond Andrew, Kyrou Kyros, Taxiarchi Chrysanthi, Bernardini Federica, O'Loughlin Samantha M., Papathanos Philippos-Aris, Nolan Tony, Windbichler Nikolai, and Crisanti Andrea. 2016. "A CRISPR-Cas9 Sex-Ratio Distortion System for Genetic Control." *Scientific Reports* 6 (8): 31139. [PubMed: 27484623]
- Gantz Valentino M., Jasinskiene Nijole, Tatarenkova Olga, Fazekas Aniko, Macias Vanessa M., Bier Ethan, and James Anthony A.. 2015. "Highly Efficient Cas9-Mediated Gene Drive for Population Modification of the Malaria Vector Mosquito Anopheles Stephensi." *Proceedings of the National Academy of Sciences of the United States of America* 112 (49): E6736–43. [PubMed: 26598698]
- Gibson Daniel G., Young Lei, Chuang Ray-Yuan, Craig Venter J, Hutchison Clyde A. 3rd, and Smith Hamilton O.. 2009. "Enzymatic Assembly of DNA Molecules up to Several Hundred Kilobases." *Nature Methods* 6 (5): 343–45. [PubMed: 19363495]
- Gilliland William D., Colwell Eileen M., Lane Fiona M., and Snouffer Ashley A.. 2014. "Behavior of Aberrant Chromosome Configurations in Drosophila Melanogaster Female Meiosis I." *G3* 5 (2): 175–82. [PubMed: 25491942]
- Gokcezade Joseph, Sienski Grzegorz, and Duchek Peter. 2014. "Efficient CRISPR/Cas9 Plasmids for Rapid and Versatile Genome Editing in Drosophila." *G3* 4 (11): 2279–82. [PubMed: 25236734]
- Gould Fred, and Schliekelman Paul. 2004. "Population Genetics of Autocidal Control and Strain Replacement." *Annual Review of Entomology* 49: 193–217.
- Gratz Scott J., Dustin Rubinstein C, Harrison Melissa M., Wildonger Jill, and O'Connor-Giles Kate M.. 2015. "CRISPR-Cas9 Genome Editing in Drosophila." *Current Protocols in Molecular Biology* / Edited by Ausubel Frederick M. ... [et al.] 111 (7): 31.2.1–20.
- Gratz Scott J., Ukken Fiona P., Dustin Rubinstein C, Thiede Gene, Donohue Laura K., Cummings Alexander M., and O'Connor-Giles Kate M.. 2014. "Highly Specific and Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in Drosophila." *Genetics* 196 (4): 961–71. [PubMed: 24478335]
- Griffiths Anthony J. F. 2005. *An Introduction to Genetic Analysis* Macmillan.

- Groth Amy C., Fish Matthew, Nusse Roel, and Calos Michele P. 2004. "Construction of Transgenic *Drosophila* by Using the Site-Specific Integrase From Phage ϕ C31." *Genetics* 166 (4): 1775–82. [PubMed: 15126397]
- Hall Andrew Brantley, Papathanos Philippos-Aris, Sharma Atashi, Cheng Changde, Akbari Omar S., Assour Lauren, Bergman Nicholas H., et al. 2016. "Radical Remodeling of the Y Chromosome in a Recent Radiation of Malaria Mosquitoes." *Proceedings of the National Academy of Sciences of the United States of America* 113 (15): E2114–23. [PubMed: 27035980]
- Hamilton WD 1967. "Extraordinary Sex Ratios. A Sex-Ratio Theory for Sex Linkage and Inbreeding Has New Implications in Cytogenetics and Entomology." *Science* 156 (3774): 477–88. [PubMed: 6021675]
- Hammond Andrew, Galizi Roberto, Kyrou Kyros, Simoni Alekos, Siniscalchi Carla, Katsanos Dimitris, Gribble Matthew, et al. 2016. "A CRISPR-Cas9 Gene Drive System Targeting Female Reproduction in the Malaria Mosquito Vector *Anopheles Gambiae*." *Nature Biotechnology* 34 (1): 78–83.
- Hearn MG, Hedrick A, Grigliatti TA, and Wakimoto BT. 1991. "The Effect of Modifiers of Position-Effect Variegation on the Variegation of Heterochromatic Genes of *Drosophila Melanogaster*." *Genetics* 128 (4): 785–97. [PubMed: 1916244]
- Hess O, and Meyer GF. 1968. "Genetic Activities of the Y Chromosome in *Drosophila* during Spermatogenesis." *Advances in Genetics* 14: 171–223. [PubMed: 4884781]
- Hickey WA, and Craig GB Jr. 1966. "Genetic Distortion of Sex Ratio in a Mosquito, *Aedes Aegypti*." *Genetics* 53 (6): 1177–96. [PubMed: 5958915]
- Holdridge C, and Dorsett D. 1991. "Repression of hsp70 Heat Shock Gene Transcription by the Suppressor of Hairy-Wing Protein of *Drosophila Melanogaster*." *Molecular and Cellular Biology* 11 (4): 1894–1900. [PubMed: 1900919]
- Horn C, and Wimmer EA. 2000. "A Versatile Vector Set for Animal Transgenesis." *Development Genes and Evolution* 210 (12): 630–37. [PubMed: 11151300]
- Huang Yunxin, Magori Krisztian, Lloyd Alun L., and Gould Fred. 2007. "Introducing Desirable Transgenes into Insect Populations Using Y-Linked Meiotic Drive - a Theoretical Assessment." *Evolution; International Journal of Organic Evolution* 61 (4): 717–26. [PubMed: 17439607]
- Irvin Nic, Hoddle Mark S., O'Brochta David A., Carey Bryan, and Atkinson Peter W.. 2004. "Assessing Fitness Costs for Transgenic *Aedes Aegypti* Expressing the GFP Marker and Transposase Genes." *Proceedings of the National Academy of Sciences of the United States of America* 101 (3): 891–96. [PubMed: 14711992]
- Jinek Martin, Chylinski Krzysztof, Fonfara Ines, Hauer Michael, Doudna Jennifer A., and Charpentier Emmanuelle. 2012. "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity." *Science* 337 (6096): 816–21. [PubMed: 22745249]
- Kohno Hiroki, Suenami Shota, Takeuchi Hideaki, Sasaki Tetsuhiko, and Kubo Takeo. 2016. "Production of Knockout Mutants by CRISPR/Cas9 in the European Honeybee, *Apis Mellifera* L." *Zoological Science* 33 (5): 505–12. [PubMed: 27715425]
- Kyrchanova Olga, Toshchakov Stepan, Podstreshnaya Yulia, Parshikov Alexander, and Georgiev Pavel. 2008. "Functional Interaction between the Fab-7 and Fab-8 Boundaries and the Upstream Promoter Region in the *Drosophila* Abd-B Gene." *Molecular and Cellular Biology* 28 (12): 4188–95. [PubMed: 18426914]
- Lemos Bernardo, Branco Alan T., and Hartl Daniel L.. 2010. "Epigenetic Effects of Polymorphic Y Chromosomes Modulate Chromatin Components, Immune Response, and Sexual Conflict." *Proceedings of the National Academy of Sciences of the United States of America* 107 (36): 15826–31. [PubMed: 20798037]
- Li Ming, Akbari Omar S., and White Bradley J.. 2018. "Highly Efficient Site-Specific Mutagenesis in Malaria Mosquitoes Using CRISPR." *G3* 8 (2): 653–58. [PubMed: 29233915]
- Li Ming, Au Lauren Yun Cook, Douglass Deema, Chong Abigail, White Bradley J., Ferree Patrick M., and Akbari Omar S.. 2017. "Generation of Heritable Germline Mutations in the Jewel Wasp *Nasonia Vitripennis* Using CRISPR/Cas9." *Scientific Reports* 7 (1): 901. [PubMed: 28424460]

- Li Ming, Bui Michelle, and Akbari Omar S.. 2017. "Embryo Microinjection and Transplantation Technique for *Nasonia Vitripennis* Genome Manipulation." *Journal of Visualized Experiments: JoVE*, no. 130 (12). 10.3791/56990.
- Li Ming, Bui Michelle, Yang Ting, Bowman Christian S., White Bradley J., and Akbari Omar S.. 2017. "Germline Cas9 Expression Yields Highly Efficient Genome Engineering in a Major Worldwide Disease vector, *Aedes Aegypti*." *Proceedings of the National Academy of Sciences of the United States of America* 114 (49): E10540–49. [PubMed: 29138316]
- Newton ME, Wood RJ, and Southern DI. 1976. "A Cytogenetic Analysis of Meiotic Drive in the Mosquito, *Aedes Aegypti* (L.)." *Genetica* 46 (3): 297–318.
- Oberstein Adam, Pare Adam, Kaplan Leah, and Small Stephen. 2005. "Site-Specific Transgenesis by Cre-Mediated Recombination in *Drosophila*." *Nature Methods* 2 (8): 583–85. [PubMed: 16094382]
- O'Tousa J1982. "Meiotic Chromosome Behavior Influenced by Mutation-Altered Disjunction in *Drosophila Melanogaster* Females." *Genetics* 102 (3): 503–24. [PubMed: 6816677]
- Papathanos PA, Windbichler N, and Akbari OS. n.d. "Sex Ratio Manipulation for Insect Population Control." In *Transgenic Insects: Techniques and Applications*, 83–100.
- Piergentili Roberto. 2010. "Multiple Roles of the Y Chromosome in the Biology of *Drosophila Melanogaster*." *TheScientificWorldJournal* 10 (9): 1749–67.
- Ramel Claes. 1962. "INTERCHROMOSOMAL EFFECTS OF INVERSIONS IN *DROSOPHILA MELANOGASTER*: I. CROSSING-OVER." *Hereditas* 48 (1–2): 1–58.
- Ramel Claes. 2009. "THE INTERCHROMOSOMAL EFFECT OF INVERSIONS ON CROSSING-OVER IN RELATION TO NON HOMOLOGOUS PAIRING IN *DROSOPHILA MELANOGASTER*." *Hereditas* 54 (3): 293–306.
- Sackton Timothy B., Montenegro Horacio, Hartl Daniel L., and Lemos Bernardo. 2011. "Interspecific Y Chromosome Introgressions Disrupt Testis-Specific Gene Expression and Male Reproductive Phenotypes in *Drosophila*." *Proceedings of the National Academy of Sciences of the United States of America* 108 (41): 17046–51. [PubMed: 21969588]
- Sinkins Steven P., and Gould Fred. 2006. "Gene Drive Systems for Insect Disease Vectors." *Nature Reviews. Genetics* 7 (6): 427–35.
- Soós István, and Szabad János. 2014. "Assaying Benzene, a Parquet Varnish, and a Synthetic Thinner with Respect to Induction of in Vivo Chromosome Loss in Wing Primordial Cells of *Drosophila*." *Mutation Research. Genetic Toxicology and Environmental Mutagenesis* 763 (3): 18–22. [PubMed: 24530351]
- Starz-Gaiano M, Cho NK, Forbes A, and Lehmann R. 2001. "Spatially Restricted Activity of a *Drosophila* Lipid Phosphatase Guides Migrating Germ Cells." *Development* 128 (6): 983–91. [PubMed: 11222152]
- Sun Dan, Guo Zhaojiang, Liu Yong, and Zhang Youjun. 2017. "Progress and Prospects of CRISPR/Cas Systems in Insects and Other Arthropods." *Frontiers in Physiology* 8 (9): 608. [PubMed: 28932198]
- Sweeny TL, and Barr AR. 1978. "Sex Ratio Distortion Caused by Meiotic Drive in a Mosquito, *Culex Pipiens* L." *Genetics* 88 (3): 427–46. [PubMed: 17248804]
- Szabad Janos, Bellen Hugo J., and Venken Koen J. T.. 2012. "An Assay to Detect in Vivo Y Chromosome Loss in *Drosophila* Wing Disc Cells." *G3* 2 (9): 1095–1102. [PubMed: 22973547]
- Szabad J, and Würzler FE. 1987. "A Genetic Assay to Detect Chromosome Gain And/or Loss in Somatic Cells of *Drosophila Melanogaster*." *Mutation Research* 180 (2): 201–6. [PubMed: 3116420]
- Venken Koen J. T., and Bellen Hugo J.. 2005. "Emerging Technologies for Gene Manipulation in *Drosophila Melanogaster*." *Nature Reviews. Genetics* 6 (3): 167–78.
- Wimmer Ernst A. 2005. "Insect Transgenesis by Site-Specific Recombination." *Nature Methods* 2 (8): 580–82. [PubMed: 16094381]
- Windbichler Nikolai, Papathanos Philippos Aris, and Crisanti Andrea. 2008. "Targeting the X Chromosome during Spermatogenesis Induces Y Chromosome Transmission Ratio Distortion and Early Dominant Embryo Lethality in *Anopheles Gambiae*." *PLoS Genetics* 4 (12): e1000291. [PubMed: 19057670]

- Wood Roger J., and Newton Martha E.. 1991. "Sex-Ratio Distortion Caused by Meiotic Drive in Mosquitoes." *The American Naturalist* 137 (3): 379–91.
- Xiang Youbin, and Scott Hawley R. 2006. "The Mechanism of Secondary Nondisjunction in *Drosophila Melanogaster* Females." *Genetics* 174 (1): 67–78. [PubMed: 16816415]
- Yang Shu Yuan, Baxter Ellen M., and Van Doren Mark. 2012. "Phf7 Controls Male Sex Determination in the *Drosophila* Germline." *Developmental Cell* 22 (5): 1041–51. [PubMed: 22595675]
- Zhang P, and Spradling AC. 1994. "Insertional Mutagenesis of *Drosophila* Heterochromatin with Single P Elements." *Proceedings of the National Academy of Sciences* 91 (9): 3539–43.
- Zhang P, and Stankiewicz RL. 1998. "Y-Linked Male Sterile Mutations Induced by P Element in *Drosophila Melanogaster*." *Genetics* 150 (2): 735–44. [PubMed: 9755204]
- Zuo Erwei, Huo Xiaona, Yao Xuan, Hu Xinde, Sun Yidi, Yin Jianhang, He Bingbing, et al. 2017. "CRISPR/Cas9-Mediated Targeted Chromosome Elimination." *Genome Biology* 18 (1): 224. [PubMed: 29178945]

Figure 1. Targeted transgene insertion on the *D. melanogaster* Y-chromosome. (A) A map of the *D. melanogaster* Y-chromosome (adapted from the UCSC Genome Browser) shows the position of attempted transgene insertions (vertical bars labelled A-J; red bars indicate successful insertions, while grey bars indicate insertion failure) relative to the location of major genes. Green lines indicate conservation of gene regions calculated via the PhastCons method, and grey bars indicate multiple alignment of gene regions, in 27 other species. (B) A schematic of the basic transgene utilized is shown. The transgene contains a marker flanked by homology arms to a specific region of the Y-chromosome, and is injected with a sgRNA that targets a sequence on the Y-chromosome between the homology arms, and a source of Cas9 (B). Following a Cas9-induced double-strand DNA break (DSB), the cell's homology-directed repair (HDR) pathway utilizes the transgene as a repair template to copy the marker between the regions of homology (B'), generating a Y-chromosome with a marker gene at the precise sgRNA-targeted site (B''). (C) The number of embryos injected for each of the Y-chromosome targeting transgenes, survival to larval stage of injected embryos, and transgenesis rate (# of independent transgenic individuals found/# of embryos injected, half of which are predicted to be male) is shown. (D) All males (and none of the females) for

each of the two recovered transgenic lines (AByF (D) and AByG (D')) express eye-specific red fluorescence.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2. Using transgenic marked Y-chromosomes to track X-chromosome meiotic nondisjunction events. (A) In *D. melanogaster*, chromosome nondisjunction can occur in meiosis I or meiosis II; in either case, nondisjunction of X-chromosomes leads to formation of two types of aberrant gametes - ones that have an extra copy of the X-chromosome, and ones that lack an X-chromosome entirely. (B) If nondisjunction occurs during meiosis in a female, she will produce four types of viable offspring when outcrossed to a normal male: normal (XX) females, normal (XY) males, sterile (XO) males, and fertile (XXY) females. (C) Outcrosses of a marked Y-chromosome line to four distinct X-chromosome balancer lines (leftmost column) resulted in progeny arising from nondisjunction events. Frequencies of XXY females (females with marked Y-chromosome; middle column) and XO males (males with no Y-chromosome; rightmost column) are shown.