Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Key Words: Traumatic Brain Injury, Neuropsychological Outcomes, Traumatic Microbleed

Research Poster 1710060

Experiences of Physical Therapists During the COVID-19 Pandemic: An Interpretive Phenomenological Analysis

Marc Campo (Mercy College), Matthew Hyland, Ruth Hansen

Research Objectives: To describe the experiences of Physical Therapists (PTs) in the initial wave of the COVID-19 pandemic in an area where the pandemic was initially very intense.

Design: This was a qualitative study with an interpretive phenomenological approach. Data were collected during 8 focus groups of 2-3 physical therapists each (n=22).

Setting: Focus groups were conducted via videoconference.

Participants: A purposive sampling strategy was used. To be included, participants had to be licensed PTs, worked in facilities in the NY metro area at the onset of the pandemic (March 1, 2020), and had to have had direct patient-care responsibilities. Participant work settings included: acute care, inpatient rehabilitation, homecare, outpatient clinics, school systems, and skilled nursing facilities.

Interventions: This was a qualitative study without interventions.

Main Outcome Measures: An open, inductive coding process was used to identify codes, categories and themes. To improve trustworthiness the investigators took notes, debriefed, and maintained an audit trail. All 3 investigators participated in the coding process.

Results: Four themes emerged including: 1) Everything was disrupted; 2) It was not safe; 3) It was overwhelming; and 4) There was a professional transformation. In hospitals and skilled nursing, participants described chaos, inadequate communication, and unsafe working environments. In outpatient settings, participants experienced furloughs and layoffs along with challenges adapting to telehealth. As the pandemic progressed, inpatient PTs gained confidence and grew into new roles as essential members of interdisciplinary care teams. Outpatient therapists adapted to telehealth and as patients returned to therapy.

Conclusions: Physical therapists experienced extreme challenges during the initial phases of the pandemic. As the pandemic progressed, they redefined practice in ways that will be instructive as the pandemic continues.

Author(s) Disclosures: The authors of this study have no disclosures to acknowledge.

Key Words: Physical Therapists, Coronavirus, Occupational Health Rehabilitation

Research Poster 1710062

The Development of a Feedback Tool To Support Post-Graduate Trainees' Competency Growth and Development in Psychiatry Ambulatory Clinics

Alice Kam (Toronto Rehabilitation Institute), Aisha Husain, Denyse Richardson

Research Objectives: To develop a feedback tool to support post-graduate trainees' competency growth and development in psychiatry ambulatory clinics.

Design: Literature review on feedback response and receptivity in medical education was carried out.

Setting: Ambulatory clinics.

Participants: Post-graduate trainees.

Interventions: Post-graduate trainees will be surveyed and their responses will be analyzed.

Main Outcome Measures: Trainees' survey response.

Results: To be finalized in the phase 2 study.

Conclusions: Phase 2 of this study will use a usability approach to further refine this clinical tool.

Author(s) Disclosures: No conflicts. No relationship or bias.

Key Words: Medical Education, Feedback, Field Based Curricula Self Assessment, Receptivity

Research Poster 1710063

Early Physical and Occupational Therapy and Discharge Disposition in Patients with Sepsis

Karen Clark (CHI Health),

Research Objectives: To explore whether implementing an early mobility protocol conducted by professionals from PT and/or occupational therapy (OT) is associated with discharge location for a group of patients with sepsis, septic shock, or severe sepsis.

Design: A quasi-experimental study design was used with a non-equivalent post-test design. The data were collected for 3 months.

Setting: Mid-sized Midwestern hospital.

Participants: The population of the usual care group were 117 adult patients admitted to any unit with a diagnosis of sepsis, septic shock, or severe sepsis from February 1, 2019, through April 30, 2019. The early rehabilitation group were 96 patients admitted May 6, 2019 through July 31, 2019.

Interventions: The usual care group received PT and OT when an order was received from the hospital medicine provider. The experimental group were treated with PT and OT per an early rehabilitation protocol.

Main Outcome Measures: The dependent variables were number of PT and OT visits and discharge disposition.

Results: There was no statistically significant ($p < 0.05$) difference found between the groups for PT treatments ($t(131) = -0.307$, $p = 0.760$) nor for the OT treatments ($t(104) = 0.925$, $p = 0.357$). The mean number of PT and OT visits in each group were approximately the same as well (PT visits 3.86, SD = 3.83 vs. 4.05 visits, SD = 3.302; OT visits, 3.85, SD = 3.764 vs. 3.25, SD = 2.855; Table 2).

The data demonstrated that the patients who received PT and OT were significantly more likely to be discharged to home/home health care ($p = 0.029$ for PT, $p = 0.047$ for OT) rather than to inpatient healthcare services (See Table 3).

Conclusions: This study demonstrates that the inclusion of PT and OT treatment in the care of patients with sepsis-related illness admitted to the hospital may influence their discharge disposition. The majority of the patients in this study were able to discharge to home.

Author(s) Disclosures: The author, Karen Clark, does not have any conflicts of interest in regards to this study.

Key Words: Sepsis, Physical Therapy, Occupational Therapy Early Mobility, Acute Care

Research Poster 1710064

Impact of Visual Discomfort Symptoms On SDMT Performance Among Persons With Multiple Sclerosis

Michelle Chen (Kessler Foundation), Kristen Costanza, Alyssa Bayley, Nancy Chiaravallotti, Yael Goverover, Silvana L Costa

Research Objectives: To examine how visual discomfort symptoms affect performance on the symbol digit modalities test (SDMT), a visually mediated information processing speed task which is considered a gold standard measure for cognition in multiple sclerosis (MS) research.

Design: Observational, cross-sectional study.

Setting: General community.

Participants: Eighty-seven persons with MS and 30 healthy controls (HCs).

Interventions: Not applicable.

Main Outcome Measures: Visual discomfort questionnaire (VDQ) and SDMT.