Published in final edited form as:

J Int Neuropsychol Soc. 2021 September ; 27(8): 761–775. doi:10.1017/S1355617721000394.

Exercise, arterial stiffness, and cerebral vascular function: potential impact on brain health

Jill N. Barnes^{1,2}, Andrew G. Pearson¹, Adam T. Corkery¹, Nicole A. Eisenmann¹, Kathleen B. Miller¹

¹Bruno Balke Biodynamics Laboratory, Department of Kinesiology, University of Wisconsin-Madison, Madison, WI, USA

²Division of Geriatrics and Gerontology, Department of Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Abstract

Exercise is associated with higher cognitive function and promising intervention to reduce the risk of dementia. With advancing age, there are changes in the vasculature that have important clinical implications for brain health and cognition. Primary aging and vascular risk factors are associated with increases in arterial stiffness and pulse pressure, and reductions in peripheral vascular function.

Objective: The purpose is to discuss the epidemiological, observational, and mechanistic evidence regarding the link between age-related changes in vascular health and brain health.

Methods: We performed a literature review and integrated with our published data.

Results: Epidemiological evidence suggests a link between age-related increases in arterial stiffness and lower cognitive function, which may be mediated by cerebral vascular function, including cerebral vasoreactivity and cerebral pulsatility. Age-associated impairments in central arterial stiffness and peripheral vascular function have been attenuated or reversed through lifestyle behaviors such as exercise. Greater volumes of habitual exercise and higher cardiorespiratory fitness are associated with beneficial effects on both peripheral vascular health and cognition. Yet, the extent to which exercise directly influences cerebral vascular function and brain health, as well as the associated mechanisms remains unclear.

Conclusion: Although there is evidence that exercise positively impacts cerebral vascular function, more research is necessary in humans to optimize experimental protocols and address methodological limitations and physiological considerations. Understanding the impact of exercise on cerebral vascular function is important for understanding the association between exercise and brain health and may inform future intervention studies that seek to improve cognition.

Keywords

aging; cardiorespiratory fitness; cerebral blood flow; cognitive function; endothelial function; imaging; pulsatility

Introduction

Epidemiological evidence consistently shows that exercise is associated with improved cognitive function throughout the lifespan (Ahlskog, Geda, Graff-Radford, & Petersen, 2011; Geda et al., 2010; Sofi et al., 2011). Thus, exercise is an attractive potential intervention to reduce the risk of cognitive decline. The mechanisms underlying exercise's protective role against cognitive decline are not well understood, though there are a few hypotheses. The neurogenesis hypothesis proposes that exercise causes an upregulation of factors called neurotrophins that directly influence neurogenesis and neuroplasticity. The vascular hypothesis is associated with the link between exercise and cognitive function arises from the coexistence of both vascular dysfunction and cognitive decline in many patients. The vascular hypothesis states that impairments in vessel function, which can range from deficiencies in the large central arteries to the brain capillaries, affect neuronal health and thus cognitive ability (de la Torre, 2004). This review will focus on the vascular hypothesis.

The brain has a vast vascular network tasked with precisely and adequately supplying blood flow to its 86 billion active neurons. The neurovascular unit (NVU) in the brain coordinates blood flow regulation and includes several cell types: endothelial cells, vascular smooth muscle cells, pericytes, astrocytes, and neurons. The majority of information regarding NVU function comes from animal models. These preclinical models suggest that early declines in vascular regulation or NVU function precede neuropathological changes (Iadecola, 2004; Kisler, Nelson, Montagne, & Zlokovic, 2017; Sweeney, Kisler, Montagne, Toga, & Zlokovic, 2018). The extent to which the mechanisms identified from preclinical models extend to humans is unknown. There is however supporting evidence that early changes in vascular biomarkers and declines in cerebral blood flow occur before common neurological biomarkers such as changes in amyloid- β , tau, or hippocampal atrophy (Albrecht et al., 2020; Iturria-Medina et al., 2016). Thus, an ideal time to initiate interventions in an effort to delay or prevent the onset of cognitive decline is midlife, when age-related declines in vascular regulation are emerging, and the neuropathological burden is low. Because exercise is considered protective against age-related vascular dysfunction, exercise may therefore reduce the risk of cognitive decline and dementia through its beneficial effect on the systemic and cerebral vasculature.

Lifestyle interventions are promising for reducing age-related morbidity and mortality. Aerobic exercise interventions have been shown to improve vascular function in the large central arteries and in the peripheral circulation (Tanaka, 2019), but the extent to which this translates to the cerebral circulation, influences neuronal function, or impacts cognition is unclear. The purpose of this review is to discuss the effect of exercise on the function of the large central arteries supplying the brain, the impact on the cerebral circulation, as well as

to discuss potential physiological mechanisms and considerations underlying the association between exercise and cognition within the context of primary aging.

Age-associated changes in structure and function of large central arteries

With advancing age, there are alterations in structure and function of the central and peripheral arteries. The structural properties of arteries begin to change with age, as evidenced by vessel wall thickening (Tanaka, Dinunno, Monahan, DeSouza, & Seals, 2001), increased collagen content (Sun, 2015), vascular calcification (Lee & Oh, 2010), deposition of advanced glycation end products (AGEs) (Lee & Oh, 2010), and remodeling of the extracellular matrix (Kohn, Lampi, & Reinhart-King, 2015). Taken together, these maladaptive changes in arterial structure contribute to an increase in large artery stiffness as evidenced by stiffening of the ascending and descending aorta, as well as the carotid arteries (Thijssen, Carter, & Green, 2016; X. Xu et al., 2017).

Because the large central arteries supply the brain with blood flow, alterations in structure and function of the large central arteries can influence hemodynamics in the cerebral circulation (de Roos, van der Grond, Mitchell, & Westenberg, 2017). Indeed, stiffening of the large central arteries is associated with reduced cerebral blood flow (Lu et al., 2011; Tomoto, Riley, Turner, Zhang, & Tarumi, 2020) and consequently hypoperfusion of the brain (Yang, Sun, Lu, Leak, & Zhang, 2017). This is evident in both animal and human models. For example, in an experimental mice model, inducing calcification predictably altered the structure and function of the carotid artery and caused detrimental hemodynamic changes in the cerebral circulation, as well as hippocampal neurodegeneration (Sadekova, Vallerand, Guevara, Lesage, & Girouard, 2013). Increased carotid arterial stiffness in mice is associated with decreased resting cerebral blood flow, a disrupted blood-brain barrier, and impaired neurovascular coupling and cerebral autoregulation (Muhire et al., 2019).

In humans, detrimental hemodynamic changes in the cerebral circulation may manifest as an increase in cerebral pulsatility (Mitchell, 2008), elevated cerebrovascular resistance, and lower cerebrovascular conductance. Cerebral pulsatility increases with age (Pase, Grima, Stough, Scholey, & Pipingas, 2014) and is associated with central arterial stiffness (T. Y. Xu et al., 2012). Indeed, large cohort studies have demonstrated associations between arterial stiffness and cerebral pulsatility (Mitchell et al., 2011). Additionally, higher arterial stiffness is associated with greater cerebrovascular resistance (Robertson, Tessmer, & Hughson, 2010), and lower cerebrovascular conductance (Jaruchart, Suwanwela, Tanaka, & Suksom, 2016). Cerebral pulsatility of the middle cerebral artery is associated with central and brachial pulse pressure (T. Y. Xu et al., 2012). In patients with leukoaraiosis, cerebral pulsatility of the middle cerebral artery is associated with aortic pulse wave velocity, a marker of arterial stiffness (Webb et al., 2012). Thus, impaired dampening of pulsatile flow caused by arterial stiffness may lead to greater transmission of pressure pulsatility to the cerebral circulation.

Unfavorable changes in large artery stiffness are associated with elevated cerebral pulsatility which could lead to impaired vascular endothelial function. Impaired vascular endothelial function is the diminished ability of the endothelial layer of the blood vessels to respond to vasoactive substances and regulate blood flow (Seals, Jablonski, & Donato, 2011).

Higher arterial stiffness, measured by aortic pulse wave velocity, is associated with impaired endothelial function (assessed by flow-mediated dilation) (McEniery et al., 2006). In addition, elevated central and peripheral pulse pressure is associated with impaired endothelial function (McEniery et al., 2006; Nigam, Mitchell, Lambert, & Tardif, 2003). Therefore, stiffening of central arteries and elevated pulse pressure could lead to changes in the structure and function of the peripheral vasculature. In the cerebral circulation, increased pulse pressure in an experimental model led to hypertrophy of cerebral arterioles and vascular remodeling (Baumbach, 1996). The study by Baumbach, et al. supports the idea that an increase transmission of pulsatile blood flow (i.e. pulse pressure) to the peripheral and cerebral circulations, results in vascular endothelial damage and impaired endothelial function (Baumbach, 1996; Mitchell, 2008; Thorin-Trescases et al., 2018) (Mitchell, 2008, Thorin-Trescases et al., 2018).

Aging is also associated with systemic endothelial dysfunction, as measured by endothelium-dependent dilation in the coronary arteries (Egashira et al., 1993) and peripheral arteries (Gerhard, Roddy, Creager, & Creager, 1996). Multiple factors are associated with the age-related decrease in systemic endothelial function including both a reduction in the bioavailability of nitric oxide (NO), a potent vasodilator, (Taddei et al., 2001), and an increase in production of endothelin-1, a potent vasoconstrictor (Thijssen et al., 2016; Thijssen, Rongen, van Dijk, Smits, & Hopman, 2007). Similarly, age-related increases in large artery stiffness and cerebral pulsatility likely impact vascular endothelial function in the cerebral circulation (Thorin-Trescases et al., 2018; Walker et al., 2015), although mechanistic data in humans are lacking. For more information on age-related changes in peripheral vascular function, there are many excellent reviews available (de Roos et al., 2017; Seals et al., 2011; Thijssen et al., 2016; X. Xu et al., 2017). Collectively, changes in the structure and function of central and peripheral arteries contribute to stiffening of the large central arteries and unfavorable alterations in cerebrovascular hemodynamics which may increase the risk of cognitive decline and dementia.

Age-related changes in vascular function influence brain health

The transmission of highly pulsatile pressure and flow to the microvasculature has been linked with tissue and organ damage (Mitchell, 2008). Because the brain and kidneys are organs with high blood flow and low impedance, the microcirculation within these organ systems are particularly susceptible to damage due to age-related changes in arterial structure (Mitchell, 2008; O'Rourke & Hashimoto, 2007). The idea that elevated arterial stiffness results in cerebrovascular damage and structural remodeling in the brain leading to impaired function is supported by human and clinical evidence (de Roos et al., 2017). For example, elevated central arterial stiffness has been associated with increased white matter hyperintensities in healthy non-hypertensive middle-aged women (Barnes et al., 2017), and adults with cognitive complaints (Kearney-Schwartz et al., 2009). In addition, arterial stiffness is associated with cerebral microbleeds (Ding et al., 2015), cerebral small vessel disease (CSVD) (Zhai et al., 2018), and lower brain volume (Palta et al., 2019). Importantly, these changes in brain structure may be mediating factors amongst the relationship between arterial stiffness and cognitive function, as shown in large clinical studies (Elias et al., 2009; Palta et al., 2019; Watson et al., 2011).

Alterations in vascular structure and function occur in both the large and small intracranial vessels. CSVD is often used to refer to the disease of the cerebral microvessels. The most common diagnostic method of CSVD is the appearance of white matter hyperintensities on MRI scans, which are areas denoting leukoaraiosis (Sargurupremraj et al., 2020). CSVD has been associated with altered cerebral hemodynamics (Robertson et al., 2019) and an increased risk of cognitive impairment (Bos et al., 2018). The relationship between CSVD and dementia has been the subject of many recent reviews and meta-analyses; however, increases in arterial stiffness and alterations in cerebral blood flow may precede the diagnosis of CSVD or CSVD related pathologies (Shi et al., 2016; van Sloten et al., 2015). While arterial stiffness is associated with structural changes in the brain in patients with CSVD, this review will focus on alterations of vascular function associated with primary aging in individuals without established CSVD.

Regular exercise and central arterial stiffness

Exercise has the ability to affect both the structure and function of the central elastic arteries; however, its effects are dependent on numerous factors, such as mode, volume, and frequency of exercise performed. Studies examining the effect of exercise on central arterial stiffness in middle-aged and older adults have largely shown beneficial results, such that central arterial stiffness is lower in exercise trained middle-aged and older adults and is inversely associated with cardiorespiratory fitness or VO_2max (Tanaka, DeSouza, & Seals, 1998; Tanaka et al., 2000; Vaitkevicius et al., 1993). Furthermore, there is evidence that elevated levels of arterial stiffness can be reversed through short-term moderate-intensity aerobic exercise interventions in this population (Tanaka et al., 2000; Yoshizawa et al., 2009). However, there may be a threshold of exercise volume necessary for the prevention of age-associated changes. Recently, Shibata et al. have reported 4-5 days per week of at least 30 minutes of aerobic exercise is required to prevent age-associated changes in central arterial stiffness (Shibata et al., 2018). This idea of an exercise volume threshold is consistent with earlier publications showing lower levels of central arterial stiffness in endurance trained older men (performing vigorous aerobic exercise at least 5 days per week and competing in local running races), but not recreationally active older men (light to moderate exercise 3 days per week) when compared with their sedentary peers (Tanaka et al., 2000). A recent meta-analysis by Ashor, et al. also concluded that higher intensity aerobic exercise resulted in lower central arterial stiffness in adults (Ashor, Lara, Siervo, Celis-Morales, & Mathers, 2014). Importantly, not all modes of exercise appear to have the same effects on central arterial stiffness. Most studies focus on aerobic exercises such as walking, running, or cycling, as the primary exercise mode when evaluating the impact on arterial stiffness. Healthy adults who participate in other modes of aerobic exercise, such as swimming or rowing, also show reduced levels of age-related arterial stiffness compared with sedentary adults (Cook et al., 2006; Nualnim, Barnes, Tarumi, Renzi, & Tanaka, 2011). In contrast, adults who regularly participate in high-intensity resistance exercise paradoxically have higher arterial stiffness (Miyachi et al., 2003). Furthermore, high-intensity progressive resistance training interventions 3-4 days per week (near 80% 1 repetition maximum) have been shown to increase central arterial stiffness in both men (Miyachi et al., 2004) and women (Cortez-Cooper et al., 2005). Although the underlying mechanisms are not clear, high intensity resistance exercise produces large fluctuations in

blood pressure during the activity, which over time can alter the load-bearing properties of the artery (Miyachi, 2013). However, this increase in arterial stiffness only appears in younger adults or when the intervention includes high intensity resistance exercise, with studies using moderate intensity resistance training showing no effect on central arterial stiffness (Ashor et al., 2014; Miyachi, 2013). Therefore, both exercise volume and modality of exercise influences arterial stiffness in the large central arteries.

Aging and vascular function of the cerebral circulation

This review has largely focused on the function of the large central arteries which supply the brain. From the large central arteries, blood travels into the intracranial vessels and through the Circle of Willis before it reaches its destination at the neuronal tissue. Blood flow to the neuronal tissue is regulated by the smaller brain blood vessels, astrocytes, pericytes and neurons that form the NVU to help control oxygen delivery to match increases in metabolic activity in the brain. The vascular smooth muscle cells at the level of the arterioles helps to control the amount of blood flowing into the capillary bed and the NVU precisely dictate the regional distribution of blood flow. Thus, the structure and function throughout the entire brain's vascular network, from the large central blood vessels to the pericytes in the brain capillaries, are essential for maintaining brain health (L. S. Brown et al., 2019; Iadecola, 2017; Kisler et al., 2017; Sweeney et al., 2018).

Many studies suggest that reduced vessel function in the cerebral circulation (or cerebral vascular function) is part of the etiology of cognitive decline. For example, individuals with lower cerebral blood flow are at a higher risk for Alzheimer's disease (Wolters et al., 2017). In addition, Alzheimer's disease patients demonstrate early vascular dysregulation (Arvanitakis, Capuano, Leurgans, Bennett, & Schneider, 2016; Iturria-Medina et al., 2016; Laing et al., 2020; Montagne et al., 2016), reduced cerebrovascular function (Silvestrini et al., 2006), and post-mortem evaluations show both vascular pathology and Alzheimer's disease pathology (Toledo et al., 2013). Furthermore, early vascular-tau associations may be exacerbated in the presence of amyloid- β , suggesting a synergistic effect of vascular and neurodegenerative pathologies (Albrecht et al., 2020). Thus, evaluating vascular health and cerebral vascular function may be important in understanding the trajectory of Alzheimer's disease. This may be especially critical in prodromal stages of cognitive decline when neuropathology and cognitive symptoms are below detection thresholds.

Vascular function in the cerebral vessels in humans can be assessed by measuring global or regional cerebral perfusion at rest or assessing the blood flow response to a vasoactive stimulus. Several vasoactive stimuli are more frequently used to assess cerebral vascular function, including alterations in arterial blood gases (O_2 and CO_2), acetazolamide, or neuronal stimulation using visual perturbations or cognitively challenging activities (Fierstra et al., 2013). The magnitude of the blood flow response to these stimuli determines cerebral vasoreactivity, with higher vasoreactivity to a vasodilatory stimulus indicating healthier cerebral blood vessels.

While there are an increasing number of studies that include cerebral vascular function, the relationship between advancing age and cerebral vascular function is still not well understood. One reason for this is because of the inherent difficulty of imaging blood flow

through the skull. Common techniques attempt to measure blood flow through the larger intracranial vessels include ultrasonography or 2D, 3D or 4D phase contrast magnetic resonance imaging (MRI). Other techniques to measure bulk flow, or infer microvascular flow, include measurement of oxy- and deoxy-hemoglobin concentration with near-infrared spectroscopy (NIRS), estimation of cerebral blood flow using an invasive tracer such as quantitative [^{15}O] water brain positron emission tomography (PET) imaging, measurement of oxy- and deoxy-hemoglobin using blood-oxygen level dependent imaging (BOLD) MRI, and estimating cerebral blood flow with an intrinsic tracer (radiofrequency labeled arterial blood) via arterial spin labeling (ASL) techniques. PET and MRI techniques can measure global or regional cerebral blood flow, or blood flow through the large intracranial vessels. However, MRI is not always available and scanning sequences cause a temporal constraint. Transcranial Doppler (TCD) ultrasound was developed in the early 1980's and has become a popular technique to measure brain blood flow because it allows for the measurement of blood velocity through large intracranial vessels with high temporal resolution to assess rapid changes in the cerebral circulation. It also allows for screening of participants who could not safely undergo MRI scans and permits experimental measurements that would be restricted in an MRI. However, TCD lacks spatial resolution, can only image one vessel at a time, and measures velocity rather than flow. The difference between blood velocity and blood flow is an important distinction, especially in the instance where a stimulus may change diameter of the vessel of interest. Currently there is no gold standard technique to measure cerebral blood flow and each technique has strengths and weaknesses (see review (Tymko, Ainslie, & Smith, 2018)).

Using the TCD technique, studies have shown that primary aging, in the absence of cardiovascular risk factors, is associated with lower cerebral vasoreactivity (Bakker et al., 2004; Barnes, Schmidt, Nicholson, & Joyner, 2012; Kastrup, Dichgans, Niemeier, & Schabet, 1998), indicating a reduction in cerebral blood vessel health. However, there are also studies using TCD that show no effect of age on cerebral vasoreactivity (Galvin et al., 2010; Madureira, Castro, & Azevedo, 2017; Miller, Howery, Harvey, Eldridge, & Barnes, 2018) and suggest preserved cerebral vascular function (see additional review (Hoiland, Fisher, & Ainslie, 2019)). These contrasting results may be due to several methodological factors (as discussed above), that may mask the effects of primary aging, including the influence of arterial blood pressure, body posture during data collection, use and timing of medication, or other experimental procedures used to deliver a vasoactive stimulus. For example, arterial blood pressure is important for the interpretation of cerebral blood flow because an increase in perfusion pressure will drive an increase cerebral blood flow. This is in part blunted if cerebral autoregulation is intact, but there is a complex relationship between arterial blood pressure and cerebral blood flow across a range of physiological conditions (Tzeng et al., 2012). Importantly, physiological factors in the aforementioned studies may also influence the conflicting results, including whether or not the study participants had high cardiorespiratory fitness or were endurance-trained (Bailey et al., 2013; Barnes, Taylor, Kluck, Johnson, & Joyner, 2013; Miller et al., 2018; Murrell et al., 2012; Thomas et al., 2013). In order to determine age-related changes in cerebral vasoreactivity in healthy adults vs. pathological declines in cerebral vasoreactivity, future studies should provide sufficient characterization of participants and

consider experimental factors such as body posture, simultaneous arterial blood pressure measurements, medication or hormone use, and consider the imaging modality best suited for the research question.

One methodological factor warrants additional discussion because of its impact on the effect of aging on cerebral vasoreactivity. TCD ultrasound, used by previous studies to evaluate cerebral vasoreactivity, has the limitation of only measuring blood velocity and not blood flow, as it cannot assess blood vessel diameter. Using MRI to measure the middle cerebral artery response to vasoactive stimuli, recent studies have shown that the middle cerebral artery vasodilates under vasoactive conditions like elevated CO₂ (Coverdale, Gati, Opalevych, Perrotta, & Shoemaker, 2014; Miller et al., 2019; Verbree et al., 2014). This would suggest that changes in blood velocity obtained by TCD may not accurately represent changes in blood flow during a vasoactive stimulus. In order to address this limitation of TCD, our group has utilized 4D flow MRI which has the advantage of measuring structural and flow data of large intracranial vessels in the same scan acquisition. This technique has been validated at rest (Rivera-Rivera et al., 2016; Schrauben et al., 2014) and during hypercapnia (Mikhail Kellawan et al., 2016) to measure blood flow in over 11 large intracranial vessels. When these changes in blood vessel diameter are evaluated with changes in blood velocity simultaneously (via 4D flow MRI) we have shown reduced cerebral vasoreactivity in healthy older adults, compared with young adults (Miller et al., 2019). This is in contrast to our previous work in healthy older adults using TCD ultrasound, when the participant characteristics were well-matched with healthy young adults (Miller et al., 2018). When using 4D flow MRI to measure the magnitude of middle cerebral artery diameter changes in response to CO₂, vasodilation was only significant in young adults (Miller et al., 2019). In other words, the middle cerebral artery (and potentially other intracranial blood vessels) diameter increases in response to a vasoactive stimulus, but only in young healthy adults. These age-associated differences could potentially explain some of the conflicting results for cerebral vasoreactivity. However, it is also unknown if factors such as cardiorespiratory fitness or habitual endurance-training influences the magnitude of cerebral artery dilation in the presence of a vasoactive stimulus like CO₂. Our hypothesis is that cardiorespiratory fitness is positively associated with the magnitude of middle cerebral artery dilation (based on our unpublished observations), but this has not been systematically tested. More research is necessary to determine how habitual exercise and cardiorespiratory fitness may mediate the effects of aging on peripheral and cerebral vascular function.

Exercise and the cerebral circulation

The impact of aging on cerebral vasoreactivity could be influenced by habitual aerobic exercise, and there are conflicting reports in the literature. Our previous work using TCD ultrasound has shown that VO₂max is associated with cerebral vasoreactivity in healthy older adults, but are not correlated in young adults (Barnes et al., 2013). Additionally, while our previous studies have shown lower cerebral vasoreactivity in healthy older adults, compared with younger adults (Barnes et al., 2012), we reported no age-related difference in cerebral vasoreactivity when including only habitually exercising adults (Miller et al., 2018). Other groups have also examined the influence of habitual exercise or VO₂max on cerebral vasoreactivity using TCD ultrasound and have reported either no association with exercise

(Braz, Fluck, Lip, Lundby, & Fisher, 2017; Zhu et al., 2013), or a positive association between VO₂max and cerebral vasoreactivity in young and older men (Bailey et al., 2013). Studies that have examined the effect of exercise on the cerebral circulation using MRI methods such as ASL or BOLD have added to the conflicting literature on the relationship between exercise and cerebral blood flow and cerebral vascular function. These studies have shown reduced cerebral vasoreactivity in highly-trained Master's athletes compared with sedentary controls (Thomas et al., 2013), and conflicting relationships between VO₂max and cerebral vasoreactivity (Furby, Warnert, Marley, Bailey, & Wise, 2019; Intzandt et al., 2020).

While we have focused on the impact of exercise on cerebral vascular function, there are studies investigating the effect of exercise on resting cerebral perfusion. Again, there are conflicting results in the literature with some studies reporting an increase in regional cerebral perfusion (Alfini, Weiss, Nielson, Verber, & Smith, 2019; Chapman et al., 2013; Guadagni et al., 2020; Kleinloog et al., 2019) or no effect of exercise training on resting cerebral blood flow (Flodin, Jonasson, Riklund, Nyberg, & Boraxbekk, 2017; van der Kleij et al., 2018).

Collectively, the majority of studies have found a neutral or favorable effect of exercise on cerebral vasoreactivity, cerebral perfusion, and thus, cerebral vascular function. However, longitudinal or intervention studies are necessary to determine the influence of habitual exercise on cerebral vasoreactivity, as the conflicting evidence may be due to: 1) methodological factors such as using TCD ultrasound vs. MRI and the method of delivering the vasoactive stimulus; 2) study design, particularly the lack of women participants in earlier studies; or 3) physiological effects of exercise that are unrelated to vascular health, such as altered chemoreceptor or baroreceptor sensitivity.

Potential mechanisms underlying the benefit of exercise

The mechanisms underlying exercise's protective role against cognitive decline are not well understood. A single bout of exercise leads to considerable changes in metabolic, cardiovascular, and immune pathways (Contrepois et al., 2020; Egan & Zierath, 2013; Pedersen & Hoffman-Goetz, 2000; Whyte & Laughlin, 2010). The effects of habitual exercise, as well as the mode, frequency, and intensity of exercise needed to invoke the favorable effects of exercise on brain health remain unclear. In addition, there are other favorable effects of exercise that will not be covered in this review. For example, psychological benefits of both acute and chronic exercise may, over the long term, affect cognition independent of vascular adaptations. Diagnosis of anxiety and/or depression is associated with an increased risk of cardiovascular disease (Cohen, Edmondson, & Kronish, 2015) and dementia (Kuring, Mathias, & Ward, 2018). However, acute exercise and exercise training can reduce psychosocial stress (O'Keefe, O'Keefe, & Lavie, 2019), which could indirectly benefit cognitive function. Another example of the favorable adaptations of exercise is the effect of learning or cognitive activity associated with exercise training, particularly when training for many competitive sports. Again, is difficult to tease out the physiological effects of movement from the neural engagement associated with learning the movement or new activity. The remainder of this section will focus on the direct physiological mechanisms underlying the benefits of exercise.

In a breakthrough study in 1999, Van Praag and colleagues described the development of new neurons in the hippocampus in mice that were exposed to wheel running (van Praag, Christie, Sejnowski, & Gage, 1999; van Praag, Kempermann, & Gage, 1999). This study is associated with the neurogenesis hypothesis, suggesting that exercise causes an upregulation of factors called neurotrophins that directly influence neurogenesis and neuroplasticity. Indeed, exercise has also been shown to upregulate neurotrophins in humans, such as brain derived neurotrophic factor (BDNF) (Church et al., 2016; P. Rasmussen et al., 2009). In humans, positive associations between greater amounts of physical activity and hippocampal volume have been reported (Erickson et al., 2009; Erickson et al., 2010). In addition to BDNF, other signaling molecules have been identified though the mechanisms underlying the cognitive benefits of exercise are still being explored. For additional reviews on this topic (Cotman, Berchtold, & Christie, 2007; Stillman, Esteban-Cornejo, Brown, Bender, & Erickson, 2020; Vivar & van Praag, 2017).

Other than BDNF, upregulation of factors that influence or regulate neural and blood vessel survival and growth, such as insulin-like growth factor (IGF-1) and vascular endothelial growth factor (VEGF) likely contribute to beneficial effect of exercise (Cotman et al., 2007; Erickson et al., 2019; Liu & Nusslock, 2018; Stillman et al., 2020). Other myokines, or cytokines synthesized and released by contracting skeletal muscle such as cathepsin B and irisin may integrate with other growth factors or work via separate pathways to preserve brain vasculature and neural tissue, though these mechanisms are only beginning to be explored (Severinsen & Pedersen, 2020). The exact mechanisms underlying the favorable effects of exercise on brain health remain unclear. Here, we briefly review considerations and potential mechanisms that may mediate the effect of exercise specifically on the cerebral circulation.

When investigating the effect of exercise on resting cerebral perfusion, studies have reported no effect of 16 weeks of moderate to vigorous intensity aerobic exercise interventions on cerebral blood flow in patients with established neurodegenerative disease (van der Kleij et al., 2018). Similarly, a study in middle-aged and older healthy adults report no effect of a low to moderate intensity exercise intervention on cerebral blood flow (Flodin et al., 2017). In contrast, Chapman, et al. reported an increase in regional cerebral perfusion in middle-aged and older healthy adults with a more intense, structured exercise intervention, representing a greater overall exercise dose (Chapman et al., 2013). Another study by Kleinloog et al., reported an increase in resting cerebral blood flow of the frontal cortex after a supervised, progressive aerobic exercise training protocol for 8 weeks in previously sedentary healthy older men aged 60-70 years (Kleinloog et al., 2019). Similarly, Poulin and colleagues recruited healthy older participants to undergo a 6-month aerobic exercise intervention progressively increasing the exercise intensity. Post-intervention, Guadagni, et al. reported increased resting cerebral blood velocity, as well as improved measures of cerebral vasoreactivity (Guadagni et al., 2020). Collectively, these studies suggest that exercise of a sufficient dose and duration of training may influence cerebral blood flow characteristics in healthy adults, at least during the preclinical phase of disease.

As previously discussed in this review, habitual exercise may reduce central arterial stiffness, which corresponds to greater compliance of the large central arteries such as the aorta. It

is possible that the beneficial effects of exercise on vascular structure may extend further than large central vessels and into the intracranial vessels. Using diffuse optical imaging to examine cerebral pulsatile waveforms in older adults, cardiorespiratory fitness was associated with greater arterial compliance of the vessels in frontoparietal brain regions (Tan et al., 2017). In young adults, cardiorespiratory fitness was positively associated with cerebral artery compliance (Furby et al., 2019). Together, these studies suggest that exercise may improve the vascular structure of both large and small intracranial vessels. As cardiorespiratory fitness is comprised of both genetic and lifestyle factors (i.e. exercise habits), future studies could evaluate if an acute exercise intervention may improve arterial compliance in the cerebral circulation. Ultimately, improved intracranial arterial compliance (or reduced arterial stiffness) could lead to reduced pressure pulsatility in the cerebral microvessels, slowed breakdown of the blood brain barrier, and better clearing of circulating amyloid- β (Kisler et al., 2017; Zhao, Nelson, Betsholtz, & Zlokovic, 2015; Zlokovic, 2011).

Together with improvements in arterial compliance, aerobic exercise improves vascular endothelial function in peripheral blood vessels. In the peripheral circulation, exercise increases NO synthesis, regulation and signaling, improves prostaglandin-mediated vasodilation, and decreases oxidative stress and inflammation (DeSouza et al., 2000; Green, Hopman, Padilla, Laughlin, & Thijssen, 2017; Seals et al., 2011; Tanaka, 2019). The endothelial function in the cerebral vessels could benefit from habitual exercise similar to peripheral blood vessels, though mechanistic studies of cerebral vessels in humans are lacking. Animal studies have provided some evidence to suggest exercise-mediated improvements in vascular endothelial function. For example, vigorous aerobic exercise training in rats resulted in higher NO-dependent cerebral vasoreactivity compared to sedentary rats (Arrick, Yang, Li, Cananzi, & Mayhan, 2014). Furthermore, voluntary wheel running in wild-type mice increased basal cerebral blood flow; however, there was no improvement in cerebral blood flow in mice that were endothelial NO synthase (eNOS) deficient (Endres et al., 2003). This suggested that improvements in cerebral blood flow are influenced by aerobic exercise-induced increases in vascular shear stress, but dependent on eNOS upregulation. The increase in blood flow during dynamic aerobic exercise increases shear stress which stimulates the eNOS enzyme to rapidly vasodilate the blood vessel (Green, Maiorana, O'Driscoll, & Taylor, 2004). Along these lines, voluntary wheel running in mice induces beneficial changes in cerebral arteries, with evidence of larger and more regularly shaped endothelial cell nuclei in the middle cerebral artery (Latimer et al., 2011). In humans, postmenopausal women with higher levels of plasma-derived measures of oxidative stress demonstrated reduced cerebrovascular conductance (Pialoux, Brown, Leigh, Friedenreich, & Poulin, 2009). Additionally, higher cardiorespiratory fitness was associated with lower markers of oxidative stress (Pialoux et al., 2009). Because oxidative stress can reduce NO bioavailability, reductions in oxidative stress or increased antioxidant pathways may be an underlying mechanism by which habitual exercise improves cerebral blood flow regulation via increased NO bioavailability (Thijssen et al., 2016). Another possible mechanism underlying exercise-mediated changes in vasodilatory capacity involves the role of prostaglandins (Nicholson, Vaa, Hesse, Eisenach, & Joyner, 2009). With our previous work in older adults, we have shown that cyclooxygenase blockade reduces cerebral vasoreactivity (Barnes et al., 2012). In addition, higher cardiorespiratory fitness

was positively associated with the magnitude of change in cerebral vasoreactivity after cyclooxygenase blockade (reducing prostaglandins) (Barnes et al., 2013). This suggests that higher cardiorespiratory fitness may upregulate the reliance on prostaglandin-mediated vasodilation in the cerebral vessels (Barnes et al., 2013).

This review largely focused on research detailing the impact of aging and exercise on the cerebral vasculature. However, altered energy metabolism is another factor that impacts neurovascular coupling and improved energy metabolism may contribute to the effect of exercise on the brain (for review see (van Praag, Fleshner, Schwartz, & Mattson, 2014)). Because glucose molecules cannot pass through the blood brain barrier without a carrier protein, this rate-limiting step in energy metabolism requires the vascular glucose transporter GLUT1. Importantly, GLUT1 is the primary glucose transporter in endothelial cells that make up the blood brain barrier (Veys et al., 2020). In addition, in the cerebral microvessels, the density and distribution of GLUT1 is correlated with regional cerebral glucose utilization (Zeller, Rahner-Welsch, & Kuschinsky, 1997). Transgenic mice overexpressing amyloid- β precursor protein (APP) with GLUT1 deficiency had lower cerebral blood flow, impaired neurovascular coupling, and increased amyloid- β pathology compared with APP mice (Winkler et al., 2015). This suggests that GLUT1 reductions may be an early modification in the neuropathological processes associated with Alzheimer's disease. In the human brain, Alzheimer's disease is associated with reduced glucose transporters in the blood brain barrier (Kalaria & Harik, 1989; Szablewski 2017).

In mice, acute aerobic exercise (voluntary wheel running) led to an increase in expression of GLUT1 in the brain (Allen & Messier, 2013). Similarly, regular exercise in mice increased expression of GLUT1 in cardiomyocytes (Jiang et al., 2020) and the central nervous system (Pang et al., 2019). In addition, upregulation of GLUT1 in the cortex was reported after a single bout of aerobic exercise in rats (Takimoto & Hamada, 2014). Because cerebral hypometabolism is often observed in Alzheimer's disease, the exercise-mediated upregulation of GLUT1 may contribute to the positive effects of exercise on cognitive and brain health (Pang et al., 2019). It is unknown if similar increases in GLUT1 expression in humans are apparent after an exercise intervention.

Recent research on the glymphatic pathway, a glial-lymphatic system that provides drainage of fluids from brain tissue that was first identified in rodents, suggests that this pathway is also present in humans (M. K. Rasmussen, Mestre, & Nedergaard, 2018). Because of the blood-brain barrier and lack of lymphatic vessels in the parenchyma, the brain cannot rely on the vascular lymphatic system to clear metabolic waste like most other organs. Briefly, the glymphatic pathway clears solutes and metabolic waste via a perivascular pathway. This pathway allows exchange between the cerebrospinal fluid and interstitial fluid, and appears to be facilitated by astrocytic aquaporin-4 (AQP4) water channels (Iliff & Nedergaard, 2013) and arterial pulsations (Mestre et al., 2018). Preclinical models show that aging and CSVD may impair glymphatic function due to attenuated periarterial influx of cerebrospinal fluid and impaired exchange between cerebrospinal and interstitial fluid (Kress et al., 2014; Mestre et al., 2018). In the context of Alzheimer's disease, AQP4 helps to "drain" extracellular amyloid- β , preventing the deposition of amyloid- β plaques (Iliff & Nedergaard, 2013). In support of the data from animal models, patients with

Alzheimer's disease or dementia have reduced tracer clearance compared with controls, indicating impaired glymphatic function, which is increasingly recognized as a key factor in the pathogenesis of Alzheimer's disease (de Leon et al., 2017; Ringstad et al., 2018; Tarasoff-Conway et al., 2015).

An emerging hypothesis is that habitual exercise assists in glymphatic flow to help clear amyloid- β , tau, and other soluble proteins and metabolites. In mice, voluntary running is associated with increased cerebrospinal fluid flow and amyloid- β clearance, indicating enhanced glymphatic function (He et al., 2017; von Holstein-Rathlou, Petersen, & Nedergaard, 2018). Therefore, in addition to the benefits of increased cerebral vascular function, one more advantage of exercise in rodents may be greater clearance of neurotoxic proteins. Yet, the translation of this information to aging humans is unclear. In a study of cognitively normal older healthy adults, those who exceeded exercise guidelines of 7.5 MET-hours/week had higher cerebrospinal fluid concentrations of A β ₄₂, the isoform of amyloid- β commonly found in brain plaques, compared with non-exercisers, which would suggest enhanced clearance (Liang et al., 2010). A study by Brown et al. reported no difference in cerebrospinal fluid concentrations of A β ₄₂ between exercise groups, even though amyloid- β deposition in the brain was lower in exercising adults at elevated risk of Alzheimer's disease (B. M. Brown et al., 2017). However, a short-term moderate to high intensity exercise training intervention in patients with Alzheimer's disease did not change A β ₄₂, tau, or APP concentrations in the cerebrospinal fluid (Steen Jensen et al., 2016). Clearly, there are still many questions in this emerging research area regarding whether exercise enhances activity of the glymphatic pathway to reduce neuropathology in the brain.

Collectively, these findings in animals and humans suggest that exercise may: 1) contribute to structural and functional improvements in the large central arteries supplying the brain and the cerebral arteries; 2) upregulate local and systemic factors that impact neurogenesis and vascular function; and 3) improvements may extend throughout the vascular tree, from large intracranial vessels to the cerebral arteries and microvasculature, and to cerebrospinal fluid flow.

Summary

Regular lifelong aerobic exercise is associated with higher cognitive function scores. In order for exercise to be considered a successful intervention to reduce the risk or delay the onset of cognitive decline, more research is necessary. Because of the time delay between the initial pathological changes and the onset of cognitive symptoms, other variables are necessary to use as targets for evaluating the success of exercise interventions on brain health. These variables may include arterial stiffness of the extracranial vessels, cerebral artery vasoreactivity, or cerebral pulsatility. Arterial stiffening in the large central arteries is associated with indicators of declining brain health and is improved by regular exercise. Therefore, the reported beneficial effects of exercise on arterial stiffness may also indicate favorable changes in the brain. Measuring cerebral vascular function offers a promising way to assess vascular health in the cerebral circulation and could be used as an early biomarker. Although there is evidence that exercise positively impacts cerebral vascular function, more research is necessary to optimize experimental protocols and address methodological

limitations and physiological considerations. Additionally, more information is necessary on the systemic and local mechanisms, the mode of exercise, as well as research on the appropriate exercise “dose” for brain health. Despite the need for more research, exercise nevertheless remains an appealing lifestyle intervention to reduce the risk of cognitive decline. Understanding the impact of exercise of vascular function in the cerebral circulation is important for understanding the association between exercise and brain health and may inform future intervention studies that seek to improve cognition.

Acknowledgements

The authors would like to acknowledge Anna Howery and Kelli England for their critical review of this manuscript. This work was supported by the Alzheimer’s Association 17-499398 (JNB) and the National Institutes of Health grants NS117746 (JNB) and T32 HL007936 (KBM).

References

- Ahlskog JE, Geda YE, Graff-Radford NR, & Petersen RC (2011). Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. *Mayo Clinic proceedings*. Mayo Clinic, 86(9), 876–884. doi:10.4065/mcp.2011.0252
- Albrecht D, Isenberg AL, Stradford J, Monreal T, Sagare A, Pachicano M, ... Pa J (2020). Associations between Vascular Function and Tau PET Are Associated with Global Cognition and Amyloid. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 40(44), 8573–8586. doi:10.1523/JNEUROSCI.1230-20.2020 [PubMed: 33046556]
- Alfini AJ, Weiss LR, Nielson KA, Verber MD, & Smith JC (2019). Resting Cerebral Blood Flow After Exercise Training in Mild Cognitive Impairment. *Journal of Alzheimer's disease : JAD*, 67(2), 671–684. doi:10.3233/JAD-180728
- Allen A, & Messier C (2013). Plastic changes in the astrocyte GLUT1 glucose transporter and beta-tubulin microtubule protein following voluntary exercise in mice. *Behavioural brain research*, 240, 95–102. doi:10.1016/j.bbr.2012.11.025 [PubMed: 23201358]
- Arrick DM, Yang S, Li C, Cananzi S, & Mayhan WG (2014). Vigorous exercise training improves reactivity of cerebral arterioles and reduces brain injury following transient focal ischemia. *Microcirculation*, 21(6), 516–523. doi:10.1111/micc.12127 [PubMed: 24617555]
- Arvanitakis Z, Capuano AW, Leurgans SE, Bennett DA, & Schneider JA (2016). Relation of cerebral vessel disease to Alzheimer's disease dementia and cognitive function in elderly people: a cross-sectional study. *Lancet Neurol*, 15(9), 934–943. doi:10.1016/S1474-4422(16)30029-1 [PubMed: 27312738]
- Ashor AW, Lara J, Siervo M, Celis-Morales C, & Mathers JC (2014). Effects of exercise modalities on arterial stiffness and wave reflection: a systematic review and meta-analysis of randomized controlled trials. *PloS one*, 9(10), e110034. doi:10.1371/journal.pone.0110034 [PubMed: 25333969]
- Bailey DM, Marley CJ, Brugniaux JV, Hodson D, New KJ, Ogoh S, & Ainslie PN (2013). Elevated aerobic fitness sustained throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke; a journal of cerebral circulation*, 44(11), 3235–3238. doi:10.1161/STROKEAHA.113.002589
- Bakker SL, de Leeuw FE, den Heijer T, Koudstaal PJ, Hofman A, & Breteler MM (2004). Cerebral haemodynamics in the elderly: the rotterdam study. *Neuroepidemiology*, 23(4), 178–184. doi:10.1159/000078503 [PubMed: 15272220]
- Barnes JN, Harvey RE, Zuk SM, Lundt ES, Lesnick TG, Gunter JL, ... Kantarci K (2017). Aortic hemodynamics and white matter hyperintensities in normotensive postmenopausal women. *Journal of neurology*, 264(5), 938–945. doi:10.1007/s00415-017-8476-1 [PubMed: 28389742]
- Barnes JN, Schmidt JE, Nicholson WT, & Joyner MJ (2012). Cyclooxygenase inhibition abolishes age-related differences in cerebral vasodilator responses to hypercapnia. *Journal of applied physiology*, 112(11), 1884–1890. doi:10.1152/jappphysiol.01270.2011 [PubMed: 22442028]

- Barnes JN, Taylor JL, Kluck BN, Johnson CP, & Joyner MJ (2013). Cerebrovascular reactivity is associated with maximal aerobic capacity in healthy older adults. *Journal of applied physiology*, 114(10), 1383–1387. doi:10.1152/jappphysiol.01258.2012 [PubMed: 23471946]
- Baumbach GL (1996). Effects of increased pulse pressure on cerebral arterioles. *Hypertension*, 27(2), 159–167. doi:10.1161/01.hyp.27.2.159 [PubMed: 8567036]
- Bos D, Wolters FJ, Darweesh SKL, Vernooij MW, de Wolf F, Ikram MA, & Hofman A (2018). Cerebral small vessel disease and the risk of dementia: A systematic review and meta-analysis of population-based evidence. *Alzheimer's & dementia : the journal of the Alzheimer's Association*, 14(11), 1482–1492. doi:10.1016/j.jalz.2018.04.007
- Braz ID, Fluck D, Lip GYH, Lundby C, & Fisher JP (2017). Impact of aerobic fitness on cerebral blood flow and cerebral vascular responsiveness to CO₂ in young and older men. *Scand J Med Sci Sports*, 27(6), 634–642. doi:10.1111/sms.12674 [PubMed: 26993209]
- Brown BM, Sohrabi HR, Taddei K, Gardener SL, Rainey-Smith SR, Peiffer JJ, ... Dominantly Inherited Alzheimer, N. (2017). Habitual exercise levels are associated with cerebral amyloid load in presymptomatic autosomal dominant Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*, 13(11), 1197–1206. doi:10.1016/j.jalz.2017.03.008
- Brown LS, Foster CG, Courtney JM, King NE, Howells DW, & Sutherland BA (2019). Pericytes and Neurovascular Function in the Healthy and Diseased Brain. *Front Cell Neurosci*, 13, 282. doi:10.3389/fncel.2019.00282 [PubMed: 31316352]
- Chapman SB, Aslan S, Spence JS, Defina LF, Keebler MW, Didehbani N, & Lu H (2013). Shorter term aerobic exercise improves brain, cognition, and cardiovascular fitness in aging. *Frontiers in aging neuroscience*, 5, 75. doi:10.3389/fnagi.2013.00075 [PubMed: 24282403]
- Church DD, Hoffman JR, Mangine GT, Jajtner AR, Townsend JR, Beyer KS, ... Stout JR (2016). Comparison of high-intensity vs. high-volume resistance training on the BDNF response to exercise. *Journal of applied physiology*, 121(1), 123–128. doi:10.1152/jappphysiol.00233.2016 [PubMed: 27231312]
- Cohen BE, Edmondson D, & Kronish IM (2015). State of the Art Review: Depression, Stress, Anxiety, and Cardiovascular Disease. *Am J Hypertens*, 28(11), 1295–1302. doi:10.1093/ajh/hpv047 [PubMed: 25911639]
- Contrepois K, Wu S, Moneghetti KJ, Hornburg D, Ahadi S, Tsai MS, ... Snyder MP (2020). Molecular Choreography of Acute Exercise. *Cell*, 181(5), 1112–1130 e1116. doi:10.1016/j.cell.2020.04.043 [PubMed: 32470399]
- Cook JN, DeVan AE, Schleifer JL, Anton MM, Cortez-Cooper MY, & Tanaka H (2006). Arterial compliance of rowers: implications for combined aerobic and strength training on arterial elasticity. *Am J Physiol Heart Circ Physiol*, 290(4), H1596–1600. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=16284225> [PubMed: 16284225]
- Cortez-Cooper MY, DeVan AE, Anton MM, Farrar RP, Beckwith KA, Todd JS, & Tanaka H (2005). Effects of high intensity resistance training on arterial stiffness and wave reflection in women. *Am J Hypertens*, 18(7), 930–934. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=16053989> [PubMed: 16053989]
- Cotman CW, Berchtold NC, & Christie LA (2007). Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends in neurosciences*, 30(9), 464–472. doi:10.1016/j.tins.2007.06.011 [PubMed: 17765329]
- Coverdale NS, Gati JS, Opalevych O, Perrotta A, & Shoemaker JK (2014). Cerebral blood flow velocity underestimates cerebral blood flow during modest hypercapnia and hypocapnia. *Journal of applied physiology*, 117(10), 1090–1096. doi:10.1152/jappphysiol.00285.2014 [PubMed: 25012027]
- de la Torre JC (2004). Is Alzheimer's disease a neurodegenerative or a vascular disorder? Data, dogma, and dialectics. *Lancet Neurol*, 3(3), 184–190. doi:10.1016/S1474-4422(04)00683-0 [PubMed: 14980533]
- de Leon MJ, Li Y, Okamura N, Tsui WH, Saint-Louis LA, Glodzik L, ... Rusinek H (2017). Cerebrospinal Fluid Clearance in Alzheimer Disease Measured with Dynamic PET. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine*, 58(9), 1471–1476. doi:10.2967/jnumed.116.187211

- de Roos A, van der Grond J, Mitchell G, & Westenberg J (2017). Magnetic Resonance Imaging of Cardiovascular Function and the Brain: Is Dementia a Cardiovascular-Driven Disease? *Circulation*, 135(22), 2178–2195. doi:10.1161/CIRCULATIONAHA.116.021978 [PubMed: 28559496]
- DeSouza CA, Shapiro LF, Clevenger CM, Dinunno FA, Monahan KD, Tanaka H, & Seals DR (2000). Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation*, 102(12), 1351–1357. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10993851> [PubMed: 10993851]
- Ding J, Mitchell GF, Bots ML, Sigurdsson S, Harris TB, Garcia M, ... Launer LJ (2015). Carotid arterial stiffness and risk of incident cerebral microbleeds in older people: the Age, Gene/Environment Susceptibility (AGES)-Reykjavik study. *Arterioscler Thromb Vasc Biol*, 35(8), 1889–1895. doi:10.1161/ATVBAHA.115.305451 [PubMed: 26112009]
- Egan B, & Zierath JR (2013). Exercise metabolism and the molecular regulation of skeletal muscle adaptation. *Cell metabolism*, 17(2), 162–184. doi:10.1016/j.cmet.2012.12.012 [PubMed: 23395166]
- Egashira K, Inou T, Hirooka Y, Kai H, Sugimachi M, Suzuki S, ... Takeshita A (1993). Effects of age on endothelium-dependent vasodilation of resistance coronary artery by acetylcholine in humans. *Circulation*, 88(1), 77–81. doi:10.1161/01.cir.88.1.77 [PubMed: 8319359]
- Elias MF, Robbins MA, Budge MM, Abhayaratna WP, Dore GA, & Elias PK (2009). Arterial pulse wave velocity and cognition with advancing age. *Hypertension*, 53(4), 668–673. doi:10.1161/HYPERTENSIONAHA.108.126342 [PubMed: 19237680]
- Endres M, Gertz K, Lindauer U, Katchanov J, Schultze J, Schrock H, ... Laufs U (2003). Mechanisms of stroke protection by physical activity. *Annals of neurology*, 54(5), 582–590. doi:10.1002/ana.10722 [PubMed: 14595647]
- Erickson KI, Hillman C, Stillman CM, Ballard RM, Bloodgood B, Conroy DE, ... For Physical Activity Guidelines Advisory, C. (2019). Physical Activity, Cognition, and Brain Outcomes: A Review of the 2018 Physical Activity Guidelines. *Med Sci Sports Exerc*, 51(6), 1242–1251. doi:10.1249/MSS.0000000000001936 [PubMed: 31095081]
- Erickson KI, Prakash RS, Voss MW, Chaddock L, Hu L, Morris KS, ... Kramer AF (2009). Aerobic fitness is associated with hippocampal volume in elderly humans. *Hippocampus*, 19(10), 1030–1039. doi:10.1002/hipo.20547 [PubMed: 19123237]
- Erickson KI, Raji CA, Lopez OL, Becker JT, Rosano C, Newman AB, ... Kuller LH (2010). Physical activity predicts gray matter volume in late adulthood: the Cardiovascular Health Study. *Neurology*, 75(16), 1415–1422. doi:10.1212/WNL.0b013e3181f88359 [PubMed: 20944075]
- Fierstra J, Sobczyk O, Battisti-Charbonney A, Mandell DM, Poubanc J, Crawley AP, ... Fisher JA (2013). Measuring cerebrovascular reactivity: what stimulus to use? *J Physiol*, 591(23), 5809–5821. doi:10.1113/jphysiol.2013.259150 [PubMed: 24081155]
- Flodin P, Jonasson LS, Riklund K, Nyberg L, & Boraxbekk CJ (2017). Does Aerobic Exercise Influence Intrinsic Brain Activity? An Aerobic Exercise Intervention among Healthy Old Adults. *Frontiers in aging neuroscience*, 9, 267. doi:10.3389/fnagi.2017.00267 [PubMed: 28848424]
- Furby HV, Warnert EA, Marley CJ, Bailey DM, & Wise RG (2019). Cardiorespiratory fitness is associated with increased middle cerebral arterial compliance and decreased cerebral blood flow in young healthy adults: A pulsed ASL MRI study. *J Cereb Blood Flow Metab*, 271678X19865449. doi:10.1177/0271678X19865449
- Galvin SD, Celi LA, Thomas KN, Clendon TR, Galvin IE, Bunton RW, & Ainslie PN (2010). Effects of age and coronary artery disease on cerebrovascular reactivity to carbon dioxide in humans. *Anaesthesia and intensive care*, 38(4), 710–717. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20715736> [PubMed: 20715736]
- Geda YE, Roberts RO, Knopman DS, Christianson TJ, Pankratz VS, Ivnik RJ, ... Rocca WA (2010). Physical exercise, aging, and mild cognitive impairment: a population-based study. *Arch Neurol*, 67(1), 80–86. doi:10.1001/archneurol.2009.297 [PubMed: 20065133]
- Gerhard M, Roddy MA, Creager SJ, & Creager MA (1996). Aging progressively impairs endothelium-dependent vasodilation in forearm resistance vessels of humans. *Hypertension*, 27(4), 849–853. doi:10.1161/01.hyp.27.4.849 [PubMed: 8613259]

- Green DJ, Hopman MT, Padilla J, Laughlin MH, & Thijssen DH (2017). Vascular Adaptation to Exercise in Humans: Role of Hemodynamic Stimuli. *Physiol Rev*, 97(2), 495–528. doi:10.1152/physrev.00014.2016 [PubMed: 28151424]
- Green DJ, Maiorana A, O'Driscoll G, & Taylor R (2004). Effect of exercise training on endothelium-derived nitric oxide function in humans. *J Physiol*, 561(Pt 1), 1–25. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=15375191> [PubMed: 15375191]
- Guadagni V, Drogos LL, Tyndall AV, Davenport MH, Anderson TJ, Eskes GA, ... Poulin MJ (2020). Aerobic exercise improves cognition and cerebrovascular regulation in older adults. *Neurology*, 94(21), e2245–e2257. doi:10.1212/WNL.0000000000009478 [PubMed: 32404355]
- He XF, Liu DX, Zhang Q, Liang FY, Dai GY, Zeng JS, ... Lan Y (2017). Voluntary Exercise Promotes Glymphatic Clearance of Amyloid Beta and Reduces the Activation of Astrocytes and Microglia in Aged Mice. *Front Mol Neurosci*, 10, 144. doi:10.3389/fnmol.2017.00144 [PubMed: 28579942]
- Hoiland RL, Fisher JA, & Ainslie PN (2019). Regulation of the Cerebral Circulation by Arterial Carbon Dioxide. *Comprehensive Physiology*, 9(3), 1101–1154. doi:10.1002/cphy.c180021 [PubMed: 31187899]
- Iadecola C (2004). Neurovascular regulation in the normal brain and in Alzheimer's disease. *Nature reviews. Neuroscience*, 5(5), 347–360. doi:10.1038/nrn1387 [PubMed: 15100718]
- Iadecola C (2017). The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease. *Neuron*, 96(1), 17–42. doi:10.1016/j.neuron.2017.07.030 [PubMed: 28957666]
- Iliff JJ, & Nedergaard M (2013). Is there a cerebral lymphatic system? *Stroke*, 44(6 Suppl 1), S93–95. doi:10.1161/STROKEAHA.112.678698 [PubMed: 23709744]
- Intzandt B, Sabra D, Foster C, Desjardins-Crepeau L, Hoge RD, Steele CJ, ... Gauthier CJ (2020). Higher cardiovascular fitness level is associated with lower cerebrovascular reactivity and perfusion in healthy older adults. *J Cereb Blood Flow Metab*, 40(7), 1468–1481. doi:10.1177/0271678X19862873 [PubMed: 31342831]
- Iturria-Medina Y, Sotero RC, Toussaint PJ, Mateos-Perez JM, Evans AC, & Alzheimer's Disease Neuroimaging I (2016). Early role of vascular dysregulation on late-onset Alzheimer's disease based on multifactorial data-driven analysis. *Nat Commun*, 7, 11934. doi:10.1038/ncomms11934 [PubMed: 27327500]
- Jaruchart T, Suwanwela NC, Tanaka H, & Suksom D (2016). Arterial stiffness is associated with age-related differences in cerebrovascular conductance. *Experimental gerontology*, 73, 59–64. doi:10.1016/j.exger.2015.11.006 [PubMed: 26571202]
- Jiang H, Jia D, Zhang B, Yang W, Dong Z, Sun X, ... Ge J (2020). Exercise improves cardiac function and glucose metabolism in mice with experimental myocardial infarction through inhibiting HDAC4 and upregulating GLUT1 expression. *Basic Res Cardiol*, 115(3), 28. doi:10.1007/s00395-020-0787-1 [PubMed: 32236769]
- Kalaria RN, & Harik SI (1989). Reduced glucose transporter at the blood-brain barrier and in cerebral cortex in Alzheimer disease. *Journal of neurochemistry*, 53(4), 1083–1088. doi:10.1111/j.1471-4159.1989.tb07399.x [PubMed: 2769254]
- Kastrup A, Dichgans J, Niemeier M, & Schabet M (1998). Changes of cerebrovascular CO₂ reactivity during normal aging. *Stroke*, 29(7), 1311–1314. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=9660378> [PubMed: 9660378]
- Kearney-Schwartz A, Rossignol P, Bracard S, Felblinger J, Fay R, Boivin JM, ... Zannad F (2009). Vascular structure and function is correlated to cognitive performance and white matter hyperintensities in older hypertensive patients with subjective memory complaints. *Stroke; a journal of cerebral circulation*, 40(4), 1229–1236. doi:10.1161/STROKEAHA.108.532853
- Kisler K, Nelson AR, Montagne A, & Zlokovic BV (2017). Cerebral blood flow regulation and neurovascular dysfunction in Alzheimer disease. *Nature reviews. Neuroscience*, 18(7), 419–434. doi:10.1038/nrn.2017.48 [PubMed: 28515434]
- Kleinloog JPD, Mensink RP, Ivanov D, Adam JJ, Uludag K, & Joris PJ (2019). Aerobic Exercise Training Improves Cerebral Blood Flow and Executive Function: A Randomized, Controlled

- Cross-Over Trial in Sedentary Older Men. *Frontiers in aging neuroscience*, 11, 333. doi:10.3389/fnagi.2019.00333 [PubMed: 31866855]
- Kohn JC, Lampi MC, & Reinhart-King CA (2015). Age-related vascular stiffening: causes and consequences. *Frontiers in genetics*, 6, 112. doi:10.3389/fgene.2015.00112 [PubMed: 25926844]
- Kress BT, Iliff JJ, Xia M, Wang M, Wei HS, Zeppenfeld D, ... Nedergaard M (2014). Impairment of paravascular clearance pathways in the aging brain. *Annals of neurology*, 76(6), 845–861. doi:10.1002/ana.24271 [PubMed: 25204284]
- Kuring JK, Mathias JL, & Ward L (2018). Prevalence of Depression, Anxiety and PTSD in People with Dementia: a Systematic Review and Meta-Analysis. *Neuropsychol Rev*, 28(4), 393–416. doi:10.1007/s11065-018-9396-2 [PubMed: 30536144]
- Laing KK, Simoes S, Baena-Caldas GP, Lao PJ, Kothiyi M, Igwe KC, ... Alzheimer's Disease Neuroimaging, I. (2020). Cerebrovascular disease promotes tau pathology in Alzheimer's disease. *Brain Commun*, 2(2), fcaa132. doi:10.1093/braincomms/fcaa132 [PubMed: 33215083]
- Latimer CS, Searcy JL, Bridges MT, Brewer LD, Popovic J, Blalock EM, ... Porter NM (2011). Reversal of glial and neurovascular markers of unhealthy brain aging by exercise in middle-aged female mice. *PloS one*, 6(10), e26812. doi:10.1371/journal.pone.0026812 [PubMed: 22046366]
- Lee HY, & Oh BH (2010). Aging and arterial stiffness. *Circulation journal : official journal of the Japanese Circulation Society*, 74(11), 2257–2262. doi:10.1253/circj.cj-10-0910 [PubMed: 20962429]
- Liang KY, Mintun MA, Fagan AM, Goate AM, Bugg JM, Holtzman DM, ... Head D (2010). Exercise and Alzheimer's disease biomarkers in cognitively normal older adults. *Annals of neurology*, 68(3), 311–318. doi:10.1002/ana.22096 [PubMed: 20818789]
- Liu PZ, & Nusslock R (2018). Exercise-Mediated Neurogenesis in the Hippocampus via BDNF. *Front Neurosci*, 12, 52. doi:10.3389/fnins.2018.00052 [PubMed: 29467613]
- Lu H, Xu F, Rodrigue KM, Kennedy KM, Cheng Y, Flicker B, ... Park DC (2011). Alterations in cerebral metabolic rate and blood supply across the adult lifespan. *Cerebral cortex*, 21(6), 1426–1434. doi:10.1093/cercor/bhq224 [PubMed: 21051551]
- Madureira J, Castro P, & Azevedo E (2017). Demographic and Systemic Hemodynamic Influences in Mechanisms of Cerebrovascular Regulation in Healthy Adults. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*, 26(3), 500–508. doi:10.1016/j.jstrokecerebrovasdis.2016.12.003 [PubMed: 28038898]
- McEniery CM, Wallace S, Mackenzie IS, McDonnell B, Yasmin, Newby DE, ... Wilkinson IB (2006). Endothelial function is associated with pulse pressure, pulse wave velocity, and augmentation index in healthy humans. *Hypertension*, 48(4), 602–608. doi:10.1161/01.HYP.0000239206.64270.5f [PubMed: 16940223]
- Mestre H, Tithof J, Du T, Song W, Peng W, Sweeney AM, ... Kelley DH (2018). Flow of cerebrospinal fluid is driven by arterial pulsations and is reduced in hypertension. *Nat Commun*, 9(1), 4878. doi:10.1038/s41467-018-07318-3 [PubMed: 30451853]
- Mikhail Kellawan J, Harrell JW, Schrauben EM, Hoffman CA, Roldan-Alzate A, Schrage WG, & Wieben O (2016). Quantitative cerebrovascular 4D flow MRI at rest and during hypercapnia challenge. *Magn Reson Imaging*, 34(4), 422–428. doi:10.1016/j.mri.2015.12.016 [PubMed: 26708027]
- Miller KB, Howery AJ, Harvey RE, Eldridge MW, & Barnes JN (2018). Cerebrovascular Reactivity and Central Arterial Stiffness in Habitually Exercising Healthy Adults. *Frontiers in physiology*, 9, 1096. doi:10.3389/fphys.2018.01096 [PubMed: 30174609]
- Miller KB, Howery AJ, Rivera-Rivera LA, Johnson SC, Rowley HA, Wieben O, & Barnes JN (2019). Age-Related Reductions in Cerebrovascular Reactivity Using 4D Flow MRI. *Frontiers in aging neuroscience*, 11, 281. doi:10.3389/fnagi.2019.00281 [PubMed: 31680935]
- Mitchell GF (2008). Effects of central arterial aging on the structure and function of the peripheral vasculature: implications for end-organ damage. *Journal of applied physiology*, 105(5), 1652–1660. doi:10.1152/jappphysiol.90549.2008 [PubMed: 18772322]
- Mitchell GF, van Buchem MA, Sigurdsson S, Gotlib JD, Jonsdottir MK, Kjartansson O, ... Launer LJ (2011). Arterial stiffness, pressure and flow pulsatility and brain structure and function: the

- Age, Gene/Environment Susceptibility--Reykjavik study. *Brain : a journal of neurology*, 134(Pt 11), 3398–3407. doi:10.1093/brain/awr253 [PubMed: 22075523]
- Miyachi M (2013). Effects of resistance training on arterial stiffness: a meta-analysis. *Br J Sports Med*, 47(6), 393–396. doi:10.1136/bjsports-2012-090488 [PubMed: 22267567]
- Miyachi M, Donato AJ, Yamamoto K, Takahashi K, Gates PE, Moreau KL, & Tanaka H (2003). Greater age-related reductions in central arterial compliance in resistance-trained men. *Hypertension*, 41(1), 130–135. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/12511542> [PubMed: 12511542]
- Miyachi M, Kawano H, Sugawara J, Takahashi K, Hayashi K, Yamazaki K, ... Tanaka H (2004). Unfavorable effects of resistance training on central arterial compliance: a randomized intervention study. *Circulation*, 110(18), 2858–2863. doi:10.1161/01.CIR.0000146380.08401.99 [PubMed: 15492301]
- Montagne A, Nation DA, Pa J, Sweeney MD, Toga AW, & Zlokovic BV (2016). Brain imaging of neurovascular dysfunction in Alzheimer's disease. *Acta neuropathologica*, 131(5), 687–707. doi:10.1007/s00401-016-1570-0 [PubMed: 27038189]
- Muhire G, Iulita MF, Vallerand D, Youwakim J, Gratuze M, Petry FR, ... Girouard H (2019). Arterial Stiffness Due to Carotid Calcification Disrupts Cerebral Blood Flow Regulation and Leads to Cognitive Deficits. *J Am Heart Assoc*, 8(9), e011630. doi:10.1161/JAHA.118.011630 [PubMed: 31057061]
- Murrell CJ, Cotter JD, Thomas KN, Lucas SJ, Williams MJ, & Ainslie PN (2012). Cerebral blood flow and cerebrovascular reactivity at rest and during sub-maximal exercise: Effect of age and 12-week exercise training. *Age*. doi:10.1007/s11357-012-9414-x
- Nicholson WT, Vaa B, Hesse C, Eisenach JH, & Joyner MJ (2009). Aging is associated with reduced prostacyclin-mediated dilation in the human forearm. *Hypertension*, 53(6), 973–978. doi:10.1161/HYPERTENSIONAHA.108.121483 [PubMed: 19414649]
- Nigam A, Mitchell GF, Lambert J, & Tardif JC (2003). Relation between conduit vessel stiffness (assessed by tonometry) and endothelial function (assessed by flow-mediated dilatation) in patients with and without coronary heart disease. *Am J Cardiol*, 92(4), 395–399. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=12914868> [PubMed: 12914868]
- Nualnim N, Barnes JN, Tarumi T, Renzi CP, & Tanaka H (2011). Comparison of central artery elasticity in swimmers, runners, and the sedentary. *Am J Cardiol*, 107(5), 783–787. doi:10.1016/j.amjcard.2010.10.062 [PubMed: 21247521]
- O'Keefe EL, O'Keefe JH, & Lavie CJ (2019). Exercise Counteracts the Cardiotoxicity of Psychosocial Stress. *Mayo Clinic proceedings*. *Mayo Clinic*, 94(9), 1852–1864. doi:10.1016/j.mayocp.2019.02.022
- O'Rourke MF, & Hashimoto J (2007). Mechanical factors in arterial aging: a clinical perspective. *J Am Coll Cardiol*, 50(1), 1–13. doi:10.1016/j.jacc.2006.12.050 [PubMed: 17601538]
- Palta P, Sharrett AR, Wei J, Meyer ML, Kucharska-Newton A, Power MC, ... Heiss G (2019). Central Arterial Stiffness Is Associated With Structural Brain Damage and Poorer Cognitive Performance: The ARIC Study. *J Am Heart Assoc*, 8(2), e011045. doi:10.1161/JAHA.118.011045 [PubMed: 30646799]
- Pang R, Wang X, Pei F, Zhang W, Shen J, Gao X, & Chang C (2019). Regular Exercise Enhances Cognitive Function and Intracerebral GLUT Expression in Alzheimer's Disease Model Mice. *Journal of Alzheimer's disease : JAD*, 72(1), 83–96. doi:10.3233/JAD-190328 [PubMed: 31561359]
- Pase MP, Grima NA, Stough C, Scholey A, & Pipingas A (2014). Association of pulsatile and mean cerebral blood flow velocity with age and neuropsychological performance. *Physiology & behavior*, 130, 23–27. doi:10.1016/j.physbeh.2014.03.015 [PubMed: 24657738]
- Pedersen BK, & Hoffman-Goetz L (2000). Exercise and the immune system: regulation, integration, and adaptation. *Physiol Rev*, 80(3), 1055–1081. doi:10.1152/physrev.2000.80.3.1055 [PubMed: 10893431]

- Pialoux V, Brown AD, Leigh R, Friedenreich CM, & Poulin MJ (2009). Effect of cardiorespiratory fitness on vascular regulation and oxidative stress in postmenopausal women. *Hypertension*, 54(5), 1014–1020. doi:10.1161/HYPERTENSIONAHA.109.138917 [PubMed: 19786647]
- Rasmussen MK, Mestre H, & Nedergaard M (2018). The glymphatic pathway in neurological disorders. *Lancet Neurol*, 17(11), 1016–1024. doi:10.1016/S1474-4422(18)30318-1 [PubMed: 30353860]
- Rasmussen P, Brassard P, Adser H, Pedersen MV, Leick L, Hart E, ... Pilegaard H (2009). Evidence for a release of brain-derived neurotrophic factor from the brain during exercise. *Experimental physiology*, 94(10), 1062–1069. doi:10.1113/expphysiol.2009.048512 [PubMed: 19666694]
- Ringstad G, Valnes LM, Dale AM, Pripp AH, Vatnehol SS, Emblem KE, ... Eide PK (2018). Brain-wide glymphatic enhancement and clearance in humans assessed with MRI. *JCI Insight*, 3(13). doi:10.1172/jci.insight.121537
- Rivera-Rivera LA, Turski P, Johnson KM, Hoffman C, Berman SE, Kilgas P, ... Wieben O (2016). 4D flow MRI for intracranial hemodynamics assessment in Alzheimer's disease. *J Cereb Blood Flow Metab*, 36(10), 1718–1730. doi:10.1177/0271678X15617171 [PubMed: 26661239]
- Robertson AD, Atwi S, Kostoglou K, Verhoeff N, Oh PI, Mitsis GD, ... MacIntosh BJ (2019). Cerebrovascular Pulsatility During Rest and Exercise Reflects Hemodynamic Impairment in Stroke and Cerebral Small Vessel Disease. *Ultrasound in medicine & biology*, 45(12), 3116–3127. doi:10.1016/j.ultrasmedbio.2019.08.019 [PubMed: 31570171]
- Robertson AD, Tessmer CF, & Hughson RL (2010). Association between arterial stiffness and cerebrovascular resistance in the elderly. *J Hum Hypertens*, 24(3), 190–196. doi:10.1038/jhh.2009.56 [PubMed: 19571826]
- Sadekova N, Vallerand D, Guevara E, Lesage F, & Girouard H (2013). Carotid calcification in mice: a new model to study the effects of arterial stiffness on the brain. *J Am Heart Assoc*, 2(3), e000224. doi:10.1161/JAHA.113.000224 [PubMed: 23782921]
- Sargurupremraj M, Suzuki H, Jian X, Sarnowski C, Evans TE, Bis JC, ... Debette S (2020). Cerebral small vessel disease genomics and its implications across the lifespan. *Nat Commun*, 11(1), 6285. doi:10.1038/s41467-020-19111-2 [PubMed: 33293549]
- Schrauben EM, Johnson KM, Huston J, Del Rio AM, Reeder SB, Field A, & Wieben O (2014). Reproducibility of cerebrospinal venous blood flow and vessel anatomy with the use of phase contrast-vastly undersampled isotropic projection reconstruction and contrast-enhanced MRA. *AJNR. American journal of neuroradiology*, 35(5), 999–1006. doi:10.3174/ajnr.A3779 [PubMed: 24287088]
- Seals DR, Jablonski KL, & Donato AJ (2011). Aging and vascular endothelial function in humans. *Clin Sci (Lond)*, 120(9), 357–375. doi:10.1042/CS20100476 [PubMed: 21244363]
- Severinsen MCK, & Pedersen BK (2020). Muscle-Organ Crosstalk: The Emerging Roles of Myokines. *Endocr Rev*, 41(4). doi:10.1210/endrev/bnaa016
- Shi Y, Thrippleton MJ, Makin SD, Marshall I, Geerlings MI, de Craen AJM, ... Wardlaw JM (2016). Cerebral blood flow in small vessel disease: A systematic review and meta-analysis. *J Cereb Blood Flow Metab*, 36(10), 1653–1667. doi:10.1177/0271678X16662891 [PubMed: 27496552]
- Shibata S, Fujimoto N, Hastings JL, Carrick-Ranson G, Bhella PS, Hearon CM Jr., & Levine BD (2018). The effect of lifelong exercise frequency on arterial stiffness. *J Physiol*, 596(14), 2783–2795. doi:10.1113/JP275301 [PubMed: 29781119]
- Silvestrini M, Pasqualetti P, Baruffaldi R, Bartolini M, Handouk Y, Matteis M, ... Vernieri F (2006). Cerebrovascular reactivity and cognitive decline in patients with Alzheimer disease. *Stroke; a journal of cerebral circulation*, 37(4), 1010–1015. doi:10.1161/01.STR.0000206439.62025.97
- Sofi F, Valecchi D, Bacci D, Abbate R, Gensini GF, Casini A, & Macchi C (2011). Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J Intern Med*, 269(1), 107–117. doi:10.1111/j.1365-2796.2010.02281.x [PubMed: 20831630]
- Steen Jensen C, Portelius E, Siersma V, Høgh P, Wermuth L, Blennow K, ... Hviid Simonsen A (2016). Cerebrospinal Fluid Amyloid Beta and Tau Concentrations Are Not Modulated by 16 Weeks of Moderate- to High-Intensity Physical Exercise in Patients with Alzheimer Disease. *Dement Geriatr Cogn Disord*, 42(3-4), 146–158. doi:10.1159/000449408 [PubMed: 27643858]

- Stillman CM, Esteban-Cornejo I, Brown B, Bender CM, & Erickson KI (2020). Effects of Exercise on Brain and Cognition Across Age Groups and Health States. *Trends in neurosciences*, 43(7), 533–543. doi:10.1016/j.tins.2020.04.010 [PubMed: 32409017]
- Sun Z (2015). Aging, arterial stiffness, and hypertension. *Hypertension*, 65(2), 252–256. doi:10.1161/HYPERTENSIONAHA.114.03617 [PubMed: 25368028]
- Sweeney MD, Kisler K, Montagne A, Toga AW, & Zlokovic BV (2018). The role of brain vasculature in neurodegenerative disorders. *Nature neuroscience*, 21(10), 1318–1331. doi:10.1038/s41593-018-0234-x [PubMed: 30250261]
- Szablewski L (2017). Glucose Transporters in Brain: In Health and in Alzheimer's Disease. *Journal of Alzheimer's disease : JAD*, 55(4), 1307–1320. doi:10.3233/JAD-160841 [PubMed: 27858715]
- Taddei S, Virdis A, Ghiadoni L, Salvetti G, Bernini G, Magagna A, & Salvetti A (2001). Age-related reduction of NO availability and oxidative stress in humans. *Hypertension*, 38(2), 274–279. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=11509489> [PubMed: 11509489]
- Takimoto M, & Hamada T (2014). Acute exercise increases brain region-specific expression of MCT1, MCT2, MCT4, GLUT1, and COX IV proteins. *Journal of applied physiology*, 116(9), 1238–1250. doi:10.1152/japplphysiol.01288.2013 [PubMed: 24610532]
- Tan CH, Low KA, Kong T, Fletcher MA, Zimmerman B, Maclin EL, ... Fabiani M (2017). Mapping cerebral pulse pressure and arterial compliance over the adult lifespan with optical imaging. *PloS one*, 12(2), e0171305. doi:10.1371/journal.pone.0171305 [PubMed: 28234912]
- Tanaka H (2019). Antiaging Effects of Aerobic Exercise on Systemic Arteries. *Hypertension, HYPERTENSIONAHA*11913179. doi:10.1161/HYPERTENSIONAHA.119.13179
- Tanaka H, DeSouza CA, & Seals DR (1998). Absence of age-related increase in central arterial stiffness in physically active women. *Arterioscler Thromb Vasc Biol*, 18(1), 127–132. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=9445266> [PubMed: 9445266]
- Tanaka H, Dinunno FA, Monahan KD, Clevenger CM, DeSouza CA, & Seals DR (2000). Aging, habitual exercise, and dynamic arterial compliance. *Circulation*, 102(11), 1270–1275. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=10982542> [PubMed: 10982542]
- Tanaka H, Dinunno FA, Monahan KD, DeSouza CA, & Seals DR (2001). Carotid artery wall hypertrophy with age is related to local systolic blood pressure in healthy men. *Arterioscler Thromb Vasc Biol*, 21(1), 82–87. Retrieved from <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&listuids=11145937> [PubMed: 11145937]
- Tarasoff-Conway JM, Carare RO, Osorio RS, Glodzik L, Butler T, Fieremans E, ... de Leon MJ (2015). Clearance systems in the brain-implications for Alzheimer disease. *Nature reviews. Neurology*, 11(8), 457–470. doi:10.1038/nrneurol.2015.119 [PubMed: 26195256]
- Thijssen DH, Carter SE, & Green DJ (2016). Arterial structure and function in vascular ageing: are you as old as your arteries? *J Physiol*, 594(8), 2275–2284. doi:10.1113/JP270597 [PubMed: 26140618]
- Thijssen DH, Rongen GA, van Dijk A, Smits P, & Hopman MT (2007). Enhanced endothelin-1-mediated leg vascular tone in healthy older subjects. *Journal of applied physiology*, 103(3), 852–857. doi:10.1152/japplphysiol.00357.2007 [PubMed: 17556493]
- Thomas BP, Yezhuvath US, Tseng BY, Liu P, Levine BD, Zhang R, & Lu H (2013). Life-long aerobic exercise preserved baseline cerebral blood flow but reduced vascular reactivity to CO₂. *Journal of magnetic resonance imaging : JMRI*, 38(5), 1177–1183. doi:10.1002/jmri.24090 [PubMed: 23526811]
- Thorin-Trescases N, de Montgolfier O, Pincon A, Raignault A, Caland L, Labbe P, & Thorin E (2018). The impact of pulse pressure on cerebrovascular events leading to age-related cognitive decline. *Am J Physiol Heart Circ Physiol*. doi:10.1152/ajpheart.00637.2017
- Toledo JB, Arnold SE, Raible K, Brettschneider J, Xie SX, Grossman M, ... Trojanowski JQ (2013). Contribution of cerebrovascular disease in autopsy confirmed neurodegenerative disease cases in the National Alzheimer's Coordinating Centre. *Brain : a journal of neurology*, 136(Pt 9), 2697–2706. doi:10.1093/brain/awt188 [PubMed: 23842566]

- Tomoto T, Riley J, Turner M, Zhang R, & Tarumi T (2020). Cerebral vasomotor reactivity during hypo- and hypercapnia across the adult lifespan. *J Cereb Blood Flow Metab*, 40(3), 600–610. doi:10.1177/0271678X19828327 [PubMed: 30764704]
- Tymko MM, Ainslie PN, & Smith KJ (2018). Evaluating the methods used for measuring cerebral blood flow at rest and during exercise in humans. *Eur J Appl Physiol*, 118(8), 1527–1538. doi:10.1007/s00421-018-3887-y [PubMed: 29767351]
- Tzeng YC, Ainslie PN, Cooke WH, Peebles KC, Willie CK, MacRae BA, ... Rickards CA (2012). Assessment of cerebral autoregulation: the quandary of quantification. *American journal of physiology Heart and circulatory physiology*, 303(6), H658–671. Retrieved from <Go to ISI>://MEDLINE:22821992 [PubMed: 22821992]
- Vaitkevicius PV, Fleg JL, Engel JH, O'Connor FC, Wright JG, Lakatta LE, ... Lakatta EG (1993). Effects of age and aerobic capacity on arterial stiffness in healthy adults. *Circulation*, 88(4 Pt 1), 1456–1462. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8403292> [PubMed: 8403292]
- van der Kleij LA, Petersen ET, Siebner HR, Hendrikse J, Frederiksen KS, Sobol NA, ... Garde E (2018). The effect of physical exercise on cerebral blood flow in Alzheimer's disease. *Neuroimage Clin*, 20, 650–654. doi:10.1016/j.nicl.2018.09.003 [PubMed: 30211001]
- van Praag H, Christie BR, Sejnowski TJ, & Gage FH (1999). Running enhances neurogenesis, learning, and long-term potentiation in mice. *Proceedings of the National Academy of Sciences of the United States of America*, 96(23), 13427–13431. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10557337> [PubMed: 10557337]
- van Praag H, Fleshner M, Schwartz MW, & Mattson MP (2014). Exercise, energy intake, glucose homeostasis, and the brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 34(46), 15139–15149. doi:10.1523/JNEUROSCI.2814-14.2014 [PubMed: 25392482]
- van Praag H, Kempermann G, & Gage FH (1999). Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nature neuroscience*, 2(3), 266–270. doi:10.1038/6368 [PubMed: 10195220]
- van Sloten TT, Protogerou AD, Henry RM, Schram MT, Launer LJ, & Stehouwer CD (2015). Association between arterial stiffness, cerebral small vessel disease and cognitive impairment: A systematic review and meta-analysis. *Neuroscience and biobehavioral reviews*, 53, 121–130. doi:10.1016/j.neubiorev.2015.03.011 [PubMed: 25827412]
- Verbree J, Bronzwaer AS, Ghariq E, Versluis MJ, Daemen MJ, van Buchem MA, ... van Osch MJ (2014). Assessment of middle cerebral artery diameter during hypocapnia and hypercapnia in humans using ultra-high-field MRI. *Journal of applied physiology*, 117(10), 1084–1089. doi:10.1152/japplphysiol.00651.2014 [PubMed: 25190741]
- Veys K, Fan Z, Ghobrial M, Bouche A, Garcia-Caballero M, Vriens K, ... De Bock K (2020). Role of the GLUT1 Glucose Transporter in Postnatal CNS Angiogenesis and Blood-Brain Barrier Integrity. *Circ Res*, 127(4), 466–482. doi:10.1161/CIRCRESAHA.119.316463 [PubMed: 32404031]
- Vivar C, & van Praag H (2017). Running Changes the Brain: the Long and the Short of It. *Physiology (Bethesda)*, 32(6), 410–424. doi:10.1152/physiol.00017.2017 [PubMed: 29021361]
- von Holstein-Rathlou S, Petersen NC, & Nedergaard M (2018). Voluntary running enhances glymphatic influx in awake behaving, young mice. *Neuroscience letters*, 662, 253–258. doi:10.1016/j.neulet.2017.10.035 [PubMed: 29079431]
- Walker AE, Henson GD, Reihl KD, Morgan RG, Dobson PS, Nielson EI, ... Donato AJ (2015). Greater impairments in cerebral artery compared with skeletal muscle feed artery endothelial function in a mouse model of increased large artery stiffness. *J Physiol*, 593(8), 1931–1943. doi:10.1113/jphysiol.2014.285338 [PubMed: 25627876]
- Watson NL, Sutton-Tyrrell K, Rosano C, Boudreau RM, Hardy SE, Simonsick EM, ... Newman AB (2011). Arterial stiffness and cognitive decline in well-functioning older adults. *J Gerontol A Biol Sci Med Sci*, 66(12), 1336–1342. doi:10.1093/gerona/glr119 [PubMed: 21768503]
- Webb AJ, Simoni M, Mazzucco S, Kuker W, Schulz U, & Rothwell PM (2012). Increased cerebral arterial pulsatility in patients with leukoaraiosis: arterial stiffness enhances transmission of aortic pulsatility. *Stroke*, 43(10), 2631–2636. doi:10.1161/STROKEAHA.112.655837 [PubMed: 22923446]

- Whyte JJ, & Laughlin MH (2010). The effects of acute and chronic exercise on the vasculature. *Acta Physiol (Oxf)*, 199(4), 441–450. doi:10.1111/j.1748-1716.2010.02127.x [PubMed: 20353494]
- Winkler EA, Nishida Y, Sagare AP, Rege SV, Bell RD, Perlmutter D, ... Zlokovic BV (2015). GLUT1 reductions exacerbate Alzheimer's disease vasculo-neuronal dysfunction and degeneration. *Nature neuroscience*, 18(4), 521–530. doi:10.1038/nn.3966 [PubMed: 25730668]
- Wolters FJ, Zonneveld HI, Hofman A, van der Lugt A, Koudstaal PJ, Vernooij MW, ... Heart-Brain Connection Collaborative Research, G. (2017). Cerebral Perfusion and the Risk of Dementia: A Population-Based Study. *Circulation*, 136(8), 719–728. doi:10.1161/CIRCULATIONAHA.117.027448 [PubMed: 28588075]
- Xu TY, Staessen JA, Wei FF, Xu J, Li FH, Fan WX, ... Li Y (2012). Blood flow pattern in the middle cerebral artery in relation to indices of arterial stiffness in the systemic circulation. *Am J Hypertens*, 25(3), 319–324. doi:10.1038/ajh.2011.223 [PubMed: 22113170]
- Xu X, Wang B, Ren C, Hu J, Greenberg DA, Chen T, ... Jin K (2017). Age-related Impairment of Vascular Structure and Functions. *Aging Dis*, 8(5), 590–610. doi:10.14336/AD.2017.0430 [PubMed: 28966804]
- Yang T, Sun Y, Lu Z, Leak RK, & Zhang F (2017). The impact of cerebrovascular aging on vascular cognitive impairment and dementia. *Ageing Res Rev*, 34, 15–29. doi:10.1016/j.arr.2016.09.007 [PubMed: 27693240]
- Yoshizawa M, Maeda S, Miyaki A, Misono M, Saito Y, Tanabe K, ... Ajisaka R (2009). Effect of 12 weeks of moderate-intensity resistance training on arterial stiffness: a randomised controlled trial in women aged 32-59 years. *Br J Sports Med*, 43(8), 615–618. doi:10.1136/bjsm.2008.052126 [PubMed: 18927168]
- Zeller K, Rahner-Welsch S, & Kuschinsky W (1997). Distribution of Glut1 glucose transporters in different brain structures compared to glucose utilization and capillary density of adult rat brains. *J Cereb Blood Flow Metab*, 17(2), 204–209. doi:10.1097/00004647-199702000-00010 [PubMed: 9040500]
- Zhai FF, Ye YC, Chen SY, Ding FM, Han F, Yang XL, ... Zhu YC (2018). Arterial Stiffness and Cerebral Small Vessel Disease. *Front Neurol*, 9, 723. doi:10.3389/fneur.2018.00723 [PubMed: 30210443]
- Zhao Z, Nelson AR, Betsholtz C, & Zlokovic BV (2015). Establishment and Dysfunction of the Blood-Brain Barrier. *Cell*, 163(5), 1064–1078. doi:10.1016/j.cell.2015.10.067 [PubMed: 26590417]
- Zhu YS, Tarumi T, Tseng BY, Palmer DM, Levine BD, & Zhang R (2013). Cerebral vasomotor reactivity during hypo- and hypercapnia in sedentary elderly and Masters athletes. *J Cereb Blood Flow Metab*, 33(8), 1190–1196. doi:10.1038/jcbfm.2013.66 [PubMed: 23591649]
- Zlokovic BV (2011). Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nature reviews. Neuroscience*, 12(12), 723–738. doi:10.1038/nrn3114 [PubMed: 22048062]

Figure.

Conceptual summary of how regular vigorous aerobic exercise may impact the central arterial stiffness, cerebral vascular function, and cerebral perfusion, with implications for future risk of Alzheimer's disease and other dementias. This figure was created in part with modified Servier Medical Art templates, which are licensed under a Creative Commons Attribution 3.0 Unported License: <https://smart.servier.com>.