Analysis of 200,000 Exome-Sequenced UK Biobank Subjects Implicates Genes Involved in Increased and Decreased Risk of Hypertension

David Curtis^{a, b}^aUCL Genetics Institute, University College London, London, UK; ^bCentre for Psychiatry, Queen Mary University of London, London, UK

Keywords

Hypertension · Biobank · Exome · *DNMT3A* · *GUCY1A1* · *GUCY1B1*

Abstract

Background: Previous analyses have identified common variants along with some specific genes and rare variants which are associated with risk of hypertension, but much remains to be discovered. **Methods and Results:** Exome-sequenced UK Biobank participants were phenotyped based on having a diagnosis of hypertension or taking anti-hypertensive medication to produce a sample of 66,123 cases and 134,504 controls. Variants with minor allele frequency (MAF) <0.01 were subjected to a gene-wise weighted burden analysis, with higher weights assigned to variants which are rarer and/or predicted to have more severe effects. Of 20,384 genes analysed, 2 genes were exome-wide significant, *DNMT3A* and *FES*. Also strongly implicated were *GUCY1A1* and *GUCY1B1*, which code for the subunits of soluble guanylate cyclase. There was further support for the previously reported effects of variants in *NPR1* and protective effects of variants in *DBH*. An inframe deletion in *CACNA1D* with MAF = 0.005, rs72556363, is associated with modestly increased risk of hypertension. Other biologically plausible genes highlighted consist of *CSK*, *AGTR1*, *ZYX*, and *PREP*. All variants implicated were rare, and cumulatively they are not predicted to make a large contribution to the population risk of hyper-

tension. **Conclusions:** This approach confirms and clarifies previously reported findings and also offers novel insights into biological processes influencing hypertension risk, potentially facilitating the development of improved therapeutic interventions. This research has been conducted using the UK Biobank Resource.

© 2021 The Author(s).

Published by S. Karger AG, Basel

Introduction

Hypertension is an important risk factor for disease that has a heritable component, and a recent large genome-wide association study of common variant effects identified 901 loci with enrichment in relevant tissues (blood vessels, heart, adrenal tissue, and adipose tissue) and pathways (angiotensinogen, calcium channels, progesterone, natriuretic peptide receptor, angiotensin-converting enzyme, angiotensin receptors, and endothelin receptors) [1]. Selection pressures tend to mean that common variants individually have small effect sizes and it can be difficult to interpret their biological effects [2]. By contrast, rare variants can potentially have large effect sizes and clear biological mechanisms, as exemplified by a number of monogenic causes of hypertension such as congenital adrenal hyperplasia, familial hyperaldosteronism and pseudohypoaldosteronism, which can be caused by variants in *CYP11B1*, *CYP11B2*, *WNK1*, *WNK4*,

KLHL3, *CUL3*, *SCNN1B*, *SCNN1G*, *CYP17A1*, *HSD11B2*, *NR3C2*, and *KCNJ5* [3]. Additionally, variants in *CACNA1H*, *CACNA1D*, and *CLCN2* have now also been identified as causes of familial hyperaldosteronism, while somatic mutations in *ATP1A1* or *ATP2B3* can produce aldosterone-producing adrenal adenomas with consequent hypertension [4]. Recessively acting variants in *GUCY1A1* (previously labelled *GUCY1A3*) can cause moyamoya disease, and 2 unrelated subjects with moyamoya disease who also had achalasia and hypertension were found to have compound heterozygote variants in this gene [5]. A more recent study has shown that moyamoya disease is itself a risk factor for hypertension [6]. Using hypertension or blood pressure as the phenotype, gene-based analyses aggregating rare, non-synonymous variants implicated *PTMT1*, *DBH* and *NPR1* in a large meta-analysis and also showed that the minor allele of a rare non-synonymous variant in *DBH*, rs3025380, was associated with lower blood pressure [7]. Another study reported that 3 individual non-synonymous variants in *NPR1* were associated with increased (rs35479618 and rs116245325) and decreased (rs61757359) blood pressure and showed that this could be explained by the effects of these variants on guanylate cyclase activity [8]. *KDM1A* codes for LSD1, which removes methyl groups from the methylated lysine 4 residue of histone 3 (H3K4), and there have been reports of association between variants in *KDM1A* and salt-sensitive hypertension in humans, while heterozygous *Lsd1* knock-out mice have salt-sensitive hypertension with increased aldosterone production [9–11].

The growing availability of sequence data means that it may become possible to study the wider effects of rare functional variants in the general population. This may implicate novel genes or may demonstrate a wider role for genes already implicated in severe familial disorders. Exome sequence data is now available for 200,000 of the 500,000 UK Biobank subjects [12]. We have recently analysed this in order to illuminate the effect of rare coding variants on susceptibility to hyperlipidaemia and a number of other common traits with complex inheritance, and we now apply the same approach to study the contribution of rare variants to risk of developing hypertension [13].

Methods

The UK Biobank dataset was downloaded along with the variant call files for 200,632 subjects who had undergone exome sequencing and genotyping by the UK Biobank Exome Sequencing

Consortium using the GRCh38 assembly with coverage 20× at 95.6% of sites on average [12]. The UK Biobank participants are volunteers intended to be broadly representative of the UK population and are not selected on the basis of having any health condition. The UK Biobank had obtained ethics approval from the North West Multi-Centre Research Ethics Committee, which covers the UK (approval number: 11/NW/0382) and had obtained informed consent from all participants. The UK Biobank approved an application for use of the data (ID 51119) and ethics approval for the analyses was obtained from the UCL Research Ethics Committee (11527/001). All variants were annotated using the standard software packages VEP, PolyPhen, and SIFT [14–16]. To obtain population principal components reflecting ancestry, version 2.0 of *plink* (<https://www.cog-genomics.org/plink/2.0/>) was run with the options *-maf 0.1 -pca 20 approx* [17, 18].

The UK Biobank sample contains 503,317 subjects, of whom 94.6% are of white ethnicity. As we have discussed previously, it has become standard practice for investigators to simply discard data from participants with other ancestries and we regard this as regrettable [19]. We demonstrated that if population principal components are included as covariates, then it is possible to include all participants, regardless of ancestry, in the type of weighted burden analysis described here without inflation of the test statistic.

To define cases, a similar approach was used as was previously implemented for the investigation of hyperlipidaemia and T2D [13, 20, 21]. The hypertension phenotype was determined from 4 sources in the dataset: self-reported diagnosis recorded as hypertension or essential hypertension; reporting taking medication for high blood pressure; reporting taking any of a list of named medications commonly used to treat high blood pressure (<https://www.nhs.uk/conditions/high-blood-pressure-hypertension/>); having an ICD-10 diagnosis of essential hypertension, hypertensive heart disease, or hypertensive renal disease in hospital records or as a cause of death. Subjects in any of these categories were deemed to be cases with hypertension, while all other subjects were taken to be controls.

The same analytic methods as had been used previously were applied, with the description repeated here for the reader's convenience. The SCOREASSOC program was used to carry out a weighted burden analysis to test whether, in each gene, sequence variants that were rarer and/or predicted to have more severe functional effects occurred more commonly in cases than in controls. Attention was restricted to rare variants with minor allele frequency (MAF) ≤0.01 in both cases and controls. As previously described, variants were weighted by overall MAF, so that variants with MAF = 0.01 were given a weight of 1, while very rare variants with MAF close to 0 were given a weight of 10 [19]. Variants were also weighted according to their functional annotation using the GENEVARASSOC program, which was used to generate input files for weighted burden analysis by SCOREASSOC [22, 23]. The weights were informed from the analysis of the effects of different categories of variant in *LDLR* on hyperlipidaemia risk [13]. Variants predicted to cause complete loss of function (LOF) of the gene were assigned a weight of 100. Non-synonymous variants were assigned a weight of 5 but if PolyPhen annotated them as possibly or probably damaging, then 5 or 10 was added to this, and if SIFT annotated them as deleterious, then 20 was added. In order to allow exploration of the effects of different types of variants on disease risk, the variants were also grouped into broader categories to be used in

Table 1. Weight, which was assigned to each type of variant as annotated by VEP, PolyPhen, and SIFT software, and the broad categories, which were used for multivariate analyses of variant effects

VEP/SIFT/PolyPhen annotation	Weight	Category
Intergenic_variant	0	Unused
Feature_truncation	0	Intronic, etc.
Regulatory_region_variant	0	Intronic, etc.
Feature_elongation	0	Intronic, etc.
Regulatory_region_amplification	1	Intronic, etc.
Regulatory_region_ablation	1	Intronic, etc.
TF_binding_site_variant	1	Intronic, etc.
TFBS_amplification	1	Intronic, etc.
TFBS_ablation	1	Intronic, etc.
Downstream_gene_variant	0	Intronic, etc.
Upstream_gene_variant	0	Intronic, etc.
Non_coding_transcript_variant	0	Intronic, etc.
NMD_transcript_variant	0	Intronic, etc.
Intron_variant	0	Intronic, etc.
Non_coding_transcript_exon_variant	0	Intronic, etc.
3_prime_UTR_variant	1	3' UTR
5_prime_UTR_variant	1	5' UTR
Mature_miRNA_variant	5	Unused
Coding_sequence_variant	0	Unused
Synonymous_variant	0	Synonymous
Stop_retained_variant	5	Unused
Incomplete_terminal_codon_variant	5	Unused
Splice_region_variant	1	Splice region
Protein_altering_variant	5	Protein altering
Missense_variant	5	Protein altering
Inframe_deletion	10	InDel, etc.
Inframe_insertion	10	InDel, etc.
Transcript_amplification	10	InDel, etc.
Start_lost	10	Unused
Stop_lost	10	Unused
Frameshift_variant	100	Disruptive
Stop_gained	100	Disruptive
Splice_donor_variant	100	Splice site variant
Splice_acceptor_variant	100	Splice site variant
Transcript_ablation	100	Disruptive
SIFT deleterious	20	Deleterious
PolyPhen possibly damaging	5	Possibly damaging
PolyPhen probably damaging	10	Probably damaging

multivariate analyses as described below. The full set of weights and categories is displayed in Table 1. As described previously, the weight due to MAF and the weight due to functional annotation were multiplied together to provide an overall weight for each variant. Variants were excluded if there were >10% of genotypes missing in the controls or cases or if the heterozygote count was smaller than both homozygote counts in the controls or cases. If a subject was not genotyped for a variant, then they were assigned the subject-wise average score for that variant. For each subject, a gene-wise weighted burden score was derived as the sum of the variant-wise weights, each multiplied by the number of alleles of the variant that the given subject possessed. For variants on the X chromosome, hemizygous males were treated as homozygotes.

For each gene, logistic regression analysis was carried out with hypertension as the dependent variable, including the first 20 population principal components and sex as covariates and a likelihood ratio test was performed comparing the likelihoods of the

models with and without the gene-wise burden score. This is a test for association between the gene-wise burden score and caseness, and the statistical significance was summarized as a signed log p value (SLP), which is the $\log_{10}$ of the p value given a positive sign if the score is higher in cases and negative if it is higher in controls. No other clinical or demographic covariates were included in the analyses.

Gene set analyses were carried out as before using the 1454 “all GO gene sets, gene symbols” pathways as listed in the file *c5.all.v5.0.symbols.gmt* downloaded from the Molecular Signatures Database at <http://www.broadinstitute.org/gsea/msigdb/collections.jsp> [24]. For each set of genes, the natural logs of the gene-wise p values were summed according to Fisher’s method to produce a χ^2 statistic with degrees of freedom equal to twice the number of genes in the set. The p value associated with this χ^2 statistic was expressed as a minus $\log_{10}p$ as a test of association of the set with the hyperlipidaemia phenotype.

Fig. 1. QQ plot of SLPs obtained for weighted burden analysis of association with hypertension showing observed against expected SLP for each gene, omitting results for *DNMT3A*, which has SLP = 8.21. QQ, quantile-quantile; SLP, signed log p value.

For selected genes, additional analyses were carried out to clarify the contribution of different categories of variant. As described previously, logistic regression analyses were performed on the counts of the separate categories of variant as listed in Table 1, again with hypertension as the dependent variable and including principal components and sex as covariates, to estimate the effect size for each category [13]. The odds ratios (ORs) associated with each category were estimated along with their standard errors, and the Wald statistic was used to obtain a p value, except for categories in which variants occurred fewer than 50 times in which case Fisher's exact test was applied to the variant counts. The associated p value was converted to an SLP again with the sign being positive if the OR was >1 , indicating that variants in that category tended to increase risk. Data manipulation and statistical analyses were performed using GENEVARASSOC, SCOREASSOC, and R [25].

Results

There were 66,123 cases and 134,504 controls. There were 20,384 genes for which there were qualifying variants. Given that there were 20,384 informative genes, the critical threshold for the absolute value of the SLP to declare a result as formally statistically significant is $-\log_{10}(0.05/20,384) = 5.61$ and this was achieved by 2 genes, *DNMT3A* (SLP = 8.21) and *FES* (SLP = 6.10). The quan-

tile-quantile plot for the SLPs obtained for all genes except *DNMT3A* is shown in Figure 1. This shows that the test appears to be well-behaved and conforms well with the expected distribution. Omitting the genes with the 100 highest and 100 lowest SLPs, which might be capturing a real biological effect, the gradient for positive SLPs is 1.096 with intercept at 0.006 and the gradient for negative SLPs is 1.080 with intercept at 0.02, indicating only modest inflation of the test statistic in spite of the fact that participants of all ancestries are included.

Table 2 shows all the genes achieving SLP with absolute value >3 , equivalent to an uncorrected p value of 0.001. Given that 20,384 genes were tested, one would expect that by chance about 20 would reach this level of significance, whereas in fact there are 42. Thus, it is possible that some of these highly ranked genes do demonstrate a biological signal, which fails to reach statistical significance after correction for multiple testing. For *NPRI*, the analysis was repeated excluding data from the previously reported individually significant variants rs35479618, rs116245325, and rs61757359. This resulted in a reduction of the SLP from 5.14 to 4.38. For *DBH*, the analysis was repeated without rs3025380, and this resulted in a change in SLP from -3.40 to -2.11 . The full list of results for all genes is provided in online suppl. Table 1; see www.karger.com/doi/10.1159/000517419 for all online suppl. material.

In order to see if any additional genes were highlighted by analysing gene sets, gene set analysis was performed as described above after first dividing the gene-wise log p values by the average inflation factor of 1.09 before combining them using Fisher's method. Given that 1,454 sets were tested, a critical minus $\log_{10}p$ to achieve to declare results significant after correction for multiple testing would be $\log_{10}(1,454 \times 20) = 4.46$, and this was achieved by just one set, labelled CHROMATIN. This set contains 35 genes, including *DNMT3A*. The full results of the gene set analyses are listed in online suppl. Table 2. Four other sets achieved SLP >3 and these and for these all genes with an absolute value SLP >1.3 (equivalent to $p < 0.05$) are listed. All of these sets also contained *DNMT3A*, and none of the other nominally significant genes appeared to be obviously relevant to hypertension. These results are presented in online suppl. Table 3.

For the genes listed in Table 2 which appeared to be of interest, additional multivariate analyses were performed to elucidate the contribution to the overall result from different categories of variant. The results of this analysis for *DNMT3A* are shown in Table 3A. From this, it can be seen that the signal comes from disruptive and splice site

Table 2. Genes with absolute value of SLP exceeding 3 or more (equivalent to $p < 0.001$) for test of association of weighted burden score with hypertension

Symbol	SLP	Name
<i>DNMT3A</i>	8.21	DNA methyltransferase 3 alpha
<i>FES</i>	6.10	FES proto-oncogene, tyrosine kinase
<i>GUCY1A1</i>	5.54	Guanylate cyclase 1 soluble subunit alpha 1
<i>NPR1</i>	5.14	Natriuretic peptide receptor 1
<i>MAG</i>	4.52	Myelin-associated glycoprotein
<i>SMAD6</i>	4.10	SMAD family member 6
<i>GUCY1B1</i>	3.93	Guanylate cyclase 1 soluble subunit beta 1
<i>SKIV2L</i>	3.87	Ski2-like RNA helicase
<i>ASXL1</i>	3.75	ASXL transcriptional regulator 1
<i>MYO10</i>	3.75	Myosin X
<i>PTGER3</i>	3.52	Prostaglandin E receptor 3
<i>POLR3H</i>	3.47	RNA polymerase III subunit H
<i>IFT172</i>	3.39	Intraflagellar transport 172
<i>LITAF</i>	3.37	Lipopolysaccharide-induced TNF factor
<i>CSK</i>	3.37	C-terminal Src kinase
<i>SNX27</i>	3.34	Sorting nexin 27
<i>TEK</i>	3.31	EK receptor tyrosine kinase
<i>PHF21A</i>	3.23	PHD finger protein 21A
<i>LOC102723675</i>	3.21	Uncharacterized LOC102723675
<i>DXO</i>	3.14	Decapping exoribonuclease
<i>RHBDL3</i>	3.12	Rhomboid like 3
<i>ASB7</i>	3.12	Ankyrin repeat and SOCS box containing 7
<i>MPP2</i>	3.11	Membrane palmitoylated protein 2
<i>ZNF549</i>	3.10	Zinc finger protein 549
<i>MAP3K14-AS1</i>	3.08	MAP3K14 antisense RNA 1
<i>ZNF600</i>	-3.01	Zinc finger protein 600
<i>SIRT2</i>	-3.05	Sirtuin 2
<i>PARGP1</i>	-3.05	Poly(ADP-ribose) glycohydrolase pseudogene 1
<i>ZNF226</i>	-3.13	Zinc finger protein 226
<i>LOC105375153</i>	-3.18	Uncharacterized LOC105375153
<i>INPPL1</i>	-3.19	Inositol polyphosphate phosphatase like 1
<i>FNDC8</i>	-3.23	Fibronectin type III domain containing 8
<i>KRT82</i>	-3.26	Keratin 82
<i>GOLGA4</i>	-3.36	Golgin A4
<i>DBH</i>	-3.40	Dopamine beta-hydroxylase
<i>ELOF1</i>	-3.59	Elongation factor 1 homolog
<i>AGTR1</i>	-3.77	Angiotensin II receptor type 1
<i>SIPA1</i>	-3.81	Signal-induced proliferation-associated 1
<i>ZYX</i>	-3.83	Zyxin
<i>ZFAT</i>	-4.00	Zinc finger and AT-Hook domain containing
<i>OSTF1</i>	-4.05	Osteoclast-stimulating factor 1
<i>PREP</i>	-5.03	Prolyl endopeptidase

SLP, signed log p value.

variants, which are predicted to cause LOF and which are between them associated with an OR of about 1.9. However, variants annotated as probably damaging by PolyPhen are also common in cases and are associated with an OR of 1.5. Table 3B shows that the result for *FES* is driven by a small number of disruptive variants which are commoner in cases with OR of 2.8. It is striking that both *GUCY1A1* and *GUCY1B1* are ranked among the top 7 genes since they code for subunits of the same guanylate cyclase,

and their results are shown in full in Table 3C and D. This shows that while the effect for *GUCY1A1* is driven by LOF variants, these are very rare in *GUCY1B1*, and for this gene there seems to be an additional contribution from an excess of 5' UTR variants among cases. These occur at 44 different locations, and detailed inspection of the output file showed these all to be individually rare, such that there was no single variant which could be seen to making a significant contribution to the overall effect. For the

Table 3. Results from logistic regression analysis showing the effects on risk of hypertension of different categories of variant within *DNMT3A*, *FES*, *GUCY1A1*, and *GUCY1B1*

Category	Separate variants, <i>n</i>	Count in controls, <i>N</i>	Mean count in controls	Count in cases, <i>N</i>	Mean count in cases	OR (95% confidence interval)	SLP
(A) Results for <i>DNMT3A</i>							
Intronic, etc.	1,417	15,625	0.116168	7,859	0.118854	0.98 (0.95–1.00)	1.04
5' UTR	71	395	0.002937	187	0.002828	0.96 (0.80–1.15)	–0.20
Synonymous	157	2,427	0.018044	1,342	0.020296	1.04 (0.97–1.11)	0.55
Splice region	63	339	0.002520	206	0.003115	1.10 (0.92–1.32)	0.54
3' UTR	19	37	0.000275	10	0.000151	0.57 (0.28–1.16)	–0.92
Protein altering	227	1,310	0.009739	729	0.011025	1.08 (0.88–1.33)	0.36
InDel, etc.	7	13	0.000097	10	0.000151	1.62 (0.69–3.77)	0.56
Disruptive	53	76	0.000565	67	0.001013	1.86 (1.33–2.60)	3.62
Splice site variant	21	43	0.000320	39	0.000590	1.87 (1.20–2.92)	2.31
Deleterious	119	1,033	0.007680	577	0.008726	0.98 (0.75–1.28)	0.06
Possibly damaging	39	659	0.004899	339	0.005127	0.99 (0.78–1.26)	0.04
Probably damaging	68	185	0.001375	143	0.002163	1.50 (1.11–2.03)	2.18
(B) Results for <i>FES</i>							
Intronic, etc.	707	10,404	0.077351	5,607	0.084797	0.99 (0.97–1.02)	0.38
5' UTR	0	1	0.000007	1	0.000015	1.37 (0.08–23.60)	0.26
Synonymous	197	5,773	0.042921	3,557	0.053794	1.00 (0.96–1.05)	0.09
Splice region	63	387	0.002877	193	0.002919	0.94 (0.79–1.13)	0.29
3' UTR	25	98	0.000729	47	0.000711	0.92 (0.65–1.32)	–0.18
Protein altering	346	3,777	0.028081	1,950	0.029490	0.92 (0.86–0.99)	1.77
InDel, etc.	4	25	0.000186	15	0.000227	1.30 (0.67–2.51)	0.21
Disruptive	29	21	0.000156	29	0.000439	2.76 (1.55–4.92)	3.38
Splice site variant	7	6	0.000045	4	0.000060	1.38 (0.38–5.06)	0.13
Deleterious	186	721	0.005360	456	0.006896	1.20 (0.97–1.47)	1.08
Possibly damaging	66	785	0.005836	403	0.006095	1.10 (0.95–1.28)	0.72
Probably damaging	122	519	0.003859	338	0.005112	1.18 (0.93–1.50)	0.80
(C) Results for <i>GUCY1A1</i>							
Intronic, etc.	355	9,682	0.071983	5,306	0.080244	1.02 (0.98–1.06)	0.57
5' UTR	27	1,610	0.011970	864	0.013067	1.06 (0.97–1.15)	0.77
Synonymous	114	4,002	0.029754	2,438	0.036871	1.02 (0.96–1.08)	0.24
Splice region	18	35	0.000260	22	0.000333	1.25 (0.72–2.15)	0.37
3' UTR	16	64	0.000476	39	0.000590	1.30 (0.86–1.96)	0.70
Protein altering	281	2,538	0.018869	1,635	0.024727	1.00 (0.92–1.09)	0.00
InDel, etc.	2	2	0.000015	0	0.000000		0.00
Disruptive	29	35	0.000260	39	0.000590	2.41 (1.51–3.85)	3.76
Splice site variant	5	13	0.000097	12	0.000181	1.84 (0.82–4.11)	0.87
Deleterious	122	395	0.002937	242	0.003660	1.11 (0.81–1.52)	0.31
Possibly damaging	41	294	0.002186	158	0.002389	1.04 (0.83–1.31)	0.14
Probably damaging	92	285	0.002119	175	0.002647	1.12 (0.79–1.61)	0.29
(D) Results for <i>GUCY1B1</i>							
Intronic, etc.	599	5,482	0.040757	3,271	0.049468	1.01 (0.97–1.06)	0.22
5' UTR	44	163	0.001212	114	0.001724	1.37 (1.07–1.75)	1.94
Synonymous	96	3,833	0.028497	1,862	0.028160	0.99 (0.93–1.05)	–0.17
Splice region	38	107	0.000796	53	0.000802	1.00 (0.71–1.40)	0.01
3' UTR	20	119	0.000885	44	0.000665	0.77 (0.54–1.10)	–0.86
Protein altering	187	990	0.007360	545	0.008242	1.06 (0.94–1.20)	0.46
InDel, etc.	1	4	0.000030	0	0.000000		–0.51
Disruptive	12	10	0.000074	11	0.000166	2.06 (0.86–4.97)	1.18
Splice site variant	10	7	0.000052	12	0.000181	3.43 (1.32–8.94)	1.92
Deleterious	87	202	0.001502	128	0.001936	1.09 (0.75–1.56)	0.19
Possibly damaging	29	68	0.000506	44	0.000665	0.98 (0.65–1.48)	0.03
Probably damaging	59	109	0.000810	75	0.001134	1.21 (0.76–1.91)	0.38

ORs for each category are estimated including principal components and sex as covariates. The SLP is also obtained from this multivariate analysis except when there are fewer than 50 variants in a category, when Fisher's exact test is used instead. ORs, odds ratios; SLP, signed log *p* value.

Table 4. Summary results of variant category analysis for additional genes of interest

Gene/category	Separate variants, <i>n</i>	Count in controls, <i>N</i>	Mean count in controls	Count in cases, <i>N</i>	Mean count in cases	OR (95% confidence interval)	SLP
<i>NPR1</i>							
Disruptive	24	50	0.000372	44	0.000665	1.75 (1.15–2.65)	2.14
Splice site variant	7	10	0.000074	5	0.000076	1.07 (0.35–3.22)	0.00
Probably damaging	105	281	0.002089	195	0.002949	1.32 (1.05–1.66)	1.81
<i>SMAD6</i>							
Synonymous	193	3,378	0.025114	1,558	0.023562	0.94 (0.88–1.00)	–1.31
Disruptive	103	92	0.000684	68	0.001028	1.48 (1.07–2.03)	1.84
Splice site variant	7	4	0.000030	3	0.000045	1.65 (0.36–7.68)	0.16
Possibly damaging	74	208	0.001546	137	0.002072	1.29 (1.02–1.63)	1.55
<i>IFT172</i>							
Protein altering	711	7,499	0.055753	4,090	0.061854	1.07 (1.01–1.13)	1.82
Disruptive	58	94	0.000699	65	0.000983	1.39 (1.01–1.91)	1.40
Splice site variant	22	72	0.000535	34	0.000514	0.98 (0.64–1.49)	–0.04
<i>CSK</i>							
Splice region	38	969	0.007204	782	0.011826	1.20 (1.01–1.42)	1.47
Disruptive	3	2	0.000015	1	0.000015	0.93 (0.08–10.86)	0.00
Splice site variant	1	1	0.000007	1	0.000015	2.02 (0.12–34.90)	0.26
Deleterious	36	193	0.001435	182	0.002752	1.45 (1.07–1.97)	1.83
<i>DBH</i>							
Disruptive	27	71	0.000528	28	0.000423	0.83 (0.53–1.29)	–0.40
Splice site variant	8	544	0.004044	256	0.003872	0.96 (0.82–1.12)	–0.24
Deleterious	186	5,384	0.040029	2,343	0.035434	0.86 (0.76–0.98)	–1.65
<i>AGTR1</i>							
Disruptive	25	64	0.000476	13	0.000197	0.41 (0.22–0.75)	–2.46
Splice site variant	0	0	0.000000	0	0.000000		0.00
<i>ZYX</i>							
Synonymous	149	3,694	0.027464	2,162	0.032697	1.05 (1.00–1.11)	1.32
Disruptive	27	58	0.000431	9	0.000136	0.32 (0.16–0.67)	–2.76
Splice site variant	6	10	0.000074	4	0.000060	0.88 (0.27–2.89)	0.00
<i>PREP</i>							
Disruptive	18	20	0.000149	5	0.000076	0.50 (0.18–1.37)	–0.69
Splice site variant	3	3	0.000022	0	0.000000		–0.26

Results are shown for disruptive and splice site variants and for any other variant categories significant at $p < 0.05$. ORs, odds ratios; SLP, signed log p value.

other genes thought to be of interest, Table 4 provides a summary of the results for LOF variants along with any other variant category individually significant at $p < 0.05$. Full results for analyses of variant categories are presented in online suppl. Table 4.

Of the previously implicated genes listed in the Introduction, aside from *GUCY1A1*, only the following were significant at $p < 0.05$: *CYP11B1* (SLP = 1.63), *NR3C2* (SLP = 1.67), and *CACNA1H* (SLP = –2.05). Analyses of the variant categories were carried out for these genes, and a summary of the results is shown in Table 5, which

provides the results for LOF variants and for other categories that were significant at $p < 0.05$. These are mostly unremarkable, and it is difficult to draw firm conclusions although a few results are worth noting. For most genes, LOF variants are very rare so that one cannot gain a clear estimate of their effect. However, the results for *WINK1*, *WINK4*, *CLCN2*, and *ATP2B3* suggest that LOF variants in these genes do not have a very major effect on risk of hypertension. The most striking result is that for *CACNA1D*, the InDel category produces SLP = 8.10 with OR = 1.30. InDel variants occur at 9 locations in this gene,

Table 5. Summary results of variant category analysis for previously implicated genes

Gene/category	Separate variants, <i>n</i>	Count in controls, <i>N</i>	Mean count in controls	Count in cases, <i>N</i>	Mean count in cases	OR (95% confidence interval)	SLP
<i>CYP11B1</i>							
Disruptive	22	68	0.000506	38	0.000575	1.14 (0.76–1.72)	0.29
Splice site variant	6	21	0.000156	7	0.000106	0.71 (0.30–1.71)	−0.37
Probably damaging	93	531	0.003948	340	0.005142	1.27 (1.03–1.57)	1.60
<i>CYP11B2</i>							
Synonymous	150	3,649	0.027129	1,830	0.027676	0.94 (0.88–0.99)	−1.72
Protein altering	268	2,193	0.016304	1,251	0.018919	1.13 (1.03–1.25)	2.07
Disruptive	19	37	0.000275	23	0.000348	1.28 (0.75–2.18)	0.45
Splice site variant	7	46	0.000342	21	0.000318	0.93 (0.55–1.59)	−0.10
SIFT	129	957	0.007115	456	0.006896	0.74 (0.61–0.91)	−2.53
<i>WNK1</i>							
Disruptive	51	157	0.001167	84	0.001270	1.08 (0.82–1.42)	0.25
Splice site variant	8	31	0.000230	20	0.000302	1.16 (0.65–2.07)	0.22
<i>WNK4</i>							
Intronic, etc.	576	12,657	0.094101	7,753	0.117251	1.04 (1.00–1.07)	1.51
Disruptive	54	301	0.002238	165	0.002495	1.12 (0.92–1.36)	0.58
Splice site variant	18	24	0.000178	8	0.000121	0.67 (0.29–1.51)	−0.34
<i>KLHL3</i>							
Disruptive	5	6	0.000045	1	0.000015	0.33 (0.04–2.85)	−0.36
Splice site variant	4	5	0.000037	2	0.000030	0.95 (0.18–5.09)	0.00
<i>CUL3</i>							
5' UTR	41	108	0.000803	85	0.001285	1.54 (1.15–2.06)	2.51
InDel, etc.	3	0	0.000000	3	0.000045		1.45
Disruptive	4	4	0.000030	1	0.000015	0.53 (0.06–4.97)	0.00
<i>SCNN1B</i>							
Disruptive	16	31	0.000230	11	0.000166	0.73 (0.36–1.48)	−0.38
Splice site variant	8	10	0.000074	7	0.000106	1.64 (0.61–4.42)	0.35
<i>SCNN1G</i>							
InDel, etc.	3	68	0.000506	18	0.000272	0.54 (0.31–0.91)	−1.72
Disruptive	12	14	0.000104	8	0.000121	1.16 (0.48–2.83)	0.09
Splice site variant	4	17	0.000126	9	0.000136	1.06 (0.46–2.43)	0.08
<i>CYP17A1</i>							
Intronic, etc.	293	3,572	0.026557	2,081	0.031472	1.06 (1.00–1.11)	1.46
Disruptive	16	57	0.000424	28	0.000423	0.96 (0.60–1.52)	−0.07
Splice site variant	2	1	0.000007	2	0.000030	4.36 (0.37–50.99)	0.59
<i>HSD11B2</i>							
Synonymous	85	2,500	0.018587	1,581	0.023910	0.93 (0.86–1.00)	−1.37
Splice region	17	803	0.005970	377	0.005701	0.86 (0.76–0.98)	−1.67
Disruptive	8	16	0.000119	5	0.000076	0.66 (0.24–1.85)	−0.31
Splice site variant	4	7	0.000052	4	0.000060	1.08 (0.30–3.80)	0.12
Deleterious	91	226	0.001680	118	0.001785	0.69 (0.50–0.96)	−1.61
Probably damaging	46	68	0.000506	59	0.000892	2.50 (1.55–4.05)	3.86
<i>NR3C2</i>							
Disruptive	3	3	0.000022	0	0.000000		−0.26
Splice site variant	1	1	0.000007	0	0.000000		0.00
<i>KCNJ5</i>							
Disruptive	10	16	0.000119	12	0.000181	1.51 (0.70–3.27)	0.50
Splice site variant	1	1	0.000007	0	0.000000		0.00

Table 5 (continued)

Gene/category	Separate variants, <i>n</i>	Count in controls, <i>N</i>	Mean count in controls	Count in cases, <i>N</i>	Mean count in cases	OR (95% confidence interval)	SLP
<i>CACNA1H</i>							
Disruptive	56	84	0.000625	35	0.000529	0.86 (0.58–1.28)	−0.34
Splice site variant	21	27	0.000201	14	0.000212	1.10 (0.57–2.13)	0.06
<i>CACNA1D</i>							
InDel	9	1,360	0.010111	860	0.013006	1.30 (1.19–1.42)	8.10
Disruptive	12	8	0.000059	4	0.000060	1.11 (0.32–3.81)	0.00
Splice site variant	6	5	0.000037	4	0.000060	1.80 (0.47–6.91)	0.31
<i>CLCN2</i>							
Synonymous	180	1,167	0.008676	523	0.007910	0.89 (0.80–0.99)	−1.45
3' UTR	34	319	0.002372	125	0.001890	0.78 (0.63–0.96)	−1.78
Disruptive	41	87	0.000647	36	0.000544	0.85 (0.57–1.26)	−0.40
Splice site variant	14	14	0.000104	8	0.000121	1.05 (0.43–2.57)	0.09
<i>ATPIA1</i>							
Synonymous	201	5,833	0.043367	3,493	0.052826	0.95 (0.91–1.00)	−1.48
Disruptive	7	4	0.000030	4	0.000060	1.95 (0.47–8.10)	0.34
Splice site variant	1	1	0.000007	0	0.000000		0.00
<i>ATP2B3</i>							
Disruptive	9	226	0.001680	74	0.001119	0.91 (0.78–1.06)	−0.64
Splice site variant	2	2	0.000015	0	0.000000		0.00
<i>KDM1A</i>							
3' UTR	28	43	0.000320	34	0.000514	1.61 (1.01–2.56)	1.41
Disruptive	32	21	0.000156	20	0.000302	1.93 (1.05–3.58)	1.35
Splice site variant	7	8	0.000059	3	0.000045	0.79 (0.20–3.08)	0.00

Results are shown for disruptive and splice site variants and for any other variant categories significant at $p < 0.05$. ORs, odds ratios; SLP, signed log p value.

and inspection of the detailed results revealed that 8 of these are very rare, so the result reflects the effect of a single inframe deletion, 3:53808664-CCTT>C. This is rs72556363, which results in the loss of a phenylalanine residue, p.Phe1923del, and according to gnomAD, it has allele frequency of 0.0043 in non-Finnish Europeans and is extremely rare or absent from other populations. In the UK Biobank subjects, we observe a frequency of 0.0049 in controls and 0.0064 in cases. Full results for analyses of variant categories in these genes are presented in online suppl. Table 5.

Discussion

These analyses provide a broad overview of some impacts of very rare genetic variants in a large sample broadly representative of the population. It should be pointed out that the analytic approach used assumes that all vari-

ants in gene have the same direction of effect. Since any random variant is more likely to impair the function of a gene than to enhance it, weighted burden analysis is not expected to detect effects of gain of function variants because these are likely to be swamped by other variants. Likewise, the method is expected to be relatively insensitive at detecting variants that have a purely recessive effect. Of course, it is quite possible that the effect of complete LOF of one gene may be modified by functional variants affecting the other copy of the gene, but such complex heterozygous effects are difficult to detect. Additionally, a population sample differs from a specially recruited case-control sample in that it is not enriched for the phenotype in question and hence will have less power to detect either rare recessively acting variants or extremely rare variants with a dominant effect. Thus, we expect that there may well be important rare variant effects in addition to the ones that this study has highlighted.

Another contrast with specifically defined case-control studies is that the phenotype needs to be derived from measures that are provided, and, in order to improve power, it is desirable to use information that is available for a large number of participants. The phenotype used here attempts to reflect a clinical diagnosis of hypertension but of course will not do this as accurately as could be achieved in a specifically ascertained sample. Some participants would be on antihypertensive medication for indications other than hypertension, some would have undiagnosed hypertension, and for some, the diagnosis would simply be incorrect. Measured blood pressure itself was not used to inform the phenotype, in part because it might reflect blood pressure on medication and in part because single measurements may be less informative than indirectly relying on the clinical decisions which have fed into assigning a diagnosis or starting a prescription. It is certainly the case that the phenotype used here does not correspond exactly with what one would use for a systematically recruited sample in a case-control study focussing explicitly on hypertension. Any lack of accuracy in the phenotype is likely to reduce power rather than to produce false-positive results. The advantages of having a large sample to some extent balance the disadvantage of a less accurate phenotype.

When considering the contribution of risk within the population, we should note that the variants involved are rare. Although an MAF threshold of 0.01 was used, the majority of variants analysed are very much rarer than this, and for the variant categories with the most severe consequences, the cumulative frequency of variants in the category is also low. This means that few subjects will carry more than one variant with a severe consequence, and we can say that the mean count of variants of a particular category is a good approximation for the proportion of the subjects who carry a variant of that category.

The result for *DNMT3A* seems unlikely to be due to chance since it would remain significant at $p = 0.00016$ even after correction for the number of genes tested. The results show that LOF variants in this gene are associated with nearly doubling the OR for hypertension and are present in about 1 in 1,000 people, while a slightly larger number will have a variant annotated as probably damaging by PolyPhen, which moderately increases hypertension risk. *DNMT3A* is a DNA methyltransferase, and non-synonymous variants in it have been reported as causes of 2 different syndromes, Tatton-Brown-Rahman syndrome with overgrowth and intellectual disability and Heyn-Sproul-Jackson syndrome with microcephalic dwarfism, neither of which has hypertension as part of the phenotype [26, 27]. Missense variants in *DNMT3A* have also been reported in

autism spectrum disorder, and mice with *Dnmt3a* haploinsufficiency produced by heterozygous deletion of exon 19 have increased body weight and some behavioural alterations [28]. In a series of 210 patients with an overgrowth syndrome similar to Sotos syndrome but with no *NSD1* mutation, 4 had de novo non-synonymous mutations in *DNMT3A* and 2 had stop variants [29]. One of the stop variants was inherited from a normal mother in whom it was thought a somatic mutation had occurred and for the other the father's DNA was not available. Given the frequency of LOF variants in *DNMT3A* observed in our samples, it seems possible that the observation of 2 patients with stop variants was coincidental. Thus, it may be that while certain specific non-synonymous variants can cause severe phenotypes, generally reduced functioning of *DNMT3A* does not cause marked problems but is associated with increased risk of hypertension.

Although *DNMT3A* variants have not previously been reported to be associated with hypertension, as reviewed recently these findings are consistent with a body of results relating to the histone lysine demethylase LSD1, which together implicate LSD1 hypofunction in salt-sensitive hypertension [30]. In our data, disruptive variants in *KDM1A*, the gene for LSD1, are associated with an OR of 1.9, but they are too rare for firm conclusions to be drawn. *DNMT3A* methylates DNA conditional on the associated H3K4 residue being unmethylated and LSD1 accomplishes this demethylation, meaning that reduced function of LSD1 is expected to reduce *DNMT3A*-dependent DNA methylation [31, 32]. In mice, *Dnmt3a* deficiency, which can be produced by knockdown or as a consequence of foetal exposure to dexamethasone or low-protein maternal diet, has been shown to lead to reduced methylation of the gene for angiotensin receptor type 1a, *Agtr1a*, leading to increased *Agtr1a* expression and salt-induced hypertension [33]. Thus, a consistent picture emerges that reduced *DNMT3A* activity can increase risk for hypertension, whether this is due to genetic variants in *DNMT3A* itself, LSD1 hypofunction, or the uterine environment.

Although *FES* (SLP = 6.10) only just reaches criteria for exome-wide significance, confidence in this result is somewhat increased by the fact that a nearby SNP, rs2521501, shows robust evidence for association [34]. However, the mechanisms by which it might influence hypertension risk are unclear as it codes for a tyrosine kinase which is involved in various signalling pathways and which may have a role in haematopoiesis and regulating the innate immune response [35].

The results for *GUCY1A1* (SLP = 5.54) and *GUCY1B1* (SLP = 3.92) are more compelling, given that they code for

2 different subunits of the same protein, soluble guanylate cyclase, and given that recessively acting variants in *GU-CY1A1* have previously been reported in cases of moyamoya disease with hypertension [5]. Soluble guanylate cyclase is responsible for detecting NO signalling in order to produce vasodilation and other responses, and the central role of this pathway in the control of blood pressure is well established from animal studies, while guanylate cyclase stimulators have been developed as treatments for pulmonary hypertension [36]. The findings reported here are the first to directly demonstrate that impaired functioning of either of these genes represents a risk factor for systemic hypertension in the general population, with nearly 1 in 1,000 people carrying a LOF variant in one of them associated with an OR of approximately 2.

The results for *NPR1* (SLP = 5.14) represent a replication of the previously reported findings and confirm that although certain non-synonymous variants such as rs61757359 may be associated with reduced blood pressure, in general variants that impair the functioning of this gene increase the risk of hypertension [7, 8]. Around 1 in 500 people carries a variant annotated by PolyPhen as probably damaging, and overall, such variants are associated with a modest increase in risk with OR = 1.32 (1.05–1.66).

SMAD6 (SLP = 4.10) has a role in signalling pathways and has not previously been clearly implicated in hypertension risk although a recent report describes how exome sequencing of 37 children with renovascular hypertension revealed a frameshift variant classified as likely pathogenic variant in *SMAD6* in 1 patient [37]. *SMAD6* variants are known to predispose to cardiovascular malformations including bicuspid aortic valve-related aortopathy [38–40]. The results reported here suggest that LOF variants in this gene may have a moderate effect on increased risk of hypertension in the general population.

Recessively acting variants in *IFT172* (SLP = 3.39) can cause ciliopathies, and there is a report of a child with compound heterozygous variant who presented with growth retardation and subsequently developed retinopathy, metaphyseal dysplasia, and, at the age of 11 years, hypertension [41]. However, there does not seem to be other evidence to implicate *IFT172* in hypertension risk, so this result would require replication in other samples.

CSK (SLP = 3.37) is located in the 15q24 locus, which, as reviewed recently, is implicated by multiple GWAS for hypertension [42]. Following up the results of eQTL analyses, these authors demonstrated that mice with gene silencing or haploinsufficiency of *Csk* had increased blood pressure and showed that this effect could be moderated

by PP2, an inhibitor of Src. Although the results reported here are not formally significant after correction for multiple testing, the additional support provided by these GWAS findings and functional studies does suggest that variants in *CSK*, in particular those annotated as deleterious by SIFT, might be a risk factor for hypertension.

The results for *DBH* (SLP = –3.40) provide further support for the previously reported findings that variants in this gene are associated with blood pressure [7]. In particular, the results suggest that variants annotated as deleterious by SIFT are on average associated with a slightly reduced risk of developing hypertension. Although these variants are individually rare, about 1 person in 20 carries one of them.

AGTR1 (SLP = –3.77) codes for a receptor for angiotensin II, so it seems very plausible from a biological point of view that variants impairing its function might be protective against hypertension in spite of the fact that no association with common variants has been detected [43]. The results suggest that very rare gene disruptive variants can about halve the OR for hypertension.

ZYX (SLP = –3.83) is potentially of interest because it codes for zyxin, the protein responsible for sensing stretch in endothelial cells and vascular smooth muscle cells, as occurs in hypertension, and mediating their response to this by changing the expression of other genes [44]. The findings reported here suggest that impaired functioning of this gene may reduce risk of hypertension.

PREP (SLP = –5.03) narrowly fails to meet conventional criteria for exome-wide significance but is clearly of interest because its product, prolyl endopeptidase, also known as prolyl oligopeptidase or post-proline cleaving enzyme, has recently been shown to be responsible for converting circulating angiotensin II to angiotensin-(1–7) in the circulation and in the lungs, a process which is largely independent of ACE2, which carries out this conversion in the kidney [45]. The results we report suggest that rare functional variants in *PREP* are protective against hypertension, but it is not clear which categories of variant are responsible, and, although LOF variants are commoner in controls, they are too rare for conclusions to be drawn. In mice, loss of this gene results in reduced ability to metabolize ACE2 and hence produces a more prolonged systemic hypertensive response to exogenously administered ACE2 [45]. While it is not obvious why impaired functioning of this gene might be protective against hypertension, these findings do seem worthy of further exploration.

The finding that an inframe deletion in *CACNA1D*, rs72556363, is associated with increased hypertension is

consistent with reports that very rare germline and somatic variants in this gene can result in aldosterone-producing adenomas and primary aldosteronism [46]. Although the variant varies in frequency between populations, essentially being restricted to those with European ancestry, this result does not seem likely to be due to an artefact of population stratification, because the frequency in controls is similar to that reported in non-Finnish Europeans, whereas the frequency in cases is even higher. It seems to represent a modest risk factor for hypertension without producing severe hyperaldosteronism, which is found in about 1% of subjects with European ancestry.

Overall, these analyses provide an overview of some of the impacts rare coding variants may have on the risk of hypertension in the general population. The validity of some novel findings will become clearer when exome sequence data is released for the remaining 300,000 UK Biobank participants or if they can be tested in other samples or followed up in functional studies. All the variants implicated are very rare and in view of their effect sizes arguably do not make an important contribution to risk from a public health point of view. Nor are they probably helpful as individual measures of risk, partly because they can only be detected by sequencing. Although it may be reasonable to assume that LOF variants within a given gene will tend to have a similar effect on phenotype, the same cannot be said of non-synonymous variants, and even within a given category the effect of such variants is likely to vary considerably. Thus, for variants that are individually extremely rare, it is in general not possible to make a clear interpretation regarding their likely effect. The main value of these findings is probably in highlighting genes and biological pathways of relevance in order to ultimately inform improved therapeutic approaches.

Preprint Availability

A preprint describing this work has been posted at medRxiv with doi 10.1101/2021.02.10.21251503 [47].

Acknowledgements

This research has been conducted using the UK Biobank Resource. The author wishes to acknowledge the staff supporting the High Performance Computing Cluster, Computer Science Department, University College London. This work was carried out in part using resources provided by BBSRC equipment Grant BB/R01356X/1. The author wishes to thank the participants who volunteered for the UK Biobank project.

Statement of Ethics

The UK Biobank had obtained ethics approval from the North West Multi-Centre Research Ethics Committee, which covers the UK (approval number: 11/NW/0382) and had obtained informed consent from all participants. The UK Biobank approved an application for use of the data (ID 51119), and ethics approval for the analyses was obtained from the UCL Research Ethics Committee (11527/001).

Conflict of Interest Statement

The author declares he has no conflict of interest.

Author Contributions

D.C. carried out the analyses and prepared the manuscript.

Availability of Data and Materials

The raw data are available on application to the UK Biobank. Detailed results with variant counts cannot be made available because they might be used for subject identification. Scripts and relevant derived variables will be deposited in the UK Biobank. Software and scripts used to carry out the analyses are also available at <https://github.com/davenomiddlenamecurtis>.

References

- 1 Evangelou E, Warren HR, Mosen-Ansorena D, Mifsud B, Pazoki R, Gao H, et al. Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. *Nat Genet*. 2018 Oct 1;50(10):1412–25.
- 2 Wang Y, Wang JG. Genome-wide association studies of hypertension and several other cardiovascular diseases. *Pulse*. 2018;6(3–4):169–86.
- 3 Ehret GB, Caulfield MJ. Genes for blood pressure: an opportunity to understand hypertension. *Eur Heart J*. 2013;34:951–61.
- 4 Scholl UI, Stölting G, Schewe J, Thiel A, Tan H, Nelson-Williams C, et al. CLCN2 chloride channel mutations in familial hyperaldosteronism type II. *Nat Genet*. 2018 Mar 1;50(3):349–54.
- 5 Wallace S, Guo DC, Regalado E, Mellor-Crummey L, Bamshad M, Nickerson DA, et al. Disrupted nitric oxide signaling due to GUCY1A3 mutations increases risk for moyamoya disease, achalasia and hypertension. *Clin Genet*. 2016 Oct 1;90(4):351–60.
- 6 Lee J, Kim S-K, Kang HG, Ha I-S, Wang K-C, Lee JY, et al. High prevalence of systemic hypertension in pediatric patients with moyamoya disease years after surgical treatment. *J Neurosurg Pediatr*. 2020 Nov 8;25(2):131–7.

- 7 Liu C, Kraja AT, Smith JA, Brody JA, Franceschini N, Bis JC, et al. Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet*. 2016 Oct 1;48(10):1162–70.
- 8 Vandenwijngaert S, Ledsy CD, Lahrouchi N, Khan MAF, Wunderer F, Ames L, et al. Blood pressure-associated genetic variants in the natriuretic peptide receptor 1 gene modulate guanylate cyclase activity. *Circ Genom Precis Med*. 2019 Aug 1;12(8):e002472.
- 9 Pojoga LH, Williams JS, Yao TM, Kumar A, Raffetto JD, do Nascimento GR, et al. Histone demethylase LSD1 deficiency during high-salt diet is associated with enhanced vascular contraction, altered NO-cGMP relaxation pathway, and hypertension. *Am J Physiol Heart Circ Physiol*. 2011 Nov;301(5):H1862.
- 10 Williams JS, Chamarthi B, Goodarzi MO, Pojoga LH, Sun B, Garza AE, et al. Lysine-specific demethylase 1: an epigenetic regulator of salt-sensitive hypertension. *Am J Hypertens*. 2012 Jul;25(7):812–7.
- 11 Huang Y, Ting PY, Yao TM, Homma T, Brooks D, Katayama Rangel I, et al. Histone demethylase LSD1 deficiency and biological sex: impact on blood pressure and aldosterone production. *J Endocrinol*. 2019 Feb 1;240(2):111–22.
- 12 Szustakowski JD, Balasubramanian S, Sasson A, Khalid S, Bronson PG, Kvistad E, et al. Advancing human genetics research and drug discovery through exome sequencing of the UK Biobank. *medRxiv*. 2020 Jan 1.
- 13 Curtis D. Analysis of 200,000 exome-sequenced UK Biobank subjects illustrates the contribution of rare genetic variants to hyperlipidaemia [Internet]. *medRxiv*. 2021. Available from: <https://www.medrxiv.org/content/10.1101/2021.02.10.21251503v3>.
- 14 McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, et al. The Ensembl variant effect predictor. *Genome Biol*. 2016 Jun 6;17(1):122.
- 15 Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen-2. *Curr Protoc Hum Genet*. 2013 Jan;Chapter 7:Unit7.20.
- 16 Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc*. 2009 Jun 25;4(8):1073–81.
- 17 Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*. 2015 Dec 25;4(1):7.
- 18 Galinsky KJ, Bhatia G, Loh PR, Georgiev S, Mukherjee S, Patterson NJ, et al. Fast principal-component analysis reveals convergent evolution of ADH1B in Europe and East Asia. *Am J Hum Genet*. 2016 Mar 3;98(3):456–72.
- 19 Curtis D. Multiple linear regression allows weighted burden analysis of rare coding variants in an ethnically heterogeneous population. *Hum Hered*. 2021 Jan 7;1–10.
- 20 Curtis D. A weighted burden test using logistic regression for integrated analysis of sequence variants, copy number variants and polygenic risk score. *Eur J Hum Genet*. 2019 Jan 26;27(1):114–24.
- 21 Curtis D. Weighted burden analysis in 200,000 exome-sequenced subjects characterises rare variant effects on risk of type 2 diabetes. *medRxiv*. 2021 Jan 21.
- 22 Curtis D. A rapid method for combined analysis of common and rare variants at the level of a region, gene, or pathway. *Adv Appl Bioinform Chem*. 2012;5:1–9.
- 23 Curtis D. Pathway analysis of whole exome sequence data provides further support for the involvement of histone modification in the aetiology of schizophrenia. *Psychiatr Genet*. 2016;26:223–7.
- 24 Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545–50.
- 25 R Core Team. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2014.
- 26 Tatton-Brown K, Seal S, Ruark E, Harmer J, Ramsay E, Del Vecchio Duarte S, et al. Mutations in the DNA methyltransferase gene DNMT3A cause an overgrowth syndrome with intellectual disability. *Nat Genet*. 2014 Mar 9;46(4):385–8.
- 27 Heyn P, Logan CV, Fluteau A, Challis RC, Auchynnikava T, Martin CA, et al. Gain-of-function DNMT3A mutations cause microcephalic dwarfism and hypermethylation of Polycomb-regulated regions. *Nat Genet*. 2019 Jan 1;51(1):96–105.
- 28 Christian DL, Wu DY, Martin JR, Moore JR, Liu YR, Clemens AW, et al. DNMT3A haploinsufficiency results in behavioral deficits and global epigenomic dysregulation shared across neurodevelopmental disorders. *Cell Rep*. 2020 Nov 24;33(8):108416.
- 29 Tlemsani C, Luscan A, Leulliot N, Bieth E, Afeenjar A, Baujat G, et al. SETD2 and DNMT3A screen in the sotos-like syndrome French cohort. *J Med Genet*. 2016 Nov 1;53(11):743–51.
- 30 Hirohama D, Fujita T. Evaluation of the pathophysiological mechanisms of salt-sensitive hypertension. *Hypertens Res*. 2019;42:1848–57.
- 31 Shi Y, Lan F, Matson C, Mulligan P, Whetstone JR, Cole PA, et al. Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell*. 2004 Dec 29;119(7):941–53.
- 32 Hashimoto H, Vertino PM, Cheng X. Molecular coupling of DNA methylation and histone methylation. *Epigenomics*. 2010;2:657–69.
- 33 Kawakami-Mori F, Nishimoto M, Reheman L, Kawarazaki W, Ayuzawa N, Ueda K, et al. Aberrant DNA methylation of hypothalamic angiotensin receptor in prenatal programmed hypertension. *JCI Insight*. 2018 Nov 2;3(21):e95625.
- 34 Ehret GB, Ehret GB, Munroe PB, Rice KM, Bochud M, Johnson AD, et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature*. 2011 Oct 6;478(7367):103–9.
- 35 Zirngibl RA, Senis Y, Greer PA. Enhanced endotoxin sensitivity in Fps/Fes-null mice with minimal defects in hematopoietic homeostasis. *Mol Cell Biol*. 2002 Apr 15;22(8):2472–86.
- 36 Buys E, Sips P. New insights into the role of soluble guanylate cyclase in blood pressure regulation. *Curr Opin Nephrol Hypertens*. 2014;23:135–42.
- 37 Viering DHHM, Chan MMY, Hoogenboom L, Iancu D, de Baaij JHF, Tullus K, et al. Genetics of renovascular hypertension in children. *J Hypertens*. 2020 Oct 1;38(10):1964–70.
- 38 Tan HL, Glen E, Töpf A, Hall D, O'Sullivan JJ, Sneddon L, et al. Nonsynonymous variants in the SMAD6 gene predispose to congenital cardiovascular malformation. *Hum Mutat*. 2012 Apr;33(4):720–7.
- 39 Luyckx I, MacCarrick G, Kempers M, Meester J, Geryl C, Rombouts O, et al. Confirmation of the role of pathogenic SMAD6 variants in bicuspid aortic valve-related aortopathy. *Eur J Hum Genet*. 2019 Jul 1;27(7):1044–53.
- 40 Gillis E, Kumar AA, Luyckx I, Preuss C, Canaerts E, van de Beek G, et al. Candidate gene resequencing in a large bicuspid aortic valve-associated thoracic aortic aneurysm cohort: SMAD6 as an important contributor. *Front Physiol*. 2017 Jun 13;8(Jun):400.
- 41 Lucas-Herald AK, Kinning E, Iida A, Wang Z, Miyake N, Ikegawa S, et al. A case of functional growth hormone deficiency and early growth retardation in a child with IFT172 mutations. *J Clin Endocrinol Metab*. 2015 Apr 1;100(4):1221–4.
- 42 Lee HJ, Kang JO, Kim SM, Ji SM, Park SY, Kim ME, et al. Gene silencing and haploinsufficiency of Csk increase blood pressure. *PLoS One*. 2016 Jan 11;11(1):e0146841.
- 43 Ji LD, Li JY, Yao BB, Cai XB, Shen QJ, Xu J. Are genetic polymorphisms in the renin-angiotensin-aldosterone system associated with essential hypertension? Evidence from genome-wide association studies. *J Hum Hypertens*. 2017;31:695–8.
- 44 Ghosh S, Kollar B, Nahar T, Suresh Babu S, Wojtowicz A, Sticht C, et al. Loss of the mechanotransducer zyxin promotes a synthetic phenotype of vascular smooth muscle cells. *J Am Heart Assoc*. 2015 Jun 12;4(6):e001712.
- 45 Serfozo P, Wysocki J, Gulua G, Schulze A, Ye M, Liu P, et al. Ang II (angiotensin II) conversion to angiotensin-(1-7) in the circulation is POP (prolyloligopeptidase)-dependent and ACE2 (angiotensin-converting enzyme 2)-independent. *Hypertens*. 2020 Jan 1;75(1):173–82.
- 46 Scholl UI, Goh G, Stölting G, De Oliveira RC, Choi M, Overton JD, et al. Somatic and germline CACNA1D calcium channel mutations in aldosterone-producing adenomas and primary aldosteronism. *Nat Genet*. 2013 Sep;45(9):1050–4.
- 47 Curtis D. Analysis of 200,000 exome-sequenced UK Biobank subjects implicates genes involved in increased and decreased risk of hypertension. *medRxiv*. 2021 Feb 16.