Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Materials/Methods: An anonymous, voluntary survey directed towards attending radiation oncologists was distributed via social media platforms (Twitter, LinkedIn, SDN, Facebook). Questions assessed practice characteristics, radiopharmaceutical prescribing patterns (Iodine-131, Radium-223, Yttrium-90, Samarium-153, Strontium-89, Phosphorus-32, “other”), prescribing interest level, and perceived barriers to implementation. Non-parametric χ^2 test was used for correlation statistics.

Results: Of the 120 respondents, 112 practiced in the U.S and were included for this analysis. Years of practice was broadly distributed with 30% (n = 34) of attendings in practice > 10 years, 21% (n = 23) 6-10 years, 25% (n = 28) 3-5 years, and 24% (n = 27) 0-2 years. Practice type included academic-main site amongst 39% (n = 44), academic-satellite in 18% (n = 20), and private practice/other in 43% (n = 48) of attendings. Forty-four percent (n = 49) of attendings served populations of > 500K, 22% (n = 25) 200-500K, 25% (n = 28) 100-200K, and 9% (n = 10) < 100K. Forty-eight percent (n = 52) reported prescribing (physically and/or billing) at least one radiopharmaceutical/unsealed source. The most to least commonly prescribed was Ra-223 (39%; mean 2.5 cases/mos.), I-131 (17%; mean 3.1 cases/mos.), Y-90 (14%; mean 3.9 cases/mos.), “other” (8%), Sm-153 (7%; mean 1.3 cases/mos.), Sr-89 and P-32 (3% each; mean 2.5 and 0.6 cases/mos., respectively). Of those who answered “other”, Lutetium-177 dotatate was most commonly prescribed (8%). No significant ($P > 0.05$) association was noted between practice type, practice location, years of practice, or practice volume with utilization of any radiopharmaceuticals/unsealed sources. The majority (55%, n = 61) of attending radiation oncologists responded they would like to actively prescribe more radiopharmaceuticals/unsealed sources, while 30% (n = 31) were indifferent and 18% (n = 20) answered “no.” Perceived barriers to implementation were varied but broadly categorized into treatment infrastructure (43%, n = 48), interspecialty relations (42%, n = 47), lack of training (21%, n = 24), and financial considerations (17%, n = 19).

Conclusion: Almost half of surveyed U.S. radiation oncologists reported prescribing at least one radiopharmaceutical/unsealed source, of which Ra-223 was the most common. The majority expressed interest in prescribing additional radiopharmaceuticals/unsealed sources. Wide-ranging barriers to implementation exist, most commonly interspecialty relations, treatment infrastructure, lack of training, and financial considerations.

Author Disclosure: I.H. Chowdhury: None. U. Shukla: None. J.S. Witt: None. M. McFarlane: None. T.J. Royce: None. C.J. Miller: None. K.E. Huber: None. M.S. Katz: Partner; Radiation Oncology Associates, PA. Stock; Dr. Reddy's Laboratories, U.S. Physical Therapy, Mazor Robotics, Healthcare Services Group, Kiniksa Pharmaceuticals, CVS Health. Oversees blog posts for both #radonc journal club on Twitter and multi-author blog posts about radiation medicine. No advertising money accepted for the blog, nonprofit. M. Chowdhury: Stock; Varian Health Systems, Teladoc Health.

2732

Implementation of a Nurse Practitioner Led CNS Follow-Up and Survivorship Clinic in a NCI-Designated Cancer Center

P. Estep,¹ B.W. Fischer-Valuck,² C. Deibert,³ S. Parker,⁴ and S.T. Kahn¹; ¹Department of Radiation Oncology, Winship Cancer Institute of Emory University, Atlanta, GA, ²Winship Cancer Institute, Department of Radiation Oncology, Emory University, Atlanta, GA, ³Emory University, Atlanta, GA, ⁴Emory St. Joseph's Hospital, Atlanta, GA, United States

Purpose/Objective(s): The role of the advanced practice provider (APP) in radiation oncology (RO) is not well defined but it has been shown that the APP can provide safe and efficient patient care. Most APPs are certified for primary care without specific concentration in oncology or RO during formal training. There are very limited fellowships available to train APPs before providing care in a specialty setting such as RO and even fewer resources for disease-site specific training. The purpose of this work is to describe our workflow and integration of a Nurse Practitioner (NP) into the RO clinic with a specific emphasis on independent management of a central nervous system (CNS) follow-up and survivorship clinic.

Materials/Methods: MDs and NPs provide services for our NCI Designated Comprehensive Cancer Center which includes 10 hospitals/clinics with services/technology including external beam radiation therapy, brachytherapy, Gamma Knife radiosurgery (GK) and proton beam therapy. The hospital which is designated for GK includes 7 radiation oncologists (3 subspecializing in CNS diseases), 2 neurosurgeons who perform GK, and 2 NPs. Within this hospital, a NP assumed leadership of our CNS follow up clinic. We herein reviewed the clinical workflow for implementing this service and evaluated metrics associated with this clinic.

Results: In 2019, a NP assumed leadership for our CNS follow up clinic. Upon development of this workflow, our CNS radiation oncologists and neurosurgeons developed standardized imaging and follow up schedules for the various CNS diseases treated. The MDs provided education to the NP on management of CNS radiation side effects and imaging review. For 3-6 months prior to implementation of our clinic, the NP saw follow ups in conjunction with the radiation oncologist and neurosurgeon. This provided confidence to the NP and MDs to ensure the images were reviewed properly and the patients were satisfied with their quality of care. Given the complexity of imaging review, the NP introduced a weekly CNS imaging review meeting during which the imaging of all follow ups are reviewed. The NP also became the organizer of a bimonthly CNS tumor conference. In 2020 the NP provided independent management of the clinic and saw 360 CNS specific follow ups (out of 1,040 of their total follow ups) accounting for 35% of their practice. This correlated with an approximate increase of 10% in the number of consultations RO providers saw during this time period. In 2020, the NP's overall patient satisfaction score was 94.9% compared to an average of 90% for all RO providers.

Conclusion: A CNS follow up clinic led by a NP was successfully implemented at our NCI Designated Cancer Center. This service was associated with excellent patient care and satisfaction and provided additional time and effort for our RO providers and neurosurgeons to see new consults and perform procedures. With specialty training from the radiation oncologist and neurosurgeon, an APP can be a critical asset to the RO team specifically in the CNS setting.

Author Disclosure: P. Estep: None. B.W. Fischer-Valuck: Research Grant; Varian. C. Deibert: None. S. Parker: None. S.T. Kahn: None.

2733

Spread of Misinformation? Quality of COVID-19 Resources for Cancer Patients

R. Jin,¹ and P.A. Ingledew²; ¹MD Undergraduate Program, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada, ²Radiation Oncology, BC Cancer Agency; Department of Surgery, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada

Purpose/Objective(s): Cancer patients are increasingly using the Internet to educate themselves about COVID-19. Recent studies have shown that cancer patients are at risk of more serious outcomes of COVID-19 compared to the general population. Some cancer treatments such as chemotherapy can impact the immune system, which may make COVID-19 infection more dangerous. This study looks to systematically examine the quality of web resources available for cancer patients about COVID-19.

Materials/Methods: The term “COVID-19 Risk and Cancer” was searched in Google and metasearch engines Yippy and Dogpile. URLs were recorded from each search and inclusion and exclusion criteria were applied. The results from the 3 lists were combined to come up with a final list based on overall average rank order. This list was analyzed using a previously validated structured rating tool with respect to accountability, currency, interactivity, readability, and content coverage and accuracy.

Results: 398 websites were identified prior (this includes overlap between the three search sites used), 37 websites were included for analysis. Out of 37 websites, only 43% disclosed authorship and 24% cited sources. Most websites (76%) revealed date of creation, and 32% were updated less than

3 months before the date of search. 68% of websites enabled questions to be sent to the author or webmaster regarding COVID-19 risk queries. 54% of websites had high school readability (8.0-12.0), 43% were at university level or above, and only one website demonstrated the recommended reading level for general public (below 8.0). Topics most commonly discussed were special consideration for cancer patients in COVID-19 (84%), COVID-19 risk factors (73%), and infection prevention (62%), while topics least covered were COVID-19 incidence/prevalence (5%), prognosis (8%), and treatment (16%).

Conclusion: There is some COVID-19 in cancer risk information available online, but quality is variable. The total number of sites with relevant information related to COVID-19 and cancer was relatively low and many sites lacked markers for accountability. Some information may not be up to date and content may be difficult to comprehend. Healthcare professionals may direct cancer patients to the most reliable online resources about COVID-19 and cancer shown in this study. In addition, this may be helpful to consider when designing comprehensive web resources regarding COVID-19.

Author Disclosure: R. Jin: None. P. Ingledew: None.

2734

COVID-19 Testing Trends: Pre-Radiation and Throughout Cancer Care

I. Messing,¹ Y.J. Rao,² D. Scully,² M. Ojong-Ntui,² S. Goyal,² and M.P. Huynh-Le²; ¹The George Washington University School of Medicine and Health Sciences, Washington, DC, ²The George Washington University, Department of Radiology, Division of Radiation Oncology, Washington, DC

Purpose/Objective(s): Timely care is critical for individuals diagnosed with cancer—despite the ongoing COVID-19 pandemic—as delays adversely affect cancer outcomes. Patients presenting for radiotherapy (RT) at a single institution who underwent COVID-19 screening procedures prior to RT were analyzed regarding treatment delays and disparities.

Materials/Methods: Our institution is an urban multidisciplinary cancer center. In April 2020, our Radiation Oncology department implemented universal COVID-19 screening protocols prior to RT initiation (with possible patient opt-out) to ensure patient/staff safety. Patients did not start RT until testing negative. We collected patient demographics and COVID-19 testing information on patients planned for RT from April to October 2020. We studied trends of other lifetime COVID-19 testing that these patients received to evaluate for overall delays. Summary statistics were analyzed within a data collection web application.

Results: 177 consecutive patients with cancer were scheduled to begin RT. 15 (8.5%) declined pre-RT COVID-19 testing and proceeded to RT. Of patients who consented to pre-RT testing, 99 (61.1%) were female; mean age was 60.6 (range 24-94). Most patients were Black (n = 98, 60.5%). 61 (37.7%) and 41 (25.3%) were insured through Medicare and Medicaid, respectively. Common primary cancer sites were breast (n = 52, 32.1%), prostate (n = 27, 16.7%), and lung (n = 23, 14.2%). 58 (35.8%) patients presented with metastatic disease; the most common metastatic sites were bone (n = 26, 44.8%) and brain (n = 12, 20.7%). Most patients received external RT (n = 152, 93.8%), primarily adjuvant (n = 66, 40.7%) and palliative (n = 59, 36.4%) RT. One (0.6%) asymptomatic patient tested "presumptive positive" for COVID-19 pre-RT. The patient was COVID-19 negative on immediate repeat testing and proceeded to RT. Overall, the 162 patients who consented to pre-RT testing received 549 total lifetime COVID-19 tests. Most of these patients received palliative RT (n = 219, 39.9%), followed by adjuvant (n = 199, 36.2%) and definitive RT (n = 101, 18.4%). Mean number of total COVID-19 tests per patient was 3.4 (range 1-14). Only 32 patients (19.8%) were administered a solitary pre-RT COVID-19 screening test. 24 (14.8%) patients had at least one positive COVID-19 test over the study period. In total, 32 (5.8%) positive COVID-19 test results were documented; 11 patients (6.8%) experienced a mean RT delay of 4.0 (range 3-8) weeks until testing negative. Other care delays

included surgical/diagnostic procedure delays (n = 4), delayed presentation due to COVID-19 fears (n = 2), and travel restrictions (n = 1).

Conclusion: The majority of patients with cancer planned for RT tested negative for COVID-19 and proceeded to RT without delay. However, delays prior to radiation oncology consultation due to diagnostic workup, imaging, or biopsies, as well as testing positive for COVID-19, could intensify underlying disparities affecting our urban patient population.

Author Disclosure: I. Messing: None. Y.J. Rao: None. D. Scully: None. M. Ojong-Ntui: None. S. Goyal: Employee; Center for Rheumatic Disease. M. Huynh-Le: None.

2735

Enhancing Follow Up Capacity and Quality at a High-Volume Academic Satellite Practice Through Integration of an Advanced Practice Provider

E.R. Miller,¹ A.W. Rajkumar,² E.T. Shinohara,¹ H.J. Jackson,¹ and E.C. Osmundson¹; ¹Vanderbilt University Medical Center, Nashville, TN, ²Department of Radiation Oncology, Vanderbilt University Medical Center, Nashville, TN

Purpose/Objective(s): Routine follow-up appointments have been limited at one of our academic facility's satellite clinics due to physician staffing limitations and high consultation volumes. We hypothesized adding an advanced practice provider (APP) on a limited schedule would permit meaningful follow up visits and improve care, while maintaining or increasing patient satisfaction.

Materials/Methods: Beginning in June 2020, an APP was embedded in our academic center satellite clinic one half-day per week to provide routine follow up care after radiation therapy (RT). Patients were provided an anonymized, voluntary satisfaction survey after each initial APP visit from June 2020-Jan 2021. Demographic information and unmet care needs at time of follow up visit were recorded. We compared characteristics of patients seen in the same clinic from June 2019-Jan 2020 and of those seen from June 2020-Jan 2021, prior to APP integration. All patients who completed definitive RT from March 2019-Oct 2019 and from March 2020-Oct 2020 were included for follow up rate comparison.

Results: In one half-day of clinic per week, an APP staffed 27.2% (44/118) of all follow up visits at the satellite clinic from June 2020-January 2021, compared to 5.2% of all follow ups seen by an APP from June 2019-Jan 2020. Over comparable timeframes, overall follow up > 90 days from definitive RT completion increased from 33.1% to 46.6% after APP addition. The greatest gains were in patients with cancers of the breast (from 28.6% to 45.7%), prostate (from 37.0% to 47.8%), head and neck (from 68.8% to 88.9%), and lung (from 15.4% to 28.6%). 20.5% of visits with the APP versus 0% with a physician incorporated formalized survivorship care. The APP saw a higher proportion of patients with female gender (64.1 vs 50.5%), age < 65 (59.0% vs 46.4%), and higher estimated median income (EMI) (83.4% vs 76.3%, EMI > \$50,000). Importantly, the APP identified and addressed an average of 2.3 unmet care needs per patient at time of follow up, which notably resulted in a new diagnosis of early-stage lung cancer in one patient. Patient satisfaction with APP-based follow up was > 95%, with 95% of patients surveyed indicating they were likely to continue follow up in the clinic.

Conclusion: Utilizing an APP at a high-volume radiation oncology satellite clinic improved follow up care by addressing unmet care needs and incorporating survivorship care. Follow up with the APP resulted in high patient satisfaction. These findings emphasize the unique value of radiation oncology-specific follow up, even among those receiving follow up with other oncology sub-specialists, and have important implications for increasing quality of care in high-volume satellite clinics. Based on this interim analysis, APP staffing will be increased to one full day per week to further increase follow up capacity. Data collection is ongoing with final analysis to be completed by July 2021.

Author Disclosure: E.R. Miller: None. A.W. Rajkumar: None. E.T. Shinohara: None. H.J. Jackson: None. E.C. Osmundson: None.