Published in final edited form as:

Atherosclerosis. 2019 November ; 290: 103–110. doi:10.1016/j.atherosclerosis.2019.09.021.

Genital *Chlamydia* infection in hyperlipidemic mouse models exacerbates atherosclerosis

Uma M. Nagarajan^{1,2}, James D. Sikes², Ramona L. Burris³, Rajneesh Jha², Branimir Popovic⁴, Pamela Fraungruber⁵, Leah Hennings⁶, Catherine L. Haggerty⁷, Shanmugam Nagarajan^{3,4,5,@}

¹Department of Pediatrics and Department of Microbiology and Immunology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

²Department of Pediatrics, University of Arkansas for Medical Sciences, Arkansas Children's Hospital Research Institute, Little Rock, AR USA

³Department of Microbiology and Immunology, University of Arkansas for Medical Sciences, and Arkansas Children's Hospital Research Institute, Little Rock, AR USA

⁴Department of Pathology, University of Pittsburgh, Pittsburgh, PA USA

⁵Department of Pathology, University of North Carolina at Chapel Hill, Chapel Hill, NC USA

⁶Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, AR USA

⁷Department of Epidemiology, University of Pittsburgh Graduate School of Public Health, Pittsburgh, PA, USA

Abstract

Background and aims.—Atherosclerosis is a chronic inflammatory disease, and recent studies have shown that infection at remote sites can contribute to the progression of atherosclerosis in hyperlipidemic mouse models. In this report, we tested the hypothesis that genital *Chlamydia* infection could accelerate the onset and progression of atherosclerosis.

Methods.—Apolipoprotein E (*ApoE*^{−/−}) and LDL receptor knockout (*Ldlr*^{−/−}) mice on a high-fat diet were infected intra-vaginally with *Chlamydia muridarum*. Atherosclerotic lesions on the aortic sinuses and in the descending aorta were assessed at 8-weeks post-infection. Systemic, macrophage, and vascular site inflammatory responses were assessed and quantified.

@ **Corresponding author:** Shanmugam Nagarajan, Department of Pathology and Laboratory Medicine, UNC at Chapel Hill, 160, N. Medical Drive, Chapel Hill, NC 27599, shannag@med.unc.edu, Tel: (919) 843 5417.

Author contributions:UMN and SN conceived and designed the experiments. JDS, RLB, RJ, BP, PF, UMN, and SN performed experiments. JDS, RLB, BP, UMN, and SN analyzed the data. LH performed pathology score of the samples, UMN, CLH, and SN wrote the manuscript with contribution from other authors.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest: The authors declared they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

Results.—Compared to the uninfected groups, infected *Apoe*^{-/-} and *Ldlr*^{-/-} mice developed significantly more atherosclerotic lesions in the aortic sinus and in the descending aorta. Increased lesions were associated with higher circulating levels of serum amyloid A-1, IL-1 β , TNF- α , and increased VCAM-1 expression in the aortic sinus, suggesting an association with inflammatory responses observed during *C. muridarum* infection. Genital infection courses were similar in *Apoe*^{-/-}, *Ldlr*^{-/-}, and wild type mice. Further, *Apoe*^{-/-} mice developed severe uterine pathology with increased dilatations. Apoe-deficiency also augmented cytokine/chemokine response in *C. muridarum* infected macrophages, suggesting that the difference in macrophage response could have contributed to the genital pathology in *Apoe*^{-/-} mice.

Conclusions.—Overall, these studies demonstrate that genital *Chlamydia* infection exacerbates atherosclerotic lesions in hyperlipidemic mouse and suggest a novel role for Apoe in full recovery of uterine anatomy after chlamydial infection.

Graphical Abstract

Keywords

atherosclerosis; *Chlamydia*; inflammation; uterine pathology

Introduction

Atherosclerosis is a chronic inflammatory disease initiated by endothelial dysfunction, macrophage activation, and lipid accumulation in the vasculature¹. Several studies provide evidence that infectious agents may accelerate atherosclerotic processes. For example, *Porphyromonas gingivalis*, *Helicobacter pylori*, HIV, and *Chlamydia pneumoniae* infection have been linked with atherosclerosis in humans²⁻⁴. Respiratory *C. pneumoniae* infection and *Chlamydia* heat shock protein have been associated with atherosclerosis in multiple

human serologic studies, providing some of the first evidence of an association between infection and atherosclerosis^{5–7}. *C. pneumoniae* has been detected in human atherosclerotic arteries in multiple studies^{5–8}. Mouse models of *Chlamydia* lung infection support the clinical observations from *C. pneumoniae* association with atherosclerosis. *C. pneumoniae* respiratory infection accelerates hyperlipidemia-induced atherosclerotic lesion development in C57BL/6J (wild type, WT)^{9, 10}, apolipoprotein-deficient (*Apoe*^{−/−})¹¹, and low-density lipoprotein receptor-deficient (*Ldlr*^{−/−}) mice^{12, 13}.

Genital *Chlamydia trachomatis* infection is the most prevalent sexually transmitted bacterial infection and is a common cause of pelvic inflammatory disease (PID)¹⁴. *Chlamydia* infection followed by inflammation of the uterine lining (termed endometritis) and/or fallopian tubes (termed salpingitis), can lead to infertility in some women¹⁵. A link between genital chlamydial infection and atherosclerosis merits investigation because of the high prevalence of *Chlamydia* infection in youth, high prevalence of cardiovascular disease in adulthood, and the socioeconomic association of both infection and cardiovascular disease^{16, 17}. However, there have been no studies investigating whether or not genital *Chlamydia trachomatis* infection primes the host for accelerated atherosclerosis. We hypothesize that the inflammatory response to genital *C. trachomatis* infection could enhance the progression of atherosclerosis. This hypothesis was tested using the mouse model for genital *Chlamydia* infection, which has served as a powerful tool in the identification of key innate and adaptive inflammatory pathways^{18, 19}. Using two independent hyperlipidemic mouse models *Apoe*^{−/−} and *Ldlr*^{−/−} mice, we show that *C. muridarum* genital infection exacerbates atherosclerotic lesions. Interestingly, *Apoe*^{−/−} mice uteri also developed severe dilatation post infection, suggesting a *Apoe*-deficiency promotes uterine pathology after chlamydial infection.

Materials and methods

Chlamydial stocks, mice, and diet:

Chlamydia muridarum, “Nigg” strain (UAMS, Little Rock, AR) was propagated as described in an earlier study¹⁸. Wild type mice (WT, C57BL/6J 000664), mice deficient in *Apoe* (*Apoe*^{−/−}, 002052) or *Ldlr* (*Ldlr*^{−/−}, 002207) in C57BL/6J background were purchased from the Jackson Laboratory. Institutional Animal Care and Use Committees at the University of Arkansas for Medical Sciences, University of Pittsburgh, and UNC at Chapel Hill approved all animal experiments. *Apoe*^{−/−} mice were fed a high-fat diet (TD88137, 42% fat and 0.5% cholesterol), and *Ldlr*^{−/−} mice were fed a high-fat (15.8% fat)/high-cholesterol (1.25%) diet (TD94059)^{20, 21}.

Genital infections of mice:

Seven days prior to infection, all mice (8-wks) received 2.5 mg of Depo-Provera subcutaneously in 100 µl of PBS. A week later, one set of mice were anesthetized with nembutal (240–250 µl of 5 mg/ml stock) and infected by administering 3×10^5 infectious forming units (IFU) of *C. muridarum* in 10 ml SPG buffer (250 mM sucrose, 10 mM sodium phosphate, 5 mM L-glutamate) into the vaginal vault. Uninfected mice did not receive SPG-buffer into the vaginal vault, since previous studies from our laboratory

have observed no inflammation or no uterine pathology from SPG buffer. Beginning one week later, uninfected and infected mice were fed a high-fat (*Apoe*^{-/-}) or a high-fat/high cholesterol diet (*Ldlr*^{-/-}) for 7 weeks (Fig 1A). Mice were euthanized at 8 weeks (56-days) post-infection. Chlamydial shedding was determined by swabbing the cervix and vaginal vault of infected mice with a calcium alginate swab (Spectrum Medical Instruments, Los Angeles, CA) at various times post-infection, and IFUs in the swabs were determined as described in an earlier study¹⁹.

Genital histopathology:

Genital tract tissues were extracted *en bloc* from mice and tissues were fixed in 10% buffered formalin and embedded in paraffin. Longitudinal sections (4 mm) were stained with hematoxylin and eosin (H&E) and evaluated by a pathologist to whom experimental design was not disclosed. Scoring was done as described in an earlier study¹⁹. Luminal distention of the uterine horns and dilatation of the oviducts was graded from 1 to 4, with a grade 4 representing peak severity or frequency of the parameter.

Atherosclerotic lesions:

The heart and descending aorta were excised and fixed in PBS/4% formalin/30% sucrose overnight before mounted in optimal cutting temperature medium and frozen at -70°C. Aortic sinus cryosections (8 µm) were stained with Oil Red O²⁰. Atherosclerotic lesions were quantified as previously described²⁰, by determining the lesion area in each of the five sections for each mouse. *En face* analysis of the descending aorta was performed after staining descending aorta with Oil Red O²⁰.

Plasma lipids, SAA, and cytokines:

Concentrations of plasma total cholesterol was determined using kits from Biovision (Milpitas, CA) as previously described²⁰. Plasma TNF-α and serum amyloid A-1 (SAA-1) levels were determined by sandwich ELISA using kits specific for mouse TNF-α (RND Systems) and SAA-1 (Invitrogen). A HEK-IL-1 receptor reporter cell line (HEK-IL-1Ra, Invivogen) was used to estimate the functionally active mature form of mouse IL-1β in the plasma. Plasma samples from uninfected and infected *Apoe*^{-/-} and *Ldlr*^{-/-} mice were used in this bioassay along with recombinant mouse IL-1β as a standard. A HEK-IL-1Ra cell line expressing IL-1Ra linked to secretory alkaline phosphatase (SEAP) as a reporter was treated for 16 h with plasma from chlamydia infected *Apoe*^{-/-} and *Ldlr*^{-/-} mice. SEAP in the supernatant collected from the HEK-IL-1Ra cell line was quantified using Quantibblue substrate (Invivogen).

Immunohistochemical analysis:

Serial aortic sinus cryosections (8 µm) were stained with goat anti-mouse vascular cell adhesion molecule-1 (VCAM-1) IgG (10 µg/ml, RND systems) using Vectastain ABC reagent (Vector Laboratories Inc.). The sections were developed with DAB (3,3'-diaminobenzidine) and counterstained with Mayer's hematoxylin. Images were captured using an Olympus microscope. Sections stained with goat IgG were used as a non-specific IgG control. The percentage of VCAM-1+ staining area was determined by measuring

the total aortic sinus area. Macrophages at the lesion site were characterized by staining cryosections with the MOMA-2 antibody²².

Chlamydia at the lesion site:

DNA was prepared from the aorta, including the aortic arches, of Chlamydia-infected *Apoe*^{-/-} mice. Detection of chlamydial DNA was determined by quantitative PCR using *C. muridarum*-specific primers for gene coding for 16s rRNA as previously described²³. Purified chlamydial DNA was used as a positive control.

Bone marrow-derived macrophages:

Bone marrow-derived macrophages (BMDM) were generated from WT, *Apoe*^{-/-} and *Ldlr*^{-/-} mice, as previously described²⁰. After 6–8 days in culture, BMDM (0.8×10^5 /well) were infected with *C. muridarum* (1 MOI). Total RNA was isolated from uninfected and infected BMDM, and expression of cytokines and chemokines were determined by quantitative RT-PCR (qRT-PCR), using primers described in earlier studies^{20, 24, 25}, and in Supplemental Table 1. To confirm infection by observation of chlamydial inclusions, BMDM (0.8×10^5 /well) were seeded onto coverslips, infected in parallel, and fixed with methanol for 10 min at room temperature at 24-hour post infection. Coverslips were stained for chlamydial inclusion using mouse anti-chlamydial immune sera, followed by anti-mouse alexaFluor-488 secondary antibody (Invitrogen) and mounted with ProLong Antifade (Invitrogen)¹⁹.

Statistical analyses:

Columns and error bars represent mean $\pm$ SD. Differences between two groups were considered significant when $p < 0.05$ using the two-tailed Student *t* test. Infection course was analyzed by 2-way ANOVA and significance of uterine pathological scores was determined using the Mann-Whitney *U* test. All data were analyzed using InStat version 3.1a for Macintosh (GraphPad, San Diego, CA).

Results

C. muridarum genital infection results in increased aortic lesions in both *Apoe*^{-/-} and *Ldlr*^{-/-} mice:

Apoe^{-/-} and *Ldlr*^{-/-} mice, fed a hyperlipidemic diet, were used to determine whether genital chlamydial infection accelerates the progression of atherosclerosis. Infection course was assessed by chlamydial IFU measurements in genital swabs from *Apoe*^{-/-} and *Ldlr*^{-/-} mice and found to be comparable to WT mice (Supplemental Fig 1A and B). Both knockout mouse strains cleared infection by day 25 and IFUs were not detected in the swabs, thereafter.

To determine whether atherosclerosis was exacerbated in mice infected with *C. muridarum*, atherosclerotic lesions were compared between infected and uninfected mice at 8 weeks post-infection that were given hyperlipidemic diet starting a week after infection (Fig. 1A). Larger lesion areas were observed in the aortic sinuses of infected *Apoe*^{-/-} mice (Fig. 1B), which was significantly different between uninfected and infected mice (Fig. 1C). *En face* analysis of the descending aorta, including aortic arch, also showed significantly increased

lesions in mice that were infected with *C. muridarum* (Fig. 1D and E). *Ldlr*^{-/-} mice were used as another model to independently test and confirm the finding from *Apoe*^{-/-} mice. The aortic sinus atherosclerotic lesion area in infected *Ldlr*^{-/-} mice was significantly larger (50% increase) than in the uninfected *Ldlr*^{-/-} mice (Fig. 2AB). *En face* analysis showed a similar increase in descending aorta of infected mice (Fig 2C). At this early time point (7 weeks on a high fat diet), Chlamydia-infected wild type mice (fed a high fat diet) did not show detectable atherosclerotic lesions (data not shown). These observations demonstrate that genital chlamydial infection exacerbates atherosclerosis in *Apoe*^{-/-} and *Ldlr*^{-/-} mice, two well-characterized hyperlipidemic mouse models.

***Chlamydia* infection induces systemic and arterial inflammation.**

Cholesterol levels were similar in uninfected and infected *Apoe*^{-/-} mice, and similar results were observed in *Ldlr*^{-/-} mice (Supplemental Fig. 2A and 2B). Therefore, increased atherosclerotic lesions in infected mice could not be attributed to plasma cholesterol levels. To evaluate if systemic inflammatory responses²⁶ were different in uninfected and infected mice, SAA-1, TNF- α , and IL-1 β levels were measured in the plasma at 8-weeks post infection. SAA-1, TNF- α , and IL-1 β levels were elevated in the plasma of *Chlamydia*-infected *Apoe*^{-/-} mice (Fig. 3A–C). Plasma levels of bioactive IL-1 β were also significantly higher in *Chlamydia*-infected *Ldlr*^{-/-} mice compared to the corresponding uninfected mice at same time post infection (Fig. 3D). To determine if inflammatory responses were different in the vascular endothelium between uninfected and infected mice, we analyzed the expression of an inflammation marker; VCAM-1 at the lesion site in the aortic sections. VCAM-1 served as a candidate inflammatory marker as VCAM-1-dependent monocyte adhesion and transmigration promote formation of fatty streak lesions²⁷. Significantly more VCAM-1+ staining was observed in the aortic lesions of infected *Apoe*^{-/-} mice compared with uninfected *Apoe*^{-/-} mice (Fig. 4A and B). A similar trend was also observed in infected *Ldlr*^{-/-} mice, compared with uninfected *Ldlr*^{-/-} mice (Fig. 4C and D). Immunohistochemical analyses of the aortic sinus using MOMA2 antibody showed more macrophages in aortic lesions of infected *Apoe*^{-/-} mice, compared to uninfected mice (Fig. 4E). These findings suggest that chlamydial infection-induced systemic inflammatory response could induce endothelial cell activation, resulting in increased adherence, and transmigration of monocytes at the lesion site in infected mice.

Clinical studies have shown that *C. pneumoniae* lung infections result in dissemination of *C. pneumoniae* at the vascular site^{28, 29}. Therefore, we investigated whether increased incidence of lesions observed in hyperlipidemic mice is caused by dissemination of *C. muridarum* to the vascular site. Quantitative PCR analyses of DNA from the aortic arch at day 8 weeks post-infection did not show chlamydial DNA (Supplemental Fig. 3A and B). These findings suggest that there is no persistent chlamydial infection in the lesion-prone vascular site. However, this does not preclude dissemination of *Chlamydia* to the vascular sites in early stages of infection.

Increased uterine pathology was observed in *C. muridarum*-infected *Apoe*^{-/-} mice.

Chlamydia infection leads to significant pathology of the oviduct, characterized by fluid filled hydrosalpinx. Additionally, some mice also develop uterine pathology with dilatation.

Excision of the genital tract at 8 weeks post-infection showed no significant differences in hydrosalpinx incidence between WT and *Ldlr*^{-/-} mice and between WT and *Apoe*^{-/-} mice, that were fed high fat diet (data not shown). However, infected *Apoe*^{-/-} mice developed severe uterine dilatation relative to infected WT mice (Fig. 5A). Histopathological observation of H&E stained section showed dilated uteruses in infected *Apoe*^{-/-} mice with severe thinning of uterine wall (Fig. 5B). To determine if feeding high fat diet exacerbated uterine pathology, in an independent study, *Apoe*^{-/-} mice were infected and fed a high fat or regular chow diets (non-hyperlipidemic diet). Histopathological scoring of genital tract tissues showed more uterine horns with a score of 4 in high fat diet fed *Apoe*^{-/-} mice. However, the differences in uterine dilatation between mice fed high fat or regular chow diet were not statistically different, although both were significantly increased compared to uninfected mice (Fig 5C). Histological scoring of inflammatory cells at this time revealed no significant differences in acute (neutrophils), chronic (mononuclear), or plasma cells in mice fed high fat (Fig. 5D) or regular diets (data not shown). Further, the severe uterine dilatation observed in infected *Apoe*^{-/-} mice, was not observed in infected *Ldlr*^{-/-} mice (data not shown). These data suggest that uterine pathology observed in infected *Apoe*^{-/-} mice is an outcome of absence of Apoe and not due to hyperlipidemia.

Inflammatory cytokine response to Chlamydia infection is increased in *Apoe*^{-/-} macrophages.

To determine if increased genital pathology in *Apoe*^{-/-} mice is a result of increased inflammatory response of immune cells to genital Chlamydia infection, we generated BMDM from WT and *Apoe*^{-/-} mice and their inflammatory gene expression in response to Chlamydia infection determined. Chlamydia-infected WT macrophages showed increased *Ifnb*, *Il1b*, *Il6*, and *Tnfa* expression (Supplemental Fig. 4A) compared to uninfected macrophages. Moreover, *Apoe*-deficient macrophages infected with Chlamydia showed higher pro-inflammatory cytokine response (Supplemental Fig. 4A) compared to infected WT macrophages. We then determined whether macrophages from two hyperlipidemic mouse models differ in cytokine response following Chlamydial infection. BMDM from *Apoe*^{-/-} and *Ldlr*^{-/-} mice were infected with Chlamydia, and their cytokine response to infection determined by qRT-PCR. Interestingly, some cytokine and chemokine expression of *Apoe*^{-/-} BMDM were significantly higher than those of *Ldlr*^{-/-} BMDM (Fig. 6). To rule out if the increased cytokine response was a result of increased infection, infected macrophages were stained for chlamydial inclusion. No differences in inclusion size or numbers were observed between *Apoe*^{-/-} and *Ldlr*^{-/-} mice BMDM macrophages (Supplemental Fig. 4B). These findings suggest that *Apoe*-deficiency augments chlamydial infection-induced pro-inflammatory cytokine response, which could contribute to genital pathology.

Discussion

In this study we addressed the hypothesis that genital chlamydial infection can accelerate atherosclerosis under hyperlipidemic conditions. Using two hyperlipidemic mouse models (*Apoe*^{-/-} and *Ldlr*^{-/-}), we observed increased atherosclerotic lesions following *Chlamydia* genital infection. The infected mice had enhanced systemic and local inflammation,

indicated by increased SAA, TNF- α , and IL-1 β in plasma, and upregulation of VCAM-1 at the lesion site. In addition, we also discovered a novel role for Apoe in the maintenance of uterine architecture after chlamydial infection is resolved in the genital tract.

There is significant but conflicting literature regarding *C. pneumoniae* lung infection promoting atherosclerosis in the mouse model. Earlier studies have reported increased atherosclerotic lesions in *Apoe*^{-/-} or *Ldlr*^{-/-} mice inoculated intra-nasally at multiple times with *C. pneumoniae*^{11, 12, 30}. However, other studies showed no differences in total lesion area^{31, 32}. Different *C. pneumoniae* strains, age of the mice, mouse strains used; the frequency and route of infections could be contributing to the conflicting reports. Unlike previous studies^{33, 34}, we have used genital *C. muridarum* infection model which requires a single inoculation to establish genital infection to examine genital *C. muridarum* infection contributing to atherosclerotic lesions in mice. Atherosclerotic lesions were analyzed as early as 8 weeks post-infection, including 7 weeks on a hyperlipidemic diet. Unlike other studies where lesions are measured at later times (> 12 weeks on a high-fat diet)^{31, 35}, we chose to collect samples at this early time when lesions are not fully developed in *Apoe*^{-/-} mice, to detect increases in lesion area post infection, before excessive lesions appeared in uninfected hyperlipidemic mice.

Bacterial infection or bacterial ligands regulate cholesterol homeostasis by regulating reverse cholesterol transporter expression³⁶. However, we found no differences in plasma cholesterol levels between infected and uninfected mice, indicating that the increased lesions in the infected groups are independent of plasma lipid levels. Infectious microbial agent-induced atherosclerosis could occur by direct effect through infection of vascular cells and/or indirectly by induction of inflammatory cytokines by infection at other sites, such as the aortic vascular endothelium. *C. pneumoniae* was detected within atherosclerotic plaques, suggesting an association between atherosclerosis and *C. pneumoniae* infection³⁷. A recent report has reported that intranasal administration of *C. muridarum* resulted in the infection of blood monocytes, which subsequently could transport *C. muridarum* to the vascular site³⁸. In our study, PCR analyses did not show chlamydial 16s ribosomal RNA in the aortic arch at 8 weeks post-infection. However, this does not rule out the possibility of early dissemination of *C. muridarum* to the mouse aorta, as we have observed early dissemination followed by rapid clearance of *C. muridarum* in the lungs and spleens of WT mice³⁹. Further, *C. muridarum* infection has been shown to persist in the gut well after it has been cleared from the genital tract^{40, 41}. Whether or not this persistence in the gut could have affected atherosclerotic lesions needs further investigation. Dissemination of *C. muridarum* is likely due to its ability to grow in macrophages⁴². However, with the exception of the less prevalent human Chlamydia strain like Lymphogranuloma venereum, genital strains of *C. trachomatis* grow poorly in human macrophages. Nevertheless, immunodeficiency or co-infection with other sexually transmitted infectious pathogens can compromise host immunity and allow *C. trachomatis* to survive in macrophages and disseminate to remote sites.

In mice, systemic inflammation induced by either intraperitoneal LPS injection or by cecal ligation puncture accelerated atherosclerosis by increased VCAM-1 expression and monocyte adhesion to the aorta^{43, 44}. Further, we have shown that *C. muridarum* infection in

Apoe^{-/-} and *Ldlr*^{-/-} mice increased expression of VCAM-1 and macrophage accumulation at the lesion site. We have also shown that chlamydial genital infection results in augmented systemic inflammatory response as evidenced by elevated levels of TNF- α and IL-1 β . Mechanistic studies have indicated that TNF- α severely accelerates lesion development by upregulating vascular cell adhesion molecules and CD36, one of the scavenger receptors implicated in foam cell formation and atherosclerosis^{45, 46}. These findings suggest that *C. muridarum*-induced production of TNF- α and IL1b could increase endothelial cell VCAM-1 expression resulting in increased monocyte adhesion, and subsequent accelerated atherosclerosis in infected mice^{27, 47}.

Genital *Chlamydia* infection in the mouse model leads to the development of fluid-filled oviduct pathology, described as hydrosalpinx. Occasionally, C57BL/6 mice also develop uterine cysts or swelling post-infection. However, in most animals, the uterine horns recover fully and resume an intact epithelial layer. A surprise finding in this study was that infected *Apoe*^{-/-} mice developed severe uterine pathology in addition to oviduct pathology. The uterine swelling led to complete loss of the epithelial architecture. We have reported similar uterine horn pathology in *Irf3*^{-/-} mice and suggested increased cellular proliferation as an associated phenotype⁴⁸. However, the exact mechanism for this phenotype is not clear. The uterine pathology observed in *Apoe*^{-/-} mice was not associated with a high fat diet, and not observed in *Ldlr*^{-/-} mice, suggesting an alternate function specifically for Apoe in uterus. Apoe is expressed in macrophages, and is essential for efficient cholesterol efflux⁴⁹. However, several new functions of Apoe have emerged. An anti-inflammatory role has been attributed to Apoe as it aids in clearance of apoptotic bodies by macrophages⁵⁰. In this report, we showed that *Chlamydia* infection augmented cytokine expression in Apoe-deficient macrophages compared to WT and Ldlr-deficient macrophages. These findings suggest that lack of anti-inflammatory function of Apoe may contribute to the augmented inflammatory cytokine response following *Chlamydia* infection. *Apoe*^{-/-} mice display impaired immunity to *Listeria monocytogene*⁵¹ and *Klebsiella pneumoniae*⁵² infection. Apoe has also been shown to inhibit cell proliferation and migration⁵³, suggesting that in the absence of Apoe, excessive proliferation of uterine stromal cells could lead to the observed dilatation.

In conclusion, our findings show that inflammatory response during a genital Chlamydia infection may be sufficient to promote the formation of atherosclerotic lesions under hyperlipidemic conditions in mouse models. To our knowledge, this is the first study to suggest a link between genital Chlamydia infection-induced PID and subclinical atherosclerosis, suggesting a causal link between genital infection and atherosclerosis. Further clinical studies are required to establish this association in humans. Additionally, the contribution of Apoe in uterine health during infection warrants further investigation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment.

The authors thank Dr. Hangpong Ng for his technical help.

Financial support: This work was supported by NIH HL086674, HL130126 (SN), AI067678, and AI124010-01 (UMN). Partly supported by intramural funds from the Department of Pathology and Vascular Medicine Institute, University of Pittsburgh, PA, and Department of Pathology and Laboratory Medicine and UNC Kidney Center, UNC at Chapel Hill, NC (SN).

References

- [1]. Libby P, Inflammation in atherosclerosis, *Nature*, 2002;420:868–874. [PubMed: 12490960]
- [2]. Moazed TC, Kuo C, Grayston JT, et al. , Murine models of *Chlamydia pneumoniae* infection and atherosclerosis, *J Infect Dis*, 1997;175:883–890. [PubMed: 9086145]
- [3]. Michelsen KS, Doherty TM, Shah PK, et al. , TLR signaling: an emerging bridge from innate immunity to atherogenesis, *J Immunol*, 2004;173:5901–5907. [PubMed: 15528321]
- [4]. Gibson FC 3rd, Hong C, Chou HH, et al. , Innate immune recognition of invasive bacteria accelerates atherosclerosis in apolipoprotein E-deficient mice, *Circulation*, 2004;109:2801–2806. [PubMed: 15123526]
- [5]. Saikku P, Leinonen M, Mattila K, et al. , Serological evidence of an association of a novel *Chlamydia*, TWAR, with chronic coronary heart disease and acute myocardial infarction, *Lancet*, 1988;2:983–986. [PubMed: 2902492]
- [6]. Thom DH, Grayston JT, Siscovick DS, et al. , Association of prior infection with *Chlamydia pneumoniae* and angiographically demonstrated coronary artery disease, *JAMA*, 1992;268:68–72. [PubMed: 1608116]
- [7]. Grayston JT, Kuo CC, Coulson AS, et al. , *Chlamydia pneumoniae* (TWAR) in atherosclerosis of the carotid artery, *Circulation*, 1995;92:3397–3400. [PubMed: 8521559]
- [8]. Thom DH, Wang SP, Grayston JT, et al. , *Chlamydia pneumoniae* strain TWAR antibody and angiographically demonstrated coronary artery disease, *Arterioscler Thromb*, 1991;11:547–551. [PubMed: 2029495]
- [9]. Tormakangas L, Erkkila L, Korhonen T, et al. , Effects of repeated *Chlamydia pneumoniae* inoculations on aortic lipid accumulation and inflammatory response in C57BL/6J mice, *Infect Immun*, 2005;73:6458–6466. [PubMed: 16177317]
- [10]. Blessing E, Campbell LA, Rosenfeld ME, et al. , *Chlamydia pneumoniae* infection accelerates hyperlipidemia induced atherosclerotic lesion development in C57BL/6J mice, *Atherosclerosis*, 2001;158:13–17. [PubMed: 11500169]
- [11]. Moazed TC, Campbell LA, Rosenfeld ME, et al. , *Chlamydia pneumoniae* infection accelerates the progression of atherosclerosis in apolipoprotein E-deficient mice, *J Infect Dis*, 1999;180:238–241. [PubMed: 10353889]
- [12]. Hu H, Pierce GN and Zhong G, The atherogenic effects of chlamydia are dependent on serum cholesterol and specific to *Chlamydia pneumoniae*, *J Clin Invest*, 1999;103:747–753. [PubMed: 10074493]
- [13]. Hauer AD, de Vos P, Peterse N, et al. , Delivery of *Chlamydia pneumoniae* to the vessel wall aggravates atherosclerosis in LDLr^{-/-} mice, *Cardiovasc Res*, 2006;69:280–288. [PubMed: 16112098]
- [14]. Workowski KA, Bolan GA, Centers for Disease C, et al. , Sexually transmitted diseases treatment guidelines, 2015, *MMWR Recomm Rep*, 2015;64:1–137.
- [15]. Eckert LO, Hawes SE, Wolner-Hanssen PK, et al. , Endometritis: the clinical-pathologic syndrome, *Am J Obstet Gynecol*, 2002;186:690–695. [PubMed: 11967492]
- [16]. Crichton J, Hickman M, Campbell R, et al. , Prevalence of chlamydia in young adulthood and association with life course socioeconomic position: birth cohort study, *PLoS One*, 2014;9:e104943. [PubMed: 25153124]
- [17]. Crichton J, Hickman M, Campbell R, et al. , Socioeconomic factors and other sources of variation in the prevalence of genital chlamydia infections: A systematic review and meta-analysis, *BMC Public Health*, 2015;15:729. [PubMed: 26224062]
- [18]. Nagarajan UM, Prantner D, Sikes JD, et al. , Type I interferon signaling exacerbates *Chlamydia muridarum* genital infection in a murine model, *Infect Immun*, 2008;76:4642–4648. [PubMed: 18663004]

- [19]. Nagarajan UM, Sikes JD, Yeruva L, et al. , Significant role of IL-1 signaling, but limited role of inflammasome activation, in oviduct pathology during *Chlamydia muridarum* genital infection, *J Immunol*, 2012;188:2866–2875. [PubMed: 22331066]
- [20]. Ng HP, Zhu X, Harmon EY, et al. , Reduced Atherosclerosis in apoE-inhibitory FcγRIIb-deficient mice Is associated with increased anti-inflammatory responses by T Cells and macrophages, *Arterioscler Thromb Vasc Biol*, 2015;35:1101–1112. [PubMed: 25792447]
- [21]. Schiller NK, Black AS, Bradshaw GP, et al. , Participation of macrophages in atherosclerotic lesion morphology in LDLr^{-/-} mice, *J Lipid Res*, 2004;45:1398–1409. [PubMed: 15175354]
- [22]. Thampi P, Stewart BW, Joseph L, et al. , Dietary homocysteine promotes atherosclerosis in apoE-deficient mice by inducing scavenger receptors expression, *Atherosclerosis*, 2008;197:620–629. [PubMed: 17950295]
- [23]. Zhang Y, Yeruva L, Marinov A, et al. , The DNA sensor, cyclic GMP-AMP synthase, is essential for induction of IFN-β during *Chlamydia trachomatis* infection, *J Immunol*, 2014;193:2394–2404. [PubMed: 25070851]
- [24]. Ng HP, Burris RL and Nagarajan S, Attenuated atherosclerotic lesions in apoE^{-/-}Fcγ-chain-deficient hyperlipidemic mouse model is associated with inhibition of Th17 cells and promotion of regulatory T cells, *J Immunol*, 2011;187:6082–6093. [PubMed: 22043015]
- [25]. Nagarajan UM, Sikes J, Prantner D, et al. , MyD88 deficiency leads to decreased NK cell gamma interferon production and T cell recruitment during *Chlamydia muridarum* genital tract infection, but a predominant Th1 response and enhanced monocytic inflammation are associated with infection resolution, *Infect Immun*, 2011;79:486–498. [PubMed: 21078858]
- [26]. Galkina E and Ley K, Vascular adhesion molecules in atherosclerosis, *Arterioscler Thromb Vasc Biol*, 2007;27:2292–2301. [PubMed: 17673705]
- [27]. Cybulsky MI, Iiyama K, Li H, et al. , A major role for VCAM-1, but not ICAM-1, in early atherosclerosis, *J Clin Invest*, 2001;107:1255–1262. [PubMed: 11375415]
- [28]. Ramirez JA, Isolation of *Chlamydia pneumoniae* from the coronary artery of a patient with coronary atherosclerosis. The *Chlamydia pneumoniae*/Atherosclerosis Study Group, *Annals of internal medicine*, 1996;125:979–982. [PubMed: 8967709]
- [29]. Hirono S and Pierce GN, Dissemination of *Chlamydia pneumoniae* to the vessel wall in atherosclerosis, *Mol Cell Biochem*, 2003;246:91–95. [PubMed: 12841348]
- [30]. Chen S, Shimada K, Zhang W, et al. , IL-17A is proatherogenic in high-fat diet-induced and *Chlamydia pneumoniae* infection-accelerated atherosclerosis in mice, *J Immunol*, 2010;185:5619–5627. [PubMed: 20935201]
- [31]. Caligiuri G, Rottenberg M, Nicoletti A, et al. , *Chlamydia pneumoniae* infection does not induce or modify atherosclerosis in mice, *Circulation*, 2001;103:2834–2838. [PubMed: 11401941]
- [32]. Ezzahiri R, Stassen FR, Kurvers HA, et al. , *Chlamydia pneumoniae* infection induces an unstable atherosclerotic plaque phenotype in LDL-receptor, ApoE double knockout mice, *European journal of vascular and endovascular surgery: the official journal of the European Society for Vascular Surgery*, 2003;26:88–95.
- [33]. Kelly KA, Walker JC, Jameel SH, et al. , Differential regulation of CD4 lymphocyte recruitment between the upper and lower regions of the genital tract during *Chlamydia trachomatis* infection, *Infect Immun*, 2000;68:1519–1528. [PubMed: 10678969]
- [34]. Nagarajan UM, Ojcius DM, Stahl L, et al. , *Chlamydia trachomatis* induces expression of IFN-γ-inducible protein 10 and IFN-β independent of TLR2 and TLR4, but largely dependent on MyD88, *J Immunol*, 2005;175:450–460. [PubMed: 15972679]
- [35]. Ezzahiri R, Nelissen-Vrancken HJ, Kurvers HA, et al. , *Chlamydia pneumoniae* (Chlamydia pneumoniae) accelerates the formation of complex atherosclerotic lesions in Apo E3-Leiden mice, *Cardiovasc Res*, 2002;56:269–276. [PubMed: 12393097]
- [36]. Azzam KM and Fessler MB, Crosstalk between reverse cholesterol transport and innate immunity, *Trends Endocrinol Metab*, 2012;23:169–178. [PubMed: 22406271]
- [37]. Assar O, Nejatizadeh A, Dehghan F, et al. , Association of *Chlamydia pneumoniae* Infection With Atherosclerotic Plaque Formation, *Glob J Health Sci*, 2015;8:260–267.

- [38]. Roufaiel M, Gracey E, Siu A, et al. , CCL19-CCR7-dependent reverse transendothelial migration of myeloid cells clears Chlamydia muridarum from the arterial intima, *Nat Immunol*, 2016;17:1263–1272. [PubMed: 27668800]
- [39]. Poston TB, O'Connell CM, Girardi J, et al. , T Cell-Independent Gamma Interferon and B Cells Cooperate To Prevent Mortality Associated with Disseminated Chlamydia muridarum Genital Tract Infection, *Infect Immun*, 2018;86.
- [40]. Shao L, Melero J, Zhang N, et al. , The cryptic plasmid is more important for Chlamydia muridarum to colonize the mouse gastrointestinal tract than to infect the genital tract, *PLoS One*, 2017;12:e0177691. [PubMed: 28542376]
- [41]. Shao L, Zhang T, Liu Q, et al. , Chlamydia muridarum with mutations in chromosomal genes *tc0237* and/or *tc0668* Is deficient in colonizing the mouse gastrointestinal tract, *Infect Immun*, 2017;85:429–417.
- [42]. Nagarajan UM, Tripathy M, Kollipara A, et al. , Differential signaling pathways are initiated in macrophages during infection depending on the intracellular fate of Chlamydia spp, *Immunol Cell Biol*, 2018;96:246–256. [PubMed: 29363185]
- [43]. Burris RL, Ng HP and Nagarajan S, Soy protein inhibits inflammation-induced VCAM-1 and inflammatory cytokine induction by inhibiting the NF-kappaB and AKT signaling pathway in apolipoprotein E-deficient mice, *Eur J Nutr*, 2013;53:135–148. [PubMed: 23468309]
- [44]. Kaynar AM, Yende S, Zhu L, et al. , Effects of intra-abdominal sepsis on atherosclerosis in mice, *Crit Care*, 2014;18:469. [PubMed: 25182529]
- [45]. Moore KJ and Freeman MW, Scavenger receptors in atherosclerosis: beyond lipid uptake, *Arterioscler Thromb Vasc Biol*, 2006;26:1702–1711. [PubMed: 16728653]
- [46]. Huh HY, Lo SK, Yesner LM, et al. , CD36 induction on human monocytes upon adhesion to tumor necrosis factor-activated endothelial cells, *J Biol Chem*, 1995;270:6267–6271. [PubMed: 7534309]
- [47]. McKellar GE, McCarey DW, Sattar N, et al. , Role for TNF in atherosclerosis? Lessons from autoimmune disease, *Nat Rev Cardiol*, 2009;6:410–417. [PubMed: 19421244]
- [48]. Prantner D, Sikes JD, Hennings L, et al. , Interferon regulatory transcription factor 3 protects mice from uterine horn pathology during Chlamydia muridarum genital infection, *Infect Immun*, 2011;79:3922–3933. [PubMed: 21788382]
- [49]. Curtiss LK and Boisvert WA, Apolipoprotein E and atherosclerosis, *Curr Opin Lipidol*, 2000;11:243–251. [PubMed: 10882339]
- [50]. Grainger DJ, Reckless J and McKilligin E, Apolipoprotein E modulates clearance of apoptotic bodies in vitro and in vivo, resulting in a systemic proinflammatory state in apolipoprotein E-deficient mice, *J Immunol*, 2004;173:6366–6375. [PubMed: 15528376]
- [51]. Roselaar SE and Daugherty A, Apolipoprotein E-deficient mice have impaired innate immune responses to *Listeria monocytogenes* in vivo, *J Lipid Res*, 1998;39:1740–1743. [PubMed: 9741685]
- [52]. de Bont N, Netea MG, Demacker PN, et al. , Apolipoprotein E knock-out mice are highly susceptible to endotoxemia and *Klebsiella pneumoniae* infection, *J Lipid Res*, 1999;40:680–685. [PubMed: 10191292]
- [53]. Hui DY and Basford JE, Distinct signaling mechanisms for apoE inhibition of cell migration and proliferation, *Neurobiol Aging*, 2005;26:317–323. [PubMed: 15639309]

Highlights

Genital chlamydia infection exacerbated atherosclerosis in *ApoE*^{-/-} and *Ldlr*^{-/-}, two independent hyperlipidemic mouse models of atherosclerosis.

Chlamydia infection increased circulating levels of TNF- α and IL-1 β .

Chlamydia infection also increased VCAM-1 expression and macrophages in atherosclerotic lesions.

Chlamydia infection increased pro-inflammatory cytokines and chemokines mRNA expression.

Interestingly, Chlamydia infection showed uterine pathology only in apoE-deficient mice.

Fig 1.

Aortic lesions were increased in *Apoe*^{-/-} mice with intra vaginal infection of *C. muridarum*. *Apoe*^{-/-} mice (N=10) were infected with 3×10^5 IFU vaginally and were fed high fat diet starting one week after infection. Mice were sacrificed 8-weeks post infection (A). Aortic sinus (N=6) and descending aorta (N=10) were stained using oil red O to visualize plaques and average lesion area plotted. Aortic sinus from a representative mouse from each group is shown and (B) lesion area in aortic sinus (C) for all mice determined. Descending aorta from a representative mouse from each group is shown (D) and lesion area in descending aorta (E) for all mice determined. Significance determined by 't' test. Figure represents one of two independent experiments.

Fig 2. Aortic lesions were increased in *Ldlr*^{-/-} mice infected with *C. muridarum*. *Ldlr*^{-/-} mice (N=10) were infected with 3×10^5 IFU vaginally and were fed high fat/high cholesterol diet starting one week after infection (as shown in Figure 1A). Mice were sacrificed 8-weeks post infection. Aortic sinus (N=5) was stained using oil red O to visualize plaques. Aortic sinus (A) from a representative mouse from each group is shown and average lesion area (B) plotted. *En face* analysis of descending aorta from a representative mouse from each group is shown (C). Uninfected *Ldlr*^{-/-} mice fed a high-fat/high cholesterol diet was used as control. Significance determined by 't' test.

Figure 3. Increased levels of SAA, TNF-α, and IL-1β in the plasma of infected hyperlipidemic mice.

Apoe^{-/-} mice (N=10) or *Ldlr*^{-/-} were infected with 3×10^5 IFU genitally, and one week after infection mice were fed hyperlipidemic diet. Mice were sacrificed 8-weeks post infection (7 weeks on a high fat diet). Plasma from uninfected mice were used as controls. Plasma levels of SAA-1 (A), TNF-α (B) in *Apoe*^{-/-} mice, and IL-1β in *Apoe*^{-/-} (C), and *Ldlr*^{-/-} (D) were determined as described in methods. Significance determined by 't' test.

Fig 4.

Increased VCAM-1 expression and macrophages in *Chlamydia*-infected mice.

Representative aortic sinus sections from uninfected or *Chlamydia*-infected *Apoe*^{-/-} (A, B) and or *Ldlr*^{-/-} (C, D) were stained with anti-mouse VCAM-1 IgG to detect VCAM-1 protein expression at the lesion site. Arrows indicate VCAM-1 positive staining. (E), Macrophages in aortic sections from uninfected and *Chlamydia*-infected or *Apoe*^{-/-} mice was determined as described the Methods section. Under similar conditions, aortic sections incubated with an isotype-matched rat IgG control were minimal. Representative of five aortic sinus sections from uninfected and chlamydia-infected *Apoe*^{-/-} or *Ldlr*^{-/-} mice are presented (100 × magnification).

Fig 5. Severe uterine dilatation in *Apoe*^{-/-} mice in comparison with WT mice.

(A), Representative photograph of genital tracts from infected WT and *Apoe*^{-/-} mice (N=10). (B) H&E staining of longitudinal sections of genital tracts WT and *Apoe*^{-/-} mice (N=5), and (C) histopathological scoring for immune cells and dilatation (N=5). (D) Dilatation scores from an independent experiment where 5 mice from each group were fed high fat diet or fed a regular chow diet. Statistics were determined by non-parametric two-tailed Mann Whitney test.

Fig 6.

Augmented inflammatory responses in Chlamydia infected *Apoe*^{-/-} macrophages. BMDM from *Apoe*^{-/-} and *Ldlr*^{-/-} were infected with *C. muridarum* (1 MOI) for 8 h, and expression of inflammatory cytokine and chemokine responses was determined by qRT-PCR. Uninfected macrophages were used as controls. Values are mean $\pm$ SD of triplicates. Representative two independent experiments are presented. Cytokine expression in Chlamydia infected *Apoe*^{-/-} BMDM was compared to infected *Ldlr*^{-/-} BMDM. Significance determined by 't' test.