ESO guideline for the management of extracranial and intracranial artery dissection

European Stroke Journal
2021, Vol. 6(3) XXXIX–LXXXVIII
© European Stroke Organisation 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/23969873211046475
journals.sagepub.com/home/eso

Stephanie Debette^{1,2} , Mikael Mazighi^{3,4,5,6,7}, Philippe Bijlenga⁸, Alessandro Pezzini⁹, Masatoshi Koga¹⁰, Anna Bersano¹¹, Janika Kõrv^{12,13} , Julien Haemmerli⁸, Isabella Canavero¹¹, Piotr Tekiel¹⁴, Kaori Miwa¹⁰, David J Seiffge¹⁵, Sabrina Schilling¹⁶, Avtar Lal¹⁶, Marcel Arnold¹⁵, Hugh S Markus¹⁷, Stefan T Engelter^{18,19} and Jennifer J Majersik¹⁴

Abstract

The aim of the present European Stroke Organisation guideline is to provide clinically useful evidence-based recommendations on the management of extracranial artery dissection (EAD) and intracranial artery dissection (IAD). EAD and IAD represent leading causes of stroke in the young, but are uncommon in the general population, thus making it challenging to conduct clinical trials and large observational studies. The guidelines were prepared following the Standard Operational Procedure for European Stroke Organisation guidelines and according to GRADE methodology. Our four recommendations result from a thorough analysis of the literature comprising two randomized controlled trials (RCTs) comparing anticoagulants to antiplatelets in the acute phase of ischemic stroke and twenty-six comparative observational studies. In EAD patients with acute ischemic stroke, we recommend using intravenous thrombolysis (IVT) with alteplase within 4.5 hours of onset if standard inclusion/exclusion criteria are met, and mechanical thrombectomy in patients with large vessel occlusion of the anterior circulation. We further recommend early endovascular or surgical intervention for IAD patients with subarachnoid hemorrhage (SAH). Based on evidence from two phase 2 RCTs that have shown no difference between the benefits and risks of anticoagulants versus antiplatelets in the acute phase of symptomatic EAD, we strongly recommend that clinicians can prescribe either option. In post-acute EAD patients with residual stenosis or dissecting aneurysms and in symptomatic IAD patients with an intracranial dissecting aneurysm and isolated headache,

¹Bordeaux Population Health research center, INSERM U1219, University of Bordeaux, Bordeaux, France

²Department of Neurology and Institute for Neurodegenerative Diseases, Bordeaux University Hospital, France

³Department of Neurology, Hôpital Lariboisière, Paris, France

⁴Interventional Neuroradiology Department, Hôpital Fondation Ophthalmologique Adolphe de Rothschild, Paris, France

⁵Université de Paris, Paris, France

⁶FHU NeuroVasc, Paris, France

⁷Laboratory of Vascular Translational Science, INSERM U1148, Paris, France

⁸Neurosurgery, Département de Neurosciences Cliniques, Hôpitaux Universitaires et Faculté de Médecine de Genève, Switzerland

⁹Department of Clinical and Experimental Sciences, Neurology Clinic, University of Brescia, Brescia, Italy

¹⁰Department of Cerebrovascular Medicine, National Cerebral and Cardiovascular Center, Suita, Osaka, Japan

¹¹Fondazione IRCCS Istituto Neurologico 'Carlo Besta', Milano

¹²Department of Neurology and Neurosurgery, University of Tartu, Tartu, Estonia

¹³Department of Neurology, Tartu University Hospital, Tartu, Estonia

¹⁴Department of Neurology, University of Utah, Salt Lake City, UT, USA

¹⁵University Hospital Bern, University of Bern, Freiburgstrasse, Bern, Switzerland

¹⁶Guidelines Methodologist, European Stroke Organization, Basel, Switzerland

¹⁷Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK

¹⁸Department of Neurology and Stroke Center, University Hospital and University of Basel, Basel, Switzerland

¹⁹Neurology and Neurorehabilitation, University Department of Geriatric Medicine FELIX PLATTER, University of Basel, Basel, Switzerland

Corresponding author:

Stephanie Debette, Bordeaux Population Health research center, INSERM U1219, University of Bordeaux, 146 rue Léo Saignat, Bordeaux 33076, France.

Email: stephanie.debette@u-bordeaux.fr

there is insufficient data to provide a recommendation on the benefits and risks of endovascular/surgical treatment. Finally, nine expert consensus statements, adopted by 8 to 11 of the 11 experts involved, propose guidance for clinicians when the quality of evidence was too low to provide recommendations. Some of these pertain to the management of IAD (use of IVT, endovascular treatment, and antiplatelets versus anticoagulation in IAD with ischemic stroke and use of endovascular or surgical interventions for IAD with headache only). Other expert consensus statements address the use of direct anticoagulants and dual antiplatelet therapy in EAD-related cerebral ischemia, endovascular treatment of the EAD/IAD lesion, and multidisciplinary assessment of the best therapeutic approaches in specific situations.

Keywords

Extracranial artery dissection, cervical artery dissection, intracranial artery dissection, stroke, subarachnoid hemorrhage, endovascular treatment, aspirin, anticoagulants, thrombolysis, mechanical thrombectomy

Date received: 17 July 2021; accepted: 26 August 2021

Introduction

Cervico-cephalic artery dissections are defined by a hematoma in the wall of a cervical or intracranial artery and represent a leading cause of stroke in the young.^{1,2} While dissection of the extracranial cervical arteries (hereafter referred to as “extracranial artery dissection”, EAD) has been extensively studied and described in increasingly large observational studies,^{3–9} less data is available about isolated intracranial artery dissection (IAD). By contrast with cervical arteries, intracranial arteries are characterized by a paucity of elastic fibers in the media, little adventitial tissue, no external elastic lamina,^{10,11} and weaker supporting tissues.¹² In internal carotid arteries, the external elastic lamina is still present in the petrous portion of the temporal bone (C3 segment) and disappears in the horizontal segment of the cavernous portion (C5); hence, typical features of IAD are found for dissections occurring in the intradural portion of the internal carotid artery—i.e., starting in C6.¹³ In vertebral arteries, the reduction of elastic fibers in the media and external elastic lamina is most pronounced in the last 0.5 cm before the intradural portion but is not complete until 0.5 cm after dural perforation (V4 segment).¹³ While many characteristics and clinical manifestations are shared between EAD and IAD, there are also important differences. For example, the mean age of occurrence is younger (44 years) for EAD than for IAD (50 years).¹³

The incidence of EAD is low in the general population, estimated around 2.6 to 3.0 / 100,000 inhabitants/year.^{14,15} The incidence of IAD is unknown, but probably lower than that of EAD in populations of European ancestry.^{14,15} The relative frequency of EAD versus IAD varies by study recruitment strategies and ascertainment methods¹³ but is also highly variable according to geographic origin. For example, the proportion of IAD amongst all cervico-cephalic dissections is estimated around 11% in European populations,¹⁶ around 27% in Latin America,¹⁷ and up to 67–78% in East Asia.^{18,19} EAD/IAD are multifactorial

conditions in the vast majority of cases, while they can rarely occur as part of an inherited connective tissue disorder or fibromuscular dysplasia.^{8,9} In EAD, hypertension and migraine are reported risk factors with common genetic risk variants in the PHACTR1 gene also associated with EAD.^{7,20} Frequently cervical trauma (most often minor) or recent infections are reported as triggers. Risk factors are less well characterized for IAD.

Clinically, EAD usually presents with “local” symptoms and signs partly due to compression of adjacent structures, including headache, cervical pain, Horner syndrome, and cranial nerve palsy. In about two thirds to three quarters of patients in published series, the EAD is complicated by cerebral ischemia (ischemic stroke or transient ischemic attack [TIA]), or more seldom retinal, or spinal cord ischemia, typically occurring several hours or days after the onset of local symptoms. Subarachnoid hemorrhage (SAH) exceptionally occurs in EAD when the dissection expands to the intradural portion of the artery.⁹ The two main clinical manifestations of IAD are SAH (in about 50–60% of patients in published series), cerebral ischemia (30%–78%), and exceptionally both. About 80% of IAD patients have prodromal headache.^{21,22} Symptoms related to brainstem or cranial nerve compression can also occur.¹³

From an imaging perspective, EAD/IAD can present as a segmental stenosis (most common presentation in EAD), an occlusion, or a dissecting aneurysm. Several features can be present at the same time, typically a dissecting aneurysm with a long tapering stenosis. In 15–20% of EAD patients, multiple cervical arteries are affected.²³ In EAD, stenosis resolution or recanalization occurs in 33–90% within 6 months;^{5,15,24–26} dissecting aneurysms are reported to resolve or decrease in size in 40–50% of patients but can also increase in size and develop anew.^{27,28} In the Cervical Artery Dissection in Stroke Study (CADISS) trial, dissecting aneurysms were present in 24 of 264 patients at baseline and in 36 of 248 patients with follow-up at 3 months: 12 of baseline dissecting aneurysms persisted and

24 new dissecting aneurysms had developed.²⁸ The time-frame of changes in imaging characteristics in IAD patients and the rates of recanalization are unknown.¹³

The low incidence of EAD/IAD has proven a challenge to clinical trials conduction, and therefore guidelines have mostly relied on indirect evidence from observational studies and expert opinion. In addition, most guidelines have focused primarily on EAD management without discussion of the treatment of IAD.²⁹⁻³¹ Recently, two randomized controlled trials (RCTs) have been published comparing anticoagulation to antiplatelet therapy in the acute phase of EAD,³²⁻³⁴ one of them while this guideline was being prepared.³⁴ Moreover, acute phase ischemic stroke management has undergone substantial developments in recent years which existing EAD/IAD guidelines have only partially included. To our knowledge, this is the first guideline taking a comprehensive approach to both EAD and IAD and the first to use the GRADE system (Grading of Recommendations Assessment, Development, and Evaluation), which allows clear separation of quality of evidence and strength of recommendation as well as a transparent process of literature search and analysis.³⁵

The aim of this guideline is to provide recommendations to guide physicians treating patients with EAD/IAD to reach therapeutic decisions when assessing patients with a suspected or confirmed EAD or IAD. It first addresses treatment in the first hours of ischemic stroke due to EAD/IAD and the question of recanalization, second the acute phase treatment of IAD for prevention of SAH and ischemic stroke, and third the prevention of longer term complications of EAD/IAD.

Methods

The guidelines for management of EAD and IAD follow the standard operations procedure (SOP) defined by the European Stroke Organisation (ESO)³⁶ and were developed using the GRADE methodology.³⁵ A Module Working Group (MWG) was established, consisting of 11 experts (SD, Chair; AP, MM, MA, PB, MK, HSM, STE, JK, AB, and JJM). The MWG was joined by five fellows during study screening (DS, KM, IC, PT, and JH) who assisted the experts with abstract and full text screening and drafting the text. The MWG included ten neurologists (of whom one is also a neurointerventionalist and one a neuro-epidemiologist) and one neurosurgeon; all eleven are experts in cerebrovascular disease with a special interest in EAD and IAD or stroke in the young. Four of the fellows were trainee or early career neurologists and one was an early career neurosurgeon. Of the 16 MWG members, 12 were European, two were Japanese, and two from the United States; this wider geographical representation was important to account for differences in epidemiological characteristics and management strategies between

continents. ESO guidelines board and ESO Executive committee approved the composition of the working group. All participants were asked to disclose any conflict of interest that could influence their participation. The group communicated using e-mail and teleconferences.

Diagnostic criteria

For cervical artery dissection, we used the term extracranial artery dissection (EAD) for clear differentiation from intracranial artery dissection (IAD). EAD refers to the dissection of a cervical carotid or vertebral artery radiologically confirmed by the presence of a mural hematoma, a dissecting aneurysm, a long tapering stenosis, an intimal flap, a double lumen, or an occlusion >2 cm above the carotid bifurcation revealing a dissecting aneurysm and/or a long tapering stenosis after recanalization.²³ The diagnosis of IAD is considered definite in presence of at least one of the following:¹³ (i) a stenosis or occlusion of an intracranial artery secondarily developing towards a fusiform or irregular aneurysmal dilation at a non-branching site; (ii) an intramural hematoma, intimal flap, or double lumen; and (iii) pathological confirmation of IAD. Of note, given the dearth of data, we did not limit our review to studies strictly applying these diagnostic criteria. Indeed, especially for IAD, these diagnostic criteria are fairly recent,¹³ and restricting our search to later studies only would have substantially reduced the number of available studies. We did not include studies on mycotic and blood blister-like aneurysms. Mycotic or oncological giant fusiform aneurysms are caused by the release of proteases by bacteria or tumor cells that break down the vessel wall but are non-dissecting. Blood blister-like aneurysms are located at non-branching sites of intracranial arteries and are caused by a degeneration of the internal elastic lamina and media without associated arterial dissection.¹³

Selection of Population, Intervention, Comparator, and Outcome (PICO)

Regarding the population, the MWG decided to focus primarily on symptomatic EAD and IAD, with ischemia (ischemic stroke, TIA, or retinal ischemia), SAH, or headache.

Interventions and comparators addressed the early acute, acute, and post-acute phase of EAD/IAD. First, we addressed recanalization at the hyperacute phase of ischemic stroke caused by EAD/IAD using intravenous thrombolysis (PICO 1) and endovascular treatment (PICO2) (versus the absence of such treatment). Second, we tackled the acute phase of IAD without cerebral ischemia, encompassing IAD with SAH (PICO3) and IAD with only headache (PICO4) for which we each assessed endovascular or surgical intervention versus medical treatment. Third, we addressed the acute phase of EAD/IAD without SAH using

anticoagulation versus antiplatelet agents (PICO5). Fourth, we considered endovascular or surgical intervention versus medical treatment for residual stenosis or dissecting aneurysm beyond the acute phase of EAD (PICO6).

We considered six separate outcomes: (i) death, (ii) functional outcome (good functional outcome defined as modified Rankin Scale [mRS] scores of 0–2 versus 3–6 and excellent functional outcome with mRS scores of 0–1 versus 2–6, or equivalent as defined in the individual studies), (iii) ischemic stroke, (iv) SAH, (v) intracerebral hemorrhage (ICH), and (vi) major bleeding, defined according to the International Society on Thrombosis and Haemostasis (ISTH).³⁷ We had initially also considered the following additional outcomes: new ischemic lesions on diffusion weighted imaging (DWI), recurrent dissection, and frequency of normalization or stability of vessel patency on imaging. Using the Delphi method, the MWG voted in a closed ballot to identify which outcomes were of highest priority, according to the GRADE methodology using a 9-point scale (7–9: critical; 4–6: important; 1–3: of limited importance). The final scores, based on the mean votes from all participants, were the following: death 8.8, functional outcome (good or excellent functional outcome) 8.7, ischemic stroke 8.0, ICH 7.6, SAH 7.5, new ischemic lesions on DWI 6.5, recurrent dissection 6.2, normalization or stability of vessel patency 5.5, major bleeding 5.0. Due to a large number of outcomes we decided to focus on the clinical outcomes only and discarded the imaging-based outcomes that had been rated by the MWG as important, but not critical for decision making.

PICO questions

The MWG formulated six main PICO (Population, Intervention, Comparator, Outcome) questions relevant for EAD and IAD management, each with several sub-questions relating to the six different outcomes defined above, different subpopulations, or intervention sub-types, as relevant to each PICO and described below in the PICO header questions ([Supplementary Panel 1](#)). These were refined following comments from the ESO Executive Committee and ESO Guidelines Board. Subsequently, ESO Executive Committee and ESO Guidelines Board approved them.

For each PICO question, search terms were identified, tested, refined, and agreed by the MWG with the ESO Guidelines methodologist (AL). Search terms are listed in the Supplementary Methods.

Identification and selection of relevant studies

A systematic review of literature was done to collect evidence to answer the PICO questions. This search was performed by the ESO Guidelines methodologist (AL). The following databases were searched: MEDLINE, EMBASE

and CINAHL, from inception to April 5, 2021. We also searched reference lists of review articles, the authors' personal reference libraries, and previous guidelines for additional relevant records. The search results were loaded into the web-based Covidence platform (Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia) for assessment by the MWG. Two or more MWG members were assigned to independently screen the titles and abstracts of publications registered in Covidence and assess the full text of studies determined to be potentially relevant. All disagreements were resolved by discussion between the two authors or by a third MWG author. We prioritized RCTs, but due to the limited data, we also considered health registry data analyses, large observational studies (minimum size: 50 subjects for EAD, 20 subjects for IAD), and systematic reviews or individual patient data meta-analyses of observational studies. We chose a more liberal minimal sample size for IAD studies based on the dearth of published data on that disease. We considered only studies in human adults (>18 years) with the full article available in English. We excluded studies on penetrating injury of cervical or intracranial arteries and publications with only conference abstracts available.

Meta-analyses and assessment of quality and risk of bias

A random effects meta-analysis was conducted using the Review Manager (RevMan) 5.3 COCHRANE Collaboration software based on raw numbers extracted from the manuscripts. Results were presented as odds ratios (ORs) with 95% confidence intervals (CIs). Since there were very few RCTs and for some PICOs also few observational comparative studies, we also derived mean event rates from large single-arm observational studies using the metaprop function in the Meta package in R.

The risk of selection, performance, detection, attrition, and reporting biases in each RCT was assessed using the Cochrane Collaboration's tool,³⁸ and heterogeneity across studies was assessed using Cochran's Q (reported as a *p* value) and *I*² statistics. The Cochrane Collaboration's tool was used to perform the assessment of risk of bias of RCT. The various components of this tool, such as risk of selection (randomization, allocation concealment), performance (blinding of participants and personal), detection (blinding of outcome assessment), attrition (incomplete outcome data), and reporting (selective reporting) bias were assessed in each RCT. For non-RCTs, study conduct, subject selection, assessment, and statistical confounding were assessed using the Scottish Intercollegiate Guidelines Network (SIGN) checklist (<https://www.sign.ac.uk/what-we-do/methodology/checklists/>). Moreover, for each PICO question and each outcome, the quality of evidence was rated using the GRADEpro Guideline Development

Tool (McMaster University, 2015; developed by Evidence Prime, Inc.) using guidelines for non-pooled data as necessary,³⁹ as high, moderate, low, or very low³⁶ by AL and verified by at least two members of the MWG and approved by the rest.

Data analysis, drafting of available evidence and recommendations

Each PICO writing group, comprising at least three MWG members, analyzed the available data and drafted two sections of text: “analysis of current evidence” which focused on relevant RCTs and/or observational studies and “additional information” to summarize indirect evidence from additional studies. Each PICO writing group formulated an “evidence-based recommendation” according to the GRADE evidence profiles and the ESO standard operating procedure,³⁶ and/or an “expert consensus statement” if the PICO group considered that not enough evidence was available for replacing an evidence-based recommendation to address specific situations. The expert consensus statements were then voted on by all expert MWG members (excluding the fellows and methodologists). These expert consensus statements should not be regarded as evidence-based recommendations, since they only reflect the opinions of the MWG.

The Guidelines document was reviewed by all MWG members and modified using a Delphi approach until consensus was reached. It was also reviewed by a member of the ESO Guidelines Board who served as External Advisor prior to submission. The document was reviewed and approved by five reviewers (2 members of the ESO Guidelines Board, 1 Executive Committee member, and 2 external reviewers).

Results

PICO 1: In extracranial artery dissection (EAD) or intracranial artery dissection (IAD) patients with acute ischemic stroke is intravenous thrombolysis (IVT) versus no intravenous thrombolysis associated with a reduced risk of death, a higher likelihood of favorable functional outcome (mRS 0–2 vs. 3–6, or 0–1 vs. 2–6, or equivalent), and no increased risk of ICH, subarachnoid hemorrhage (SAH), or any major bleeding?

Analysis of current evidence

Intravenous thrombolysis is effective and safe in acute ischemic stroke.^{31,40,41} EAD/IAD was not a specific exclusion criterion in RCTs of IVT versus placebo, but because dissection is uncommon the number of randomized patients with dissection is likely to be very low and no specific subgroup analysis has been published.³¹ There is a

theoretical concern that thrombolysis may increase the risk of enlargement of an intramural hematoma in the dissected artery and thus impair cerebral hemodynamics, or promote dissecting aneurysm formation or vessel rupture.^{42,43}

Our systematic review identified no randomized data on the efficacy and safety of IVT in patients with EAD/IAD. Only 4 observational studies assessing the effect of IVT on clinical outcome in EAD/IAD patients met our inclusion criteria (Figure 1 and Table 1).^{44–47} Of note, all except one of the aforementioned studies (which was based on electronic health records with no indication on thrombolytic agent types⁴⁷) consistently refer to alteplase, with no data available for other thrombolytic agents (e.g., tenecteplase). In total, these studies gather 593 EAD/IAD patients receiving IVT and 7573 EAD/IAD patients without IVT in the acute phase of EAD/IAD-related ischemic stroke. The vast majority of these patients were reported in a single study based on electronic health records from the Nationwide Inpatient Sample (NIS) in the United States (488 and 7374 EAD/IAD patients with IVT and without IVT, respectively),⁴⁷ where the diagnosis of dissection was based on International Classification of Diseases (ICD)-9 codes, thus not allowing a distinction between EAD and IAD. The other three studies focused exclusively on EAD patients.^{44–46}

Mortality was assessed at 3 months for three studies,^{44–46} and seven days for two studies,^{45,47} of which three contributed to the meta-analysis, due to a lack of events in two studies: Overall, the meta-analysis showed no difference between the use of IVT compared to no-IVT treatment on mortality (OR, 1.95 [95% CI, 0.32–11.99]; $p = 0.47$; $I^2 = 83\%$; Figure 2(a)).^{45–48} At 7 days,^{47,49} mortality was significantly higher in EAD/IAD patients who received IVT compared to those who didn’t (OR, 4.16 [95% CI, 3.06–5.65]; $p \leq 0.001$; $I^2 = 0\%$; Figure 2(b)). This result was largely driven by the NIS based on electronic health records.⁴⁷ In this study, patients with arterial dissections receiving IVT had a higher rate of medical co-morbidities, adjunctive procedures, and medical complications, compared to patients with dissection not receiving IVT. This study also included patients with ischemic stroke unrelated to arterial dissection, with ($N = 47,411$) and without IVT ($N = 2,964,253$) and, among all patients with ischemic stroke, there was no significant interaction between dissection and thrombolytic treatment for predicting in-hospital mortality ($p = 0.78$).

Regarding functional outcome (105 and 199 EAD patients with and without IVT),^{44–46} there were no differences between IVT and no-IVT patients, both using the endpoint excellent functional outcome (mRS 0–1 vs. 2–6; OR, 0.95 [95% CI, 0.46–1.96]; $p = 0.90$; $I^2 = 0\%$; Figure 3(a))^{44,45} and good functional outcome (mRS 0–2 vs. 3–6; OR, 1.19 [95% CI, 0.70–2.01]; $p = 0.52$; $I^2 = 0\%$; Figure 3(b))^{44,46,48} at 3 months. NIHSS at admission was higher in EAD patients with than those without IVT (Table 1).^{44,46}

Figure 1. PRISMA flow chart of study selection for PICO1. “Wrong study design, patient population, intervention, comparator, and control group” in COVidence corresponds to studies that do not match criteria for this PICO.

Finally, IVT with alteplase was found to increase the odds of intracranial hemorrhage (OR, 5.35 [95% CI, 6.62–7.92], $p < 0.00001$, $I^2 = 0\%$, [Figure 4\(a\)](#)), which became non-significant after removing the NIS study that likely included a mix of EAD and IAD cases and was based on electronic health records only (OR, 7.64 [95% CI, 0.91–63.86]; $p = 0.08$; $I^2 = 0\%$; [Figure 4\(b\)](#)).^{44–46,48} There was no significant association with major bleedings (OR, 3.92 [95% CI, 0.60–25.66]; $p = 0.15$; $I^2 = 0\%$; [Figure 5](#)).^{44,46}

The results of these small observational studies should be viewed with extreme caution, as there are serious

inconsistencies, very serious imprecision due to small sample size, and strong suspicion of publication bias, leading overall to very low certainty ([Table 2](#) and [Supplementary Table 1.1](#)).

The high rate of cervical internal carotid artery occlusions and associated supra-clinoid internal carotid artery (tandem) occlusions seen in patients with underlying dissection could potentially, partly, account for the apparent lack of benefit with IVT.^{44,46} However, the results are very difficult to interpret in this observational setting where EAD/IAD patients undergoing IVT were more

Table 1. Summary of observational studies findings relevant for PICO1.

Study	Population	Mean age	Intervention		Comparison		Duration of follow-up	Outcome	IPD		IPD control	
			Type	NIHSS at admission	Type	NIHSS at admission			event	N	event	N
Bernardo 2019 ⁴⁶	EAD patients with acute ischemic stroke and intracranial occlusion	Intravenous thrombolysis: 53 (48–59)* No revascularization treatment; 48 (41–56)*	IVT	14 (8–18)*	No revascularization treatment	8 (2–21)*	3 months	Mortality mRS 0–2 Intracranial hemorrhage Major bleeding	3 26 2 2	38 38 38 1	5 24 0 1	44 44 43 39
Dziewas 2003 ⁴⁵	EAD patients with acute ischemic stroke	43.2 ± 11.2	IVT	NA	Anticoagulation with heparin followed by coumarin or antiplatelets	NA	Max 6 months	Mortality mRS 0–1 Intracranial hemorrhage	0 2 0	3 3 3	1 56 1	91 91 91
Engelter 2012 ⁴⁴	EAD patients with acute ischemic stroke	With thrombolysis: 45 (interquartile range 36–50)*; Without thrombolysis : 44.5 (38.5–51)* ^a	IVT	16 (interquartile range 10–19)*	No thrombolysis	14 (interquartile range 7–18.5)*	3 months	Mortality mRS 0–1 mRS0–2 Intracranial hemorrhage Major bleeding	0 19 35 4 4	64 64 64 64 64	0 20 35 0 0	64 64 64 64 64
Qureshi 2011 ⁴⁷	Patients with ischemic stroke and arterial dissection (ICD–9 codes 443.2, 443.21, 443.24 ^b ; EAD and IAD not distinguished	Patients treated with thrombolytics: 50.13 Patients not treated with thrombolytics : 50.1	IVT	NA	No thrombolysis	NA	7.87 days for patient who received thrombolytics; 6.9 days for patients who did not receive thrombolytics	Mortality ^b Intracranial hemorrhage	55 34	488 488	214 103	7374 7374

Values are mean ± SD or mean (range) unless otherwise specified; * median (IQR).

^aInternational Classification of Diseases (ICD)9 codes 443.2: other arterial dissection (excluding aortic and coronary dissection), 443.21: dissection of carotid artery, and 443.24: dissection of vertebral artery.

^bIn hospital mortality, which was approximated as 7 day mortality for the meta-analysis, based on the mean duration of hospital stays.

Figure 2. (a) Meta-analysis of effects of intravenous thrombolysis in observational studies on mortality at 3 months. (b) Meta-analysis of effects of intravenous thrombolysis in observational studies on mortality at 7 days.

Figure 3. (a) Meta-analysis of effects of intravenous thrombolysis in observational studies on excellent functional outcome (mRS 0–1 vs. mRS 2–6) at 3–6 months. Functional outcome was assessed after a maximum of 6 months in Dziewas et al.⁴⁵ and at 3 months in Engelter et al.⁴⁴. (b) Meta-analysis of effects of intravenous thrombolysis in observational studies on good functional outcome (mRS 0–2 vs. mRS 3–6) at 3 months.

severe than EAD/IAD patients not receiving IVT, as explained above.

In this context, it is important to consider studies that compared patients with EAD and ischemic stroke treated with IVT to patients with non-EAD ischemic stroke treated with IVT. In the Safe Implementation of Thrombolysis in

Stroke International Stroke Thrombolysis Register (SITS-ISTR), compared with non-EAD ischemic stroke patients matched for age and stroke severity, EAD patients receiving thrombolytic therapies (180 patients of whom 67% received IVT and 33% intra-arterial thrombolysis) showed no significant differences in terms of safety and prognosis.⁵⁰ In

Figure 4. (a) Meta-analysis of effects of intravenous thrombolysis in observational studies on the risk of intracranial hemorrhage. (b) Meta-analysis of effects of intravenous thrombolysis in observational studies on the risk of intracranial hemorrhage, without the NIS study.⁴⁷

Figure 5. Meta-analysis of effects of intravenous thrombolysis in observational studies on the risk of major bleedings.

particular, only one patient with an intramural hematoma expansion after IVT treatment was reported, who had an excellent outcome. In the NIS study,⁴⁷ no increase in the principal safety end point of post-thrombolytic intracranial hemorrhage was observed between EAD/IAD ischemic stroke patients and ischemic stroke patients without dissection. The lack of interaction between IVT and dissection on the risk of death and the absence of excessive rates of post-thrombolytic hemorrhages suggested the use of thrombolytic treatment in EAD/IAD patients was not subject to safety concerns.⁴⁷ This is in line with results from the Swiss IVT databank,⁵¹ where 55 EAD patients undergoing IVT were compared with 1007 IVT-treated ischemic stroke patients without EAD. Intracranial hemorrhages were equally frequent in EAD (14%) and non-EAD patients (14%, $p = 0.99$), and recurrent ischemic strokes occurred in 1.8% of EAD patients and in 3.7% of non-EAD patients ($p = 0.71$). These findings are also

supported by those of a recent prospective multicenter study and a meta-analysis of data from patients with dissection-related ischemic stroke treated with IVT, which confirmed that the risk of ICH in these patients is overall quite low (2% in the meta-analysis), while an excellent functional outcome (mRS 0–1 at hospital discharge) was observed in 41% of patients.⁵²

Thus, despite the low level of evidence and the limited data available, considering the overwhelming evidence of a benefit during the acute phase of ischemic stroke in general, and in line with a recent ESO guideline on IVT,³¹ we believe that the treatment of EAD-induced acute ischemic stroke using IV alteplase within 4.5 hours of onset is safe.

Of note, in the previously published recent ESO IVT guideline,³¹ the authors decided not to perform a meta-analysis of observational studies comparing EAD patients with IVT versus no IVT due to notable differences in study design, important risk of selection bias and confounding by

Table 2. Quality of results in observational studies relevant for PICO1.

Certainty assessment			No. of patients		Effect		Certainty	Importance				
No. of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations			PICO 1: Thrombolytics	Control	Relative (95% CI)	Absolute (95% CI)
PICO 1.1 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a reduced risk of death at 3 months?												
3	observational studies	not serious	serious ^a	not serious	very serious ^b	publication bias strongly suspected ^c	58/590 (9.8%)	219/7482 (2.9%)	OR 1.95 (0.32 to 11.99)	26 more per 1,000 (from 20 fewer to 236 more)	⊕○○○ VERY LOW	CRITICAL
PICO 1.1 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a reduced risk of death at 7 days?												
2	observational studies	not serious	not serious	not serious	not serious	publication bias strongly suspected strong association ^{c,d}	58/526 (11.0%)	216/7421 (2.9%)	OR 4.16 (3.06 to 5.65)	82 more per 1,000 (from 55 more to 116 more)	⊕○○○ VERY LOW	CRITICAL
PICO 1.2 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a higher likelihood of good functional outcome (mRS 0–2 vs. 3–6)?												
2 ^g	observational studies	not serious	not serious	not serious	serious ^e	publication bias strongly suspected ^c	61/102 (59.8%)	59/108 (54.6%)	OR 1.25 (0.71 to 2.19)	55 more per 1,000 (from 85 fewer to 179 more)	⊕○○○ VERY LOW	CRITICAL
PICO 1.2 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a higher likelihood of excellent functional outcome (mRS 0–1 vs. 2–6)?												
2	observational studies	not serious	not serious	not serious	serious ^e	publication bias strongly suspected ^c	21/67 (31.3%)	76/155 (49.0%)	OR 0.95 (0.46 to 1.96)	13 fewer per 1,000 (from 184 fewer to 163 more)	⊕○○○ VERY LOW	CRITICAL

(continued)

Table 2. (continued)

Certainty assessment			No. of patients		Effect		Certainty	Importance					
No. of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations			PICO 1: Thrombolytics	Control	Relative (95% CI)	Absolute (95% CI)	
PICO1.3 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a reduced risk of intracranial hemorrhage?													
3	observational studies	not serious	not serious	not serious	not serious	publication bias strongly suspected	very strong association ^{c,f}	40/590 (6.8%)	103/7481 (1.4%)	OR 5.35 (3.62 to 7.92)	56 more per 1,000 (from 34 more to 86 more)	⊕⊕⊕○ MODERATE	CRITICAL
PICO1.4 Is intravenous thrombolysis versus no intravenous thrombolysis associated with a reduced risk of major bleeding?													
2 ^g	observational studies	not serious	not serious	not serious	very serious ^b	publication bias strongly suspected ^c		6/102 (5.9%)	1/103 (1.0%)	OR 3.92 (0.60 to 25.66)	27 more per 1,000 (from 4 fewer to 191 more)	⊕⊕○○ LOW	CRITICAL

CI: Confidence interval; OR: Odds ratio; MD: Mean difference

^aSignificant heterogeneity, I² = 83%.

^bVery wide confidence intervals.

^cThree or less studies reported this outcome.

^dLarge effect.

^eWide confidence intervals.

^fVery large effect.

^gDziewas 2003 was not meta-analyzed due to zero events and only 3 patients in the intervention group.

indication. As study selection is specific to each ESO guideline, slightly different study inclusion criteria were used in the IVT and in the present dissection guideline. In the absence of RCTs for EAD/IAD except for PICO5, our MWG had indeed decided to analyze and critically interpret all available evidence, as described above. Importantly, our conclusion and evidence-based recommendation regarding use of IVT in EAD patients is the same as in the ESO IVT guideline.³¹

Additional information

Even more than for EAD, the management of IAD is controversial because RCTs and large observational studies are not available.¹³ Only anecdotal reports (case reports or small case series of fewer than 10 patients) have been published on the safety and efficacy of IVT in patients with IAD presenting with acute ischemic stroke.^{49,53,54} The scarce information available, therefore, does not allow conduction of a meta-analysis. In a consecutive single-center series of 181 EAD/IAD ischemic stroke patients,⁴⁹ 10 were due to IAD and five of these received IVT. Among these, there were no cases of SAH nor symptomatic ICH after IVT. One patient had an asymptomatic hemorrhagic infarct type 1 and two patients died within 7 days from ischemic mass effect. The other three patients had favorable clinical outcomes at 3 months.⁴⁹

By contrast with extracranial arteries, intradural arteries are characterized by a well-developed internal elastic lamina, a paucity of elastic fibers in the media, little adventitial tissue, and no external elastic lamina.¹¹ These features, and weaker supporting tissues than cervical arteries,¹² may make intracranial arteries increasingly more prone to bleeding than extracranial arteries. Based on these observations, there is an obvious theoretical concern that IAD may result in sub-adventitial extension of the hematoma and increase the risk of SAH (especially when located in the posterior circulation) or that brain ischemia might progress to a hemorrhagic transformation as a consequence of thrombolysis, which has made many clinicians reluctant to use IVT in known or suspected IAD.⁵⁵ It should be noted, however, that the pathognomonic radiological findings of IAD (mural hematoma, intimal flap, and double lumen) are not easily detectable, especially in the hyperacute phase of the disease. In most cases, as described in the introduction, the definite diagnosis of IAD needs the combination of arterial wall and lumen imaging and also the comparison between baseline and follow-up imaging. This implies that, in clinical practice, the scenario in which the neurologist must decide whether or not to administer IVT in a patient with acute ischemic stroke because of a definite diagnosis of IAD is extremely uncommon.

The recent ESO guideline on IVT³¹ suggested to refrain from IVT in IAD patients with acute ischemic stroke (approved by 6 of 9 group members). We believe that caution is warranted in patients with a suspected diagnosis of IAD. Indeed, IAD diagnosis is only very rarely confirmed in the time window for IVT in acute ischemic stroke patients. There is no evidence in the current literature that IVT in IAD patients without SAH is harmful and we fear that systematically refraining from IVT in patients with a suspicion of possible IAD may lessen the chances of ischemic stroke patients of benefiting from an efficient therapy. Therefore, we suggest that in patients with an acute ischemic stroke suspected to be caused by IAD, IVT should probably not be withheld after ruling out subtle signs of SAH on initial brain imaging. Our expert consensus statement is not in contradiction with the expert consensus statement of the IVT guideline that focused on the rare situation of patients with a confirmed diagnosis of IAD with 4.5 hours of ischemic stroke onset.

Evidence-based recommendation

In patients with symptomatic EAD with acute ischemic stroke within 4.5 hours of onset, we suggest using IVT with alteplase, if the standard inclusion / exclusion criteria are met.

Quality of evidence: Low ⊕ ⊕

Strength of recommendation: Weak for an intervention ↑?

In patients with symptomatic IAD with acute ischemic stroke within 4.5 hours of onset, there is insufficient data to provide a recommendation.

Quality of evidence: Very low ⊕

Strength of recommendation: -

Expert consensus statement

In patients with an acute ischemic stroke suspected to be caused by IAD, all but one expert suggest that IVT should be considered, after ruling out standard contra-indications, including subtle signs of sub-arachnoid bleeding on brain imaging.

PICO 2: In extracranial artery dissection (EAD) & intracranial artery dissection (IAD) patients with acute ischemic stroke is endovascular treatment (stenting and/or thrombectomy) versus no endovascular treatment (with or without IV thrombolysis) associated with a reduced risk of death, a higher likelihood of favorable functional outcome (mRS 0–2 vs 3–6, or 0–1 vs 2–6, or equivalent), and no increased risk of ICH, or SAH?

Analysis of current evidence

No randomized controlled trial was available to address this PICO question. After exclusion of single-arm studies and systematic reviews, we identified five comparative observational studies including 463 EAD patients that met our inclusion criteria (190 patients with EVT and 373 with no EVT or [in 71 patients] without the primary EVT of interest [in that case carotid stenting]; [Figure 6](#) and [Table 3](#)).^{46,48,56-58} No study on this question in IAD was identified.

Of note, two steps in EVT need to be considered: intracranial clot removal (i.e., mechanical thrombectomy [MT]) and management of the EAD lesion and related stenosis. Most of the identified studies focused predominantly on the first step. Different types of EVT were used in the five identified observational studies. One study (48/48 EAD patients with/without intervention) focused on EVT with MT only compared with medical treatment and did not describe if adjunct procedures were used.⁵⁶ One study included only EAD patients undergoing EVT (the strategy of which was left to the discretion of the interventionist) and

Figure 6. PRISMA flow chart of study selection for PICO2. “Wrong study design, setting, intervention, outcomes, and (patient) population” in COVidence corresponds to studies that do not match criteria for this PICO

Table 3. Summary of findings from observational studies relevant for PICO2.

Study	Population	Mean age	Intervention		Comparison		Follow-up	Outcome	IPD			IPD control		
			Type	NIHSS at admission	Type	NIHSS at admission			event	N	total	event	N	total
Bernardo, 2019 ⁴⁶	EAD with acute ischemic stroke and intracranial occlusion	50 (42–57)*	Endovascular treatment (different types ^a)	15 (12–19)*	IVT alone or no revascularization treatment	IVT: 14 (8–18)* No revascularization: 8 (2–21)*	3 months	mRS 0–2 Symptomatic ICH Mortality	9 1 1	21 24 21	50 2 8	82 81 82		
Jensen, 2017 ⁴⁸	EAD with acute ischemic stroke	Intra-arterial thrombolysis: 52 (43–60)*; Intravenous thrombolysis: 52 (37–59)*; No anti-arterial thrombolysis: 44 (35–54)*	Intra-arterial thrombolysis (with or without intravenous thrombolysis) ^a	13 (12–16)*	Intravenous thrombolysis alone or no anti-arterial thrombolysis ^b	Intravenous thrombolysis: 10 (7–12)*; No anti-arterial thrombolysis: 3 (1–7)*	90 days	mRS 0–2 Mortality	2 3	20 15	93 5	137 137		
Li 2018 ⁵⁶	EAD with acute ischemic stroke and with large artery occlusion	With thrombectomy : 48.8 ± 12.6; without thrombectomy : 49.4 ± 8.7	Endovascular thrombectomy	14.0 (12.0–18.0)*	Without thrombectomy (medical treatment alone, 15% IVT)	14.5 (10.3–17.8)*	90 days	mRS 0–2 Symptomatic ICH Mortality	32 4 5	48 48 48	19 2 3	48 48 48		
Marnat 2020 ⁵⁷	EAD (internal carotid artery) with tandem occlusion (TITAN & ETIS)	51.9 ± 11.7	Endovascular treatment (emergency carotid artery stenting)	15 (10–19)*	No carotid artery stenting	17 (13–20)*	90 days	mRS 0–2 Symptomatic ICH Mortality	35 7 5	65 65 65	44 4 4	71 71 71		
Traenka 2018 ⁵⁸	EAD with acute ischemic stroke	48.8 (44–55.2) *	Endovascular treatment (different types ^c)	16 (13.5–18)*	Intravenous thrombolysis	14.5 (8.0–17.8)*	3 months	mRS 0–2 Symptomatic ICH Mortality	15 5 6	38 38 38	13 0 0	24 24 24		

* Several endovascular techniques were used, including intra-arterial thrombolysis (n = 1, 4.2%), thrombectomy with stent-retriever (n = 17, 70.8%), thromboaspiration (n = 8, 33.3%), angioplasty (n = 7, 29.2%), and cervical stent placement (n = 11, 45.8%). Most patients (n = 13, 54.2%) had more than one technique for EVT procedure.

^aSeveral endovascular techniques were used: the majority of patients had multiple IAT techniques, including thrombectomy (n = 19), IA thrombolysis (n = 17), stenting (n = 14), and angioplasty (n = 7); 11 patients were also treated with IVT.

^b: The comparison group here comprised patients with EAD not receiving intra-arterial thrombolysis (n = 137, of whom 11 received IVT).

^cSeveral endovascular techniques were used, including mechanical thrombectomy with devices available 2007–2012 (stent retrievers), thrombus aspiration, or percutaneous transluminal angioplasty; intra-arterial application of urokinase or tissue plasminogen activator (tPA) as well as intracranial or extracranial stent placement was performed in selected patients.

compared EAD patients in whom the EVT included carotid artery stenting to EAD patients undergoing EVT without carotid stenting (65/71).⁵⁷ Finally, three studies used mixed endovascular approaches,^{46,48,58} with mostly more than one technique per procedure (comparing in total 77/254 patients with/without EVT). A majority of patients in these studies underwent MT with stent retrievers, other interventions included intra-arterial thrombolysis, thrombus aspiration, percutaneous transluminal angioplasty, and cervical stent placement (45.8% of patients in one of the studies⁴⁶). In these studies, it was difficult to identify patients who only had one technique or a combination of these.

The meta-analysis conducted on these five EAD studies^{46,48,56-58} showed similar odds ratio (OR) for good functional outcome (mRS 0–2 or good recovery on Glasgow Outcome Scale), OR, 0.56 [95% CI, 0.19–1.59], $I^2 = 84\%$, $p = 0.27$, Figure 7) after EVT compared to no EVT. This finding is in accordance with a previous meta-analysis,⁵⁸ which included 8 studies and 212 patients (110 IVT and 102 EVT [including smaller studies than allowed in our inclusion criteria (Methods)]), showing no significant difference between the two therapeutic approaches (OR for good functional outcome, mRS 0–2, 0.97 [95% CI, 0.38–2.44]), despite the fact that EVT was associated with higher rates of recanalization (respectively, 84.2% for EVT vs. 66.7% for IVT; OR, 3.2 [95% CI, 0.9–11.38]) in that study.⁵⁸ Of note, the difference in NIHSS between patients

treated with EVT and those who were not, in three of the five studies included in our meta-analysis,^{46,48,58} shows that EVT was performed in patients with more severe strokes and represents, therefore, an important selection bias (Table 3). The study by Li et al. that applied propensity score matching of 48 EAD patients undergoing EVT with 48 EAD patients without EVT, enabling a similar severity profile in both groups (Table 3), is the only study that showed a superiority of EVT, with 66.7% versus 39.6% of patients reaching a good functional outcome at 90 days ($p = 0.008$).⁵⁶

Of note, in the study by Marnat et al.,⁵⁷ on the databases of Endovascular Treatment in Ischemic Stroke (ETIS) and Thrombectomy in Tandem Lesion (TITAN),^{59,60} including 65 EAD patients treated with emergency carotid stenting and 71 with EVT without stenting, EVT with EAD stenting was associated with greater rates of successful reperfusion (adjusted OR [aOR], 2.24 [95% CI, 1.33–2.77]). However, in sensitivity analyses restricted to patients with successful reperfusion, EVT with EAD stenting was not associated with improved rates of favorable outcome (aOR, 0.70 [95% CI, 0.45–1.08]), as in the overall group (Figure 7).

With respect to safety, EVT in EAD was associated with a non-significant increased risk of all symptomatic ICH (OR, 2.27 [95% CI, 0.92–5.61], $I^2 = 0\%$, $p = 0.08$, Figure 8). Previous lines of evidence showed a trend towards more ICH in the EVT group (20.5% vs. 5.3%, $p = 0.072$), but not

Figure 7. Meta-analysis of effects of endovascular treatment in observational studies on good functional outcome (mRS 0–2 vs. 3–6) at 3 months.

Figure 8. Meta-analysis of effects of endovascular treatment in observational studies on risk of symptomatic ICH at 3 months.

symptomatic ICH (sICH, 4.2% vs. 5.3%, $p = 0.669$).⁴⁶ Of note, Traenka et al. and Bernardo et al. reported sICH nearly solely among EVT patients who had bridging therapy with IVT (80–100% of the cases).^{46,58} Mortality did not differ significantly between EVT and non EVT-treated patients overall (OR, 2.15 [95% CI, 0.86–5.37], $I^2 = 28\%$, $p = 0.10$, Figure 9), and at 7 days (OR, 2.44 [95% CI, 0.25–23.89], $I^2 = 66\%$, $p = 0.44$), Figure 9).

It is important to acknowledge that most of the studies are retrospective, conducted at a single center, and lack matched control groups. In addition, EVT strategies, stenting/angioplasty protocols, and antiplatelet regimens were not standardized. These facts underline the significant risk of bias, to be considered when interpreting the data (Table 4 and Supplementary Table 1.2 for risk bias assessment).

EVT with MT plus best medical management over best medical management alone is recommended for clot extraction to improve functional outcome in adults with anterior circulation large vessel occlusion-related acute ischemic stroke,⁶¹ irrespectively of the presence of a dissection. Previous ESO/ESMINT guidelines on MT have not addressed the specific situation of EAD/IAD patients.⁶¹ Of note, based on a large single-center dataset of 445 consecutive patients receiving EVT (i.e., MT, IA thrombolysis, stent, and angioplasty), Jensen and collaborators compared the outcome between 24 EAD patients and 421 patients with another cause of acute ischemic stroke.⁴⁸ There was no significant difference in the odds of a good functional outcome (mRS 0–2) at 90 days between the two groups, accounting for differences in age and stroke severity.⁴⁸ Similarly, in the prospective, multicenter Prognostic

Factors Related to Clinical Outcome Following Thrombectomy in Ischemic Stroke (RECAST) study, the rate of good functional outcome (mRS 0–2) was similar (as were recanalization rates) in 20 carotid EAD patients with tandem occlusion undergoing MT ($\pm$ IVT) compared to 201 non-EAD patients with isolated intracranial large vessel occlusion.⁶² In a more recent study, including the analysis of a database of 1422 acute ischemic stroke patients, with 43 EAD patients matched to 86 patients with other etiologies, safety and efficacy of EVT with MT were comparable in both groups: sICH and mortality rates were similar (OR, 0.85 [95% CI 0.21–3.49], $p = 0.82$; OR, 1.54 [95% CI 0.33–2.79], $p = 0.58$; OR, 0.18 [95% CI 0.02–1.46], $p = 0.11$, respectively), as well as rates of favorable functional outcome (OR, 1.26 [95% CI 0.61–2.64], $p = 0.53$).⁶³

Thus, despite the dearth and heterogeneity of (purely observational) data assessing EVT in EAD patients, considering the overwhelming benefit of MT in the setting of acute ischemic stroke with large vessel occlusion of the anterior circulation in general, we believe that the treatment of EAD-induced acute ischemic stroke with anterior circulation large vessel occlusion using MT is efficient and safe.

Additional information

The comparison between primary MT and bridging strategies of IVT immediately followed by MT in acute ischemic stroke patients with EAD is beyond the scope of this guideline but deserves special attention in future research and careful multidisciplinary assessment in current practice. As mentioned above, the few cases of sICH after MT were

Figure 9. (a) Meta-analysis of effects of endovascular treatment in observational studies on mortality, at 3 months. (b) Meta-analysis of effects of endovascular treatment in observational studies on mortality at 7 days.

Table 4. Quality of results in observational studies relevant for PICO2.

Certainty assessment			No. of patients		Effect		Certainty	Importance			
No. of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations			Endovascular	Control	Relative (95% CI)
PICO 2.2.1 Is endovascular treatment (stenting and/or thrombectomy) (I) versus no endovascular treatment (with or without IV thrombolysis) (C), associated with a reduced risk of death at 3 months?											
5	observational studies	not serious	not serious	not serious	not serious	publication bias strongly suspected ^a	20/187 (10.7%)	20/362 (5.5%)	OR 2.15 (0.86 to 5.37)	56 more per 1,000 (from 7 fewer to 184 more)	⊕○○○ VERY LOW CRITICAL
PICO 2.2.1 Is endovascular treatment (stenting and/or thrombectomy) (I) versus no endovascular treatment (with or without IV thrombolysis) (C), associated with a reduced risk of death at 7 days (subgroup)?											
2	observational studies	not serious	not serious	not serious	very serious ^b	publication bias strongly suspected ^a	4/39 (10.3%)	10/222 (4.5%)	OR 2.44 (0.25 to 23.84)	58 more per 1,000 (from 33 fewer to 484 more)	⊕○○○ VERY LOW CRITICAL
PICO 2.2.2 Is endovascular treatment (stenting and/or thrombectomy) (I) versus no endovascular treatment (with or without IV thrombolysis) (C), associated with a higher likelihood of good functional outcome (mRS 0–2 vs. 3–6 or equivalent) at 3 months?											
5	observational studies	not serious	serious ^c	not serious	serious	publication bias strongly suspected ^a	93/192 (48.4%)	219/362 (60.5%)	OR 0.56 (0.19 to 1.59)	143 fewer per 1,000 (from 380 fewer to 104 more)	⊕○○○ VERY LOW CRITICAL
PICO 2.2.3 Is endovascular treatment (stenting and/or thrombectomy) (I) versus no endovascular treatment (with or without IV thrombolysis) (C), associated with increased risk of symptomatic ICH at 3 months?											
4	observational studies	not serious	not serious	not serious	not serious	publication bias strongly suspected ^a	17/175 (9.7%)	8/224 (3.6%)	OR 2.27 (0.92 to 5.61)	42 more per 1,000 (from 3 fewer to 136 more)	⊕○○○ VERY LOW CRITICAL

CI: Confidence interval; OR: Odds ratio; MD: Mean difference

^aSix or less studies reported this outcome.^bVery wide confidence intervals.^cSignificant heterogeneity, $I^2 > 65\%$.^dWide confidence intervals.^eVery large effect.

reported mostly among EVT patients who had bridging therapy with IVT and some authors have listed dissections as a factor potentially favoring primary MT.⁶⁴

Whether the EAD requires a specific EVT (e.g., angioplasty and stenting) before or after intracranial large vessel clot extraction remains to be determined. Treating the EAD lesion during the same procedure may be required in case of hemodynamic compromise (e.g., carotid occlusion with incomplete circle of Willis) or recurrent embolism during EVT.⁶⁵ Whenever possible, these complex cases should be discussed between the interventionalist and the neurologist, as the implantation of a stent requires a specific antithrombotic regimen (to prevent stent thrombosis that could cause ischemic stroke), with an increased bleeding risk in the setting of acute ischemic stroke.

For patients with IAD, beyond the bleeding risk inherent to the antiplatelet therapy needed by the stenting, the benefit-risk ratio of EVT (i.e., intracranial stenting) needs to be discussed also on the basis of the IAD location, considering anatomic specificities (e.g., occlusion risk of perforating arteries of the M1 segment of the middle cerebral artery).

Evidence-based Recommendation

In acute ischemic stroke patients with EAD and large vessel occlusion of the anterior circulation, we suggest using MT.

Quality of evidence: Very low ⊕

Strength of recommendation: Weak for an intervention ↑?

In acute ischemic stroke patients with IAD, there is insufficient data to provide a recommendation regarding the use of EVT.

Quality of evidence:

Very low ⊕

Strength of recommendation: -

Expert consensus statement

In acute ischemic stroke with EAD and large vessel occlusion of the anterior circulation, all but one expert suggest EVT (other than MT) for the treatment of the EAD lesion in case of carotid occlusion without patent circle of Willis or in case of recurrent embolism.

In acute ischemic stroke with IAD and large vessel occlusion of the anterior circulation, all experts suggest EVT (other than MT) for the treatment of the IAD lesion at the hyperacute phase after assessing the risk/benefit ratio based on the location of the dissection and bleeding risk.

PICO 3: In patients with an intracranial dissecting aneurysm and a subarachnoid hemorrhage (SAH) does endovascular or surgical treatment of the aneurysm versus optimal medical treatment alone reduce the risk of SAH recurrence, ICH, death and increase the likelihood of favorable functional outcome (mRS 0–2 vs. 3–6, or 0–1 vs. 2–6, or equivalent)?

Analysis of current evidence

No randomized control trial comparing intervention (endovascular or surgical) versus optimal medical treatment was identified. Only observational studies, mostly single-arm studies, were found. A single study reported on the natural history of the disease.⁶⁶ Four studies, all retrospective, comparing intervention with medical treatment, were identified (Table 5, Figure 10 and Supplementary Table 1.3 for risk bias assessment) and in all studies allocation to medical treatment was justified either by a poor initial condition of the patient or by the risk and complexity of the intervention making it futile.^{67–70} The overall risk of bias was rated as serious, inconsistency not serious, publication bias was strongly suspected, and the observation certainty was rated as very low and the importance as critical (Table 6). Intervention was associated with a lower risk of SAH recurrence (OR, 0.06 [95% CI, 0.02–0.22], $I^2 = 44\%$, $p < 0.0001$, Figure 11) and with a lower risk of death (OR, 0.15 [95% CI, 0.04–0.56], $I^2 = 40\%$, $p = 0.005$, Figure 12). Good functional outcomes with mRS 0–2 tended to be more frequent in patients treated by intervention compared to patients under medical treatment (OR, 2.32 [95% CI, 0.95–5.68], $I^2 = 0\%$, $p = 0.07$, Figure 13). The mean follow-up time was 28.8 months.

Additional information

Due to the small number of comparative observational studies,^{67–70} we chose to also report the mean frequency of PICO3 outcomes across all identified observational studies (comparative and single-arm), by treatment group (endovascular/surgical or medical). In addition to the four comparative observational studies, twenty-three single-arm studies providing data regarding mortality, functional outcome, and SAH recurrence (rebleeding) rates were identified and included for descriptive analyses (one with medical treatment only⁶⁶ and all others with endovascular or surgical treatment only).^{66–92} In one study, medical and interventional management of cases was merged and both groups could not be unequivocally analyzed separately, this study was therefore only included in the analysis of the overall rebleeding rate.⁹³ Most studies were from Japan (11/27) followed by Korea (4/27), China (4/27), France (1/27), and

Table 5. Summary of observational studies findings relevant for PICO3.

Study	Population	Mean age	Intervention		Comparison		Follow-up	Outcome	IPD intervention			IPD comparison			
			Type	Severity	Type	Severity			N	event	N	total	N	event	total
Anxionnat 2003 ⁶⁷	SAH related to IAD	48 (16–74)	Endovascular treatment (occlusion using coil; proximal balloon occlusion); wrapping	GOS 5: 1; GOS 4: 5; GOS 1: 1 Hunt and Hess: grade IV: 1; grade III: 3; grade II: 8; grade I: 5	Conservative treatment	GOS 5: 6; GOS 4: 1; GOS 1: 3 Hunt and Hess: grade IV: 1; grade III: 1; grade II: 1; grade I: 3	1 year (N = 14)	Mortality Recurrent SAH mRS 0–2	1	17	3	10			
Mizutani 1995 ⁶⁹	SAH related to IAD	53.4 (32–77)	Surgical: proximal vertebral artery obliteration; trapping; wrapping; bleb clipping; bleb clipping combined with wrapping	NA ^a	No surgery	NA ^a	1 week	Mortality Recurrent SAH mRS 0–2	4	29	11	13			
Rabinov 2003 ⁶⁸	IAD (intradural vertebralbasilar dissecting aneurysms)	52 (17–87)	Endovascular treatment (trapping or proximal occlusion) or surgery	Hunt and Hess grade 0: 4; grade I: 7, grade III: 1; grade IV: 4; grade V: 1	Untreated	Hunt and Hess: grade II: 1; grade III: 3; grade IV > V: 1; grade V: 1	3.5 years	Mortality Recurrent SAH mRS 0–2	5	22	2	6			
Zhao 2007 ⁷⁰	SAH related to IAD (intracranial vertebralbasilar dissection)	42.9 (6–67)	Endovascular treatment (proximal occlusion; parent artery embolization; endovascular trapping); surgery	Glasgow coma scale: GCS 15: 6; GCS 14: 3; GCS 10: 2; GCS 7: 1; GCS 6: 1; GCS 5: 1 Hunt and Hess: Grade IV: 2; Grade III: 1; Grade II: 6; Grade I: 2	Conservative treatment	Glasgow coma scale: GCS 15: 4; GCS 12: 1; GCS 6: 1; GCS 5: 1 Hunt and Hess: Grade IV: 2; Grade III: 1; Grade II: 3; Grade I: 1	Up to 7.5 years	Mortality mRS 0–2	2	14	3	7			

Values are mean (range) unless otherwise specified; GOS: Glasgow outcome scale.

^aOf the 42 patients, 31 (73.8%) remained in good or fair neurological condition after the initial attack.

Figure 10. PRISMA flow chart of study selection for PICO3.

USA (2/27). As there was important heterogeneity in the frequency of reported outcomes across studies, a random effects model was used to compute the mean frequencies.

The natural history and outcome of IAD-related SAH is similar to aneurysmal SAH. In identified studies, IAD with SAH was associated with a high mortality under medical treatment only, estimated on average at 55% (95% CI [34%–76%]) (Figure 14).^{66-70, 94,95} The strongest predictors of mortality are clinical condition on admission and rebleeding.⁹⁶ The rebleeding rate (SAH recurrence) was high (46%, 95% CI [18%–75%]) in the aforementioned studies (Figure 15).⁶⁶⁻⁶⁹ Most rebleedings occurred within the first hours to days after the initial event. In the unique study

focusing on the outcome of patients managed conservatively by Yamada et al.,⁶⁶ most rebleeding occurred within the first 6 hours (10 out of 14) and the vast majority within the first 24 hours (13 out of 14). Eleven of the 14 patients who suffered rebleeding died.⁶⁶ In the study reported by Mizutani et al., 10 out of 13 patients treated conservatively suffered rebleeding and 6 died subsequently. Nineteen of 29 patients suffered rebleeding prior to intervention. Of the 30 patients with rebleeding, 17 occurred within the first 24 hours and 24 during the first week. Rebleeding was lethal in 47% of cases.⁶⁹

Management of patients with interventions reduced the rate of rebleeding to 32% (95% CI, 18%–48%)

Table 6. Quality of results in observational studies relevant for PICO3.

Certainty assessment		No. of patients					Effect					
No. of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Endovascular/ surgical treatment	Medical	Relative (95% CI)	Absolute (95% CI)	Certainty	Importance
PICO 3.1 Does endovascular or surgical treatment of the aneurysm versus optimal medical treatment alone reduce the risk of death?												
4	observational studies	serious ^a	not serious	not serious	not serious	publication bias strongly suspected ^b	12/82 (14.6%)	19/36 (52.8%)	OR 0.15 (0.04 to 0.56)	384 fewer per 1,000 (from 485 fewer to 143 fewer)	⊕○○○ VERY LOW	CRITICAL
PICO 3.2 Is endovascular or surgical treatment of the aneurysm versus optimal medical treatment alone associated with a higher likelihood of good functional outcome (mRS 0–2 or equivalent)?												
4	observational studies	serious ^a	not serious	not serious	not serious	publication bias strongly suspected ^b	45/82 (54.9%)	13/35 (37.1%)	OR 2.32 (0.95 to 5.68)	207 more per 1,000 (from 12 fewer to 399 more)	⊕○○○ VERY LOW	CRITICAL
PICO 3.3 Does endovascular or surgical treatment of the aneurysm versus optimal medical treatment alone reduce the risk of major bleeding?												
2	observational studies	serious ^a	not serious	not serious	not serious	publication bias strongly suspected ^b	21/46 (45.7%)	0/0	not pooled	see comment	⊕○○○ VERY LOW	CRITICAL

Figure 11. Meta-analysis of effects of endovascular or surgical treatment in observational studies on the risk of rebleeding (SAH recurrence).

Figure 12. Meta-analysis of effects of endovascular or surgical treatment in observational studies on mortality. Anxionnat 2003⁶⁷: deaths occurred in the acute or subacute phase; Mizutani 1995⁶⁹: deaths occurred in the acute or subacute phase, the latest death occurred within a month; Rabinov 2003⁶⁸: four of five deaths occurred in the initial hospital course, the last death occurred in a delayed fashion due to an unknown cause; Zhao 2007⁷⁰: most deaths occurred in the acute or subacute phase, except one death that occurred at 16 months due to the rupture of another IAD.

Figure 13. Meta-analysis of effects of endovascular or surgical treatment in observational studies on good functional outcome (mRS 0–2 vs. 3–6 or equivalent).

(Figure 16).^{67,69,71,72,74,77,81,84,87,91-93} Despite patients being allocated to interventions, 31% (95% CI, 16%–47%) of cases rebled prior to intervention (Figure 17),^{67,69,71,72,74,76,77,81,84,87,91,92} and only very few rebled after the lesion was secured (2%, 95% CI, 1%–4%) (Figure 18).^{67-69,71-75,77-79,81-92}

Twenty-six studies reported mortality rates in patients with SAH secondary to IAD.^{66-75,77-92} One study focused only on patients managed medically,⁶⁶ 4 studies reported mortality for both patients managed medically or with

interventions,⁶⁷⁻⁷⁰ while the remaining studies described survival rates only in patients undergoing intervention. The mortality of IAD with SAH, when treated by intervention, was 10% (95% CI, 7%–12%) (Figure 19) as compared to 55% (95% CI, 34%–76%) in absence of intervention (Figure 14).

Twenty-five studies reported functional outcome associated with intervention,^{67-88,90-92} one study focusing only on cases managed medically⁶⁶ and 4 studies reporting outcomes for both cases managed medically and with

Figure 14. Mortality rate in IAD patients with SAH under medical treatment only.

Figure 15. Rebleeding rate (SAH recurrence) in IAD patients with SAH under medical treatment only.

Figure 16. Rebleeding rate (SAH recurrence) in IAD patients with SAH undergoing endovascular or surgical treatment.

Figure 17. Rebleeding rate (SAH recurrence) in IAD patients with SAH undergoing endovascular or surgical treatment, prior to intervention.

Figure 18. Rebleeding rate (SAH recurrence) in IAD patients with SAH undergoing endovascular or surgical treatment, after intervention.

Figure 19. Mortality rate in IAD patients with SAH undergoing endovascular or surgical treatment.

intervention.⁶⁷⁻⁷⁰ The rate of good functional outcomes defined as mRS 0–2 was observed in 70% (95% CI, 64%–76%) of patients managed with an intervention (Figure 20) and 35% (95% CI, 21–51%) of patients managed medically (Figure 21).

Overall, the very high rate of early rebleeding and subsequent fatality in the absence of intervention and the reduction of rebleeding and lower mortality associated with intervention (although based on limited small studies) are strong reasons for recommending early surgical or endovascular treatment in patients suffering IAD with SAH.

There are no RCTs and no observational studies systematically comparing different types of endovascular or surgical interventions. Those aiming at mechanically securing IAD consist most often in the trapping of the diseased vessel segment,^{67,84,96} with or without downstream revascularisation (e.g., selective aneurysmal sac occlusion through surgical clipping, surgical bypass, endovascular

[stent-assisted] coiling or stent placement, including flow-diverter stents).^{83,89,90} Another possible intervention is occlusion of the vessel proximally to the disease segment. Vessel wrapping or side wall clipping offer only a limited protection against rebleeding and should be avoided.^{69,92,97} Interdisciplinary collaboration is essential to offer the best possible treatment adapted to each situation.

Evidence-based Recommendation

In patients suffering IAD with SAH, we suggest early surgical or endovascular intervention. There is insufficient data to provide a recommendation on the type of intervention to prioritize and the precise time window.

Quality of evidence: Very low ⊕

Strength of recommendation: Weak for an intervention ↑?

Figure 20. Rate of good functional outcome in IAD patients with SAH undergoing endovascular or surgical treatment.

Figure 21. Rate of good functional outcome in IAD patients with SAH under medical treatment only.

Expert consensus statement

Different types of surgical and endovascular treatment methods can be used for treating IAD with SAH. In the absence of RCTs and considering the limited data from observational studies with high risk of bias, all experts suggest that the choice of intervention type in acute IAD-related SAH should ideally be the result of a multidisciplinary assessment.

PICO 4: In patients with an intracranial dissecting aneurysm and isolated headache (no transient ischemic attack [TIA], acute ischemic stroke, or subarachnoid hemorrhage [SAH]), does endovascular or surgical treatment of the aneurysm versus optimal medical treatment alone reduce the risk of ischemic stroke, SAH, intracerebral hemorrhage (ICH), death, and increase the likelihood of favorable functional outcome (mRS 0–2 vs. 3–6, or 0–1 vs. 2–6, or equivalent)?

Analysis of current evidence

We found no study matching our inclusion criteria to answer this question (Figure 22). The literature search did not find any randomized controlled trial of endovascular or surgical treatment in symptomatic IAD patients with an unruptured intracranial dissecting aneurysm and isolated headache or any observational studies meeting our minimal sample size criteria. The main reason for literature exclusion was small sample size (Methods) and that most studies on IAD focused on either patients with ischemic stroke, SAH, or mixed group of patients.

Additional information

Five observational studies with small sample sizes of patients with an intracranial vertebral artery dissecting aneurysm and isolated headache (of which two only were comparative) were reported from Japan or Korea (Table 7).^{98–102} None of 21 patients with an intracranial dissecting aneurysm and isolated headache who had endovascular (n = 20) or surgical treatment (n = 1) experienced intracranial hemorrhage or ischemic stroke. All 21 patients had an excellent (mRS 0–1) or good (mRS 0–2 or equivalent) functional outcome, and none died during the observation period. In contrast, 2 out of 68 patients who did not have endovascular or surgical treatment suffered SAH on the day following headache or neck pain onset, 1 experienced new clinical symptoms due to mass effect, but none of the 68 patients had an ischemic stroke. Twelve patients, including the 2 with new-onset SAH, experienced good functional outcome, but there was no information on functional outcome in the remaining 52

patients, including 1 with new-onset clinical symptoms due to mass effect. None died during the observation period.

Supporting Information to the Expert consensus statement

Although data are scarce, in recent years most IAD patients with intracranial dissecting aneurysm and isolated headache are treated medically. Because the natural course of these patients is generally favorable, and because of the competing risks of both SAH and ischemic stroke, no antithrombotic treatment, but close monitoring has been proposed.^{13,102,103} Endovascular or surgical treatment might be considered if the dissecting aneurysm increases in size or signs of compression occur.

Evidence-based Recommendation

For symptomatic IAD patients with an intracranial dissecting aneurysm and isolated headache (no TIA, no acute ischemic stroke, no SAH), there is uncertainty over the benefits and risks of endovascular or surgical treatment and therefore it is not possible to make a recommendation.

Quality of evidence: Very low ⊕

Strength of recommendation: -

Expert consensus statement

For symptomatic IAD patients with an intracranial aneurysm and isolated headache, all but one expert suggest against endovascular or surgical treatment unless the aneurysmal size increases significantly on follow-up imaging, or signs of compression occur.

PICO 5: In symptomatic extracranial artery dissection (EAD) and intracranial artery dissection (IAD) patients with ischemic stroke, transient ischemic attack (TIA), retinal ischemia, or local symptoms only, and without subarachnoid hemorrhage (SAH), is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of ischemic stroke, death, higher likelihood of favorable functional outcome (mRS 0–2 vs. 3–6, or 0–1 vs. 2–6, or equivalent), and no increased risk of intracerebral hemorrhage (ICH), SAH, or other major bleeding?

Analysis of current evidence

We identified 2 completed RCTs addressing PICO5 (Figure 23, Table 8),^{32–34} both of which compared antiplatelets to vitamin K antagonists [VKA] in EAD, and of

Figure 22. PRISMA flow chart of study selection for PICO4. “Wrong patient population” in COVIDENCE corresponds to studies that do not match criteria for this PICO.

which one has two publications (with 3 months and 1 year follow-up),^{32,33} and no ongoing RCT. We found no RCT data on the use of direct acting oral anticoagulants (DOACs) in EAD nor any RCTs on the IAD population.

The CADISS study (Cervical Artery Dissection in Stroke Study) was a multicenter, randomized controlled, open label study designed to show feasibility of a RCT in EAD patients. CADISS included 250 EAD patients and randomly allocated participants to either antiplatelets or anticoagulant (VKA with or without bridging with unfractionated heparin [UFH] or low molecular weight heparin [LMWH]).³² The

specific choice of drug within either treatment arm was left to the discretion of the treating physicians. The intention-to-treat population comprised 126 patients in the antiplatelet group and 124 patients in the anticoagulation group. Antiplatelet treatment was heterogeneous with 22% of patients receiving aspirin alone, 33% receiving clopidogrel alone, 28% receiving both aspirin and clopidogrel, 16% receiving aspirin and dipyridamole, and one patient receiving dipyridamole only. In the anticoagulation group, 90% of patients received heparin and warfarin, whereas 10% received warfarin alone. Regarding the primary study endpoint

Table 7. Summary of observational studies findings relevant for PICO4.

Study	Population	Mean age (SD)	Intervention	Comparison	Follow-up	Outcome	IPD					
							intervention			IPD control		
							N	N	N	event	total	event
Ahn 2006 ¹⁰⁰	Intracranial VA dissection with and without SAH (n = 14), Korea	49.0 (8.7)	Endovascular, n = 5 (Stent placement alone, n = 4; stent-assisted coiling, n = 1)	-	NA	Mortality Excellent functional outcome	0	5	0	0	0	0
						Intracranial hemorrhage	0	5	0	0	0	0
						Ischemic stroke	0	5	0	0	0	0
Naito 2002 ⁹⁹	Intracranial VA dissection without SAH (n = 21), Japan	53.6 (9.4)	Endovascular, n = 1 (Proximal occlusion using balloons)	Conservative	NA	Mortality Good functional outcome	0	1	0	8	8	8
						Intracranial hemorrhage	0	1	2	8	8	8
						Ischemic stroke	0	1	0	8	8	8
Nakazawa 2011 ⁹⁸	Intracranial VA dissection with and without SAH (n = 47), Japan	51.2 (7.4)	Endovascular, n = 4 (Stent only, n = 2; stent-assisted coiling, n = 2) and surgery, n = 1 (VA ligation with OA-PICA anastomosis, n = 1)	Observation	NA	Mortality Good functional outcome	0	5	0	4	4	4
						Intracranial hemorrhage	0	5	0	4	4	4
Nam 2015 ¹⁰¹	Intracranial VA dissection with and without SAH (n = 26), Korea	Median 50s (range, 30s to 60s)	Endovascular, n = 10 (Proximal occlusion, n = 1; internal trapping, n = 1; stent only, n = 1; stent-assisted coiling, n = 1; modified semi-jailing technique with stent and coiling, n = 5; coiling followed by stent, n = 1)	-	30.4 months (SD, 22.9)	Mortality Excellent functional outcome	0	10	0	0	0	0
						Intracranial hemorrhage	0	10	0	0	0	0
						Ischemic stroke	0	10	0	0	0	0
Kobayashi 2014 ¹⁰²	Vertebral IAD without stroke and with headache (out of n = 113 patients in total with non-stroke vertebral IAD), Japan	51.1 (10.6)	-	Conservatively managed without intervention and antithrombotic therapy	mean follow-up of 2.9 years (range, 27 days to 8 years)	Mortality Intracranial hemorrhage	0	0	0	56	56	56
						Ischemic stroke	0	0	0	56	56	56

Figure 23. PRISMA flow chart of study selection for PICO5. *these included 2 consecutive publications on the same trial (3 months and 1 year follow-up)^{32,33} and 4 observational studies on partly overlapping samples.^{16,17,108,113}; “wrong population” in COVIDENCE corresponds to studies that do not match criteria for this PICO.

(ipsilateral stroke or death), there was no statistically significant group difference.³² Within the 3-month study period, ischemic stroke occurred in three (of 126) patients in the antiplatelet group and in one (of 124) patient in the anticoagulation group (OR, 0.335 [95% CI, 0.006–4.233], $p = 0.63$). There was one major hemorrhage (SAH) in the anticoagulation group. No major hemorrhage was observed in the antiplatelet group.³² In the subsequent 12-month follow-up analysis, there were two additional ischemic strokes (one in each treatment arm) yielding again no statistically significant difference in the primary endpoint between groups.³³

In about 20% of the participants, the diagnosis of EAD was not confirmed by central adjudication either due to an alternative cause being identified or because imaging was not of sufficient quality to be confident of the diagnosis. However, per-protocol analysis excluding these subjects showed similar results.^{32,33} Although this is beyond the scope of PICO5, it can be noted that secondary analyses showed no association between treatment allocation (antiplatelets vs. anticoagulants) and whether dissecting aneurysms at baseline persisted at follow-up or whether new dissecting aneurysms

Table 8. Summary of RCT (per-protocol) findings relevant for PICO5.

Trial	Study	Population	Intervention			Comparison			F-up	Outcome	IPD				Effect estimate (OR/RR/HR)
			Mean age $\pm$ SD (yrs)	Type	NIHSS at baseline	N patients with IS at admission*	NIHSS at baseline	N patients with IS at admission*			Intervention		IPD Control		
											Type	Type	event	N total	
Engelter 2021 ³⁴	TREAT-CAD; RCT (multicentre, randomized, open-label, non-inferiority trial, testing non-inferiority of ASP to VKA)	symptomatic, MRI-verified EAD with symptom onset within 2 weeks	ASP group: 46.7 $\pm$ 10.2; VKA group: 45.5 $\pm$ 11.6	AC (VKA ^a) for 90 days	2.5 $\pm$ 4.3	43 (52%)	2.1 $\pm$ 2.9	47 (52%)	3 months	IS	0	82	7	91	0.07 (0.00–1.21)
Markus 2019 ³³	CADISS; RCT (multicenter, randomized controlled, open label study to show feasibility of a RCT in EAD patients)	EAD with symptom onset within 7 days	49 $\pm$ 12 (range 18–87) AC group: 48.1 $\pm$ 11; Antiplatelet group: 48.5 $\pm$ 12	AC (heparin followed by warfarin) for 3 months	NA	77 (80%)	NA	74 (73%)	1 year	IS	1	96	4	101	0.26 (0.03–2.33)
Markus 2015 ³²									3 months	(Any) stroke	1	96	3	101	0.346 (0.01–4.39)

*presenting signs and symptoms.

^aPhenprocoumon, acenocoumarol, or warfarin; target international normalized ratio [INR] 2.0–3.0 (heparin or LMWH recommended until target INR is reached); AC: anticoagulants; AP: antiplatelet; ASP: aspirin; ICH: intracerebral hemorrhage; IS: ischemic stroke; VKA: vitamin K antagonists.

developed,²⁸ or whether stenosis present at baseline showed recanalization.³³

Power calculations based on the per-protocol results from CADISS and the composite endpoint of stroke, death and major bleeding suggested a sample size of about 10000 (4876 per arm) would be required for a definitive phase 3 RCT.³²

TREAT-CAD was a multicenter randomized controlled therapy trial comparing aspirin to VKA in the treatment of EAD. Participants were randomly assigned to receive either aspirin (300mg/d) or VKA (with or without bridging with UFH or LMWH) for 3 months. To increase the number of endpoints and therefore power it included imaging marker events (new diffusion weighted imaging [DWI] or susceptibility weighted imaging [SWI]/T2*-lesions during follow-up as compared to baseline imaging),¹⁰⁴ as well as clinical endpoints (acute ischemic stroke, major intra- or extracranial hemorrhage, and death) in a composite study endpoint. The main results of TREAT-CAD were published in March 2021.³⁴ The primary analyses in TREAT-CAD were performed in the per-protocol population which comprised 173 patients (of 194 in the intention-to-treat population) of which 91 were allocated to aspirin and 82 were allocated to VKA. The primary composite endpoint occurred in 21 (23%) patients in the aspirin group and in 12 (15%) in the VKA group (absolute difference 8% [95% CI, -4 to 21], non-inferiority $p = 0.55$).³⁴ Accordingly, non-inferiority of aspirin was not shown. All ischemic strokes

($n = 7$) occurred in the ASA group, whereas the only major—though extracranial (gastrointestinal bleeding)—hemorrhage occurred in the VKA group. There were no deaths in either group. Five of the 7 ischemic strokes in the aspirin group occurred (or recurred) on day 1 after treatment onset, suggesting the importance of early initiation of antithrombotic treatment—whichever the clinician might choose.

As active treatment was stopped at 3 months in both trials, our meta-analyses combined the per-protocol results from CADISS and TREAT-CAD at 3 months follow-up^{32,34} for ischemic stroke (Figure 24), major bleeding (Figure 25), and the composite outcome of ischemic stroke, major bleeding, or death (Figure 26). There was no significant difference between the two treatment groups for the composite endpoint, ischemic stroke or major bleeding. In the anticoagulation group, the odds of developing the composite endpoint was OR, 0.35 (95% CI, 0.08–1.63), while that of developing ischemic stroke was OR, 0.18 (95% CI, 0.03–1.10), and that of major bleeding OR, 3.28 (95% CI, 0.34–31.80). There were no deaths at 3 months in either study. Information on functional outcome was available in TREAT-CAD only, where no difference was observed between both arms for excellent or good functional outcome.³⁴ It should be noted that both were phase 2 RCTs and underpowered to show small, but still important, differences between the two treatment regimens.

Figure 24. Meta-analysis of effects on risk of ischemic stroke of anticoagulant versus antiplatelet treatment at the acute phase of EAD in RCTs (at 3 months).

Figure 25. Meta-analysis of effects on risk of major bleeding of anticoagulant versus antiplatelet treatment at the acute phase of EAD in RCTs (at 3 months).

Figure 26. Meta-analysis of effects on risk of ischemic stroke, major bleeding or death of anticoagulant versus antiplatelet treatment at the acute phase of EAD in RCTs (at 3 months).

Figure 27. Risk of bias of RCTs for PICO5.

We have based our grading recommendation on data from RCTs alone. The risk of bias is described in [Figure 27](#), [Table 9](#)). The overall risk of bias was rated as not serious, inconsistency not serious, indirectness non-serious, imprecision mostly as serious to very serious due to small sample sizes in these phase 2 trials; the observation certainty was rated as moderate and the importance as critical.

Overall, the two phase 2 RCTs have shown no difference between the benefits and risks of anticoagulants versus antiplatelets in the acute phase of symptomatic EAD. Although these were underpowered to show smaller differences in risks and benefits between the two approaches, they indicate that, based on current evidence, clinicians can prescribe either option.

Table 9. Quality of results in RCTs relevant for PICO5.

Certainty assessment												Certainty	Importance
No. of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	No. of patients	Anticoagulants	Antiplatelets	Effect Relative (95% CI)	Absolute (95% CI)		
PICO 5.1 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of ischemic stroke major bleeding and death at 3 months in RCTs?													
2	randomized trials	not serious	not serious	not serious	serious ^a	none	3/178 (1.7%)	110/192 (57.3%)	OR 0.35 (0.08 to 1.63)	253 fewer per 1,000 (from 476 fewer to 113 more)		⊕⊕⊕⊕○ MODERATE	CRITICAL
PICO 5.2 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of death at 3 months in RCTs?													
2	randomized trials	not serious	not serious	not serious	serious ^b	none	0/178 (0.0%)	0/192 (0.0%)	not estimable			⊕⊕⊕⊕○ MODERATE	CRITICAL
PICO 5.3 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a higher likelihood of mRS 0–2 at 3 months in RCTs?													
1	randomized trials	not serious	not serious	not serious	very serious ^c	none	80/82 (97.6%)	88/91 (96.7%)	OR 1.36 (0.22 to 8.37)	9 more per 1,000 (from 101 fewer to 29 more)		⊕⊕⊕⊕○ LOW	CRITICAL
PICO5.4 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of recurrent ischemic stroke at 3 months in RCTs?													
2	randomized trials	not serious	not serious	not serious	serious ^a	none	1/178 (0.6%)	10/192 (5.2%)	OR 0.18 (0.03 to 1.10)	42 fewer per 1,000 (from 50 fewer to 5 more)		⊕⊕⊕⊕○ MODERATE	CRITICAL
PICO5.5 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of intracranial hemorrhage at 3 months in RCTs?													
1	randomized trials	not serious	not serious	not serious	serious ^b	none	0/82 (0.0%)	0/91 (0.0%)	not estimable			⊕⊕⊕⊕○ MODERATE	CRITICAL
PICO5.6 Is anticoagulant versus antiplatelet therapy in the acute phase associated with a reduced risk of major hemorrhage at 3 months in RCTs?													
2	randomized trials	not serious	not serious	not serious	very serious ^c	none	2/178 (1.1%)	0/192 (0.0%)	OR 3.28 (0.34 to 31.80)	0 fewer per 1,000 (from 0 fewer to 0 fewer)		⊕⊕⊕⊕○ LOW	CRITICAL

CI: Confidence interval; OR: Odds ratio.

^aWide confidence intervals.^bOdds ratio not calculated because of zero events in the intervention and control arms.^cVery wide confidence intervals.

It should be noted that, while there is no data demonstrating that antithrombotic treatment is better than no treatment in preventing stroke in EAD patients, as antiplatelets have been shown to be superior to placebo in preventing stroke after TIA or ischemic stroke in general,^{105,106} it wouldn't be possible to carry out a trial on this in EAD patients with ischemic stroke or TIA. Prior to the CADISS trial, a survey of physicians was conducted on what they would use to treat EAD (with and without cerebral ischemia).¹⁰⁷ It shows that almost all would give either anticoagulants or antiplatelets emphasizing the fact that a trial would not be possible. Further adding to this evidence, a Cochrane review across observational case series of EAD showed that most patients with EAD of the internal carotid artery were treated with some type of antithrombotic treatment.⁴ Although in an observational setting prone to bias, it is noteworthy that the small group of patients with no antithrombotic treatment had a much higher mortality rate (25%) than patients under antiplatelets or anticoagulants (each 1.8%).⁴

Additional information

In addition to the 2 RCTs, we identified 13 observational studies (Figure 23, Table 10), reporting on some of the outcomes mentioned in the PICO question.^{16,17,45,108-117} When these studies were conducted on overlapping samples, only the largest study available for a given outcome was used in the analysis.^{16,17,108,113} These were uncontrolled studies and suffered from potential bias, particularly selection bias in choice of treatment (Supplementary Table 1.5). Most studies comprised EAD patients only, while a few included a minority of patients with IAD.^{16,17,110,112,115} However, there was insufficient information on outcomes in EAD or IAD patients specifically; hence, no separate meta-analysis could be conducted.

No significant differences in the treatment approaches were shown for any of the outcomes, except for a borderline significant higher rate of good functional outcome with anticoagulation based on three small studies (Figures 28–33). Previous systematic reviews and meta-analyses across observational studies, using different selection criteria and heterogeneous methodologies, did not indicate clear superiority of either approach but showed inconsistent findings.^{4,114,118-120} Compared to the present study, most of these meta-analyses had used much lower sample size thresholds for inclusion (Methods), including encompassing small case series.

Only one retrospective observational study fulfilling our inclusion criteria compared use of DOACs to traditional anticoagulants (VKA or heparin) and antiplatelet agents in EAD patients.¹¹¹ Of the 149 included patients, 39, 70, and 40 were treated with a DOAC, traditional anticoagulant, and antiplatelet agent, respectively. There was no significant

difference between the DOAC and other treatment groups for the risk of recurrent stroke (2 in the DOAC group vs. 1 in each of the other groups, $p = 0.822$). There were more major bleeding events in the traditional anticoagulant group (11.4%) compared to the DOAC (0.0%) and antiplatelet (2.5%) groups ($p = 0.034$). Three patients treated with DOAC and none in the other groups had a worsened degree of stenosis on follow-up imaging (8.6 vs. 0.0 vs. 0.0%, $p = 0.019$), but patients treated with DOAC more often had a severe stenosis or occlusion at baseline than patients in the other treatment groups.¹¹¹

Dual antiplatelet therapy with clopidogrel and aspirin has been shown to reduce the recurrent stroke rate in TIA and minor stroke in two large RCTs (CHANCE¹²¹ and POINT¹²²) when given within 24 hours of symptom onset, compared to aspirin alone. These trials have shown a reduction the risk of stroke¹²¹ or major ischemic events (ischemic stroke, myocardial infarction, or death from an ischemic vascular event)¹²² at 3 months with a significant increase in the risk of major bleeding in the POINT trial.¹²² Most stroke events, and the separation in stroke incidence curves between the aspirin plus clopidogrel arm and aspirin alone arm occurred within 10 days of randomization, while the separation in incidence of bleeding continued to increase throughout the treatment period and there was no net benefit from continuing treatment beyond 3 weeks.^{105,123} Similarly, the THALES trial showed that in patients with a mild to moderate acute non-cardioembolic ischemic stroke or TIA who were not undergoing thrombolysis, the risk of the composite outcome of stroke or death was lower with the combination of ticagrelor and aspirin than with aspirin alone within 30 days, although severe bleeding was more frequent with ticagrelor.¹²⁴ No subgroup information was available on patients with EAD/IAD in these trials. Recent ESO guidelines on TIA management recommend short term dual antiplatelet therapy with aspirin and clopidogrel over monotherapy, subsequently followed by monotherapy, in patients with acute non-cardioembolic high risk TIA.¹⁰⁵

There is no good information on the duration of antithrombotic treatment in EAD patients. The ESO-Karolinska guideline recommended (with a grade C) to pursue antithrombotic treatment for 6–12 months.³⁰ It was further suggested that in patients in whom full recanalization of the dissected artery has occurred and there have been no recurrent symptoms, stopping antithrombotic treatment may be considered.³⁰ In case of a residual dissecting aneurysm or stenosis, long-term antiplatelet treatment was recommended.³⁰ There is no new data justifying an update of this. Of note, in the two RCTs on EAD the randomized intervention (anticoagulants or antiplatelets) had a duration of 3 months.

Regarding IAD, there is no RCT comparing antiplatelet agents to anticoagulants at the acute phase. As proposed previously,¹³ higher theoretical risk of SAH than EAD and

Table 10. Summary of observational study findings relevant for PICO5.

Study	Population	Mean age	Intervention		Comparison	Duration of follow-up	Outcome	IPD				
			Type	Anticoagulants				Type	intervention		IPD control	
									N	event	N	event
Arauz 2006 ¹⁰⁸	EAD	35.4	Anticoagulants	Aspirin	Median 19 (6–120) months	Mortality mRS 0–2 Ischemic stroke Intracranial hemorrhage Major bleeding	Mortality mRS 0–2 Ischemic stroke Major bleeding	0 34 3 0	48 48 48 48	4 38 3 0	82 82 82 82	
Arauz 2013** ¹⁷	EAD+IAD (vertebral artery dissection and ischemic stroke)	37.9 ± 8.5	Anticoagulants	Aspirin	Median 46.4 months (range : 6 to 175)	mRS 0–2 Ischemic stroke Major bleeding	mRS 0–2 Ischemic stroke Major bleeding	0 39 1	48 49 49	0 43 0	82 50 50	
Arnold 2006** ¹⁶	EAD+IAD (vertebral artery dissection and ischemic stroke)	43 ± 9	Warfarin	Aspirin	3 months	mRS 0–1	mRS 0–1	48	58	27	33	
Ast 1993 ¹⁰⁹	EAD (internal carotid artery)	NA	Anticoagulants ^a	Antiplatelets ^a	NA	Mortality ^a Ischemic stroke ^a Intracranial hemorrhage ^a	Mortality ^a Ischemic stroke ^a Intracranial hemorrhage ^a	0 1 0	30 30 30	0 0 0	21 21 21	
Beletsky 2003** ¹¹⁰	EAD+IAD	Men: 46 ± 12 yrs Women: 42 ± 10 yrs	Anticoagulants	Aspirin	10.0±3.5 months	Mortality Ischemic stroke	Mortality Ischemic stroke	2 2	71 71	0 1	23 23	
Caprio 2014 ¹¹¹	EAD	Anticoagulants : 41.4 ± 15.0 Antiplatelets : 48.1 ± 13.2	AC	Aspirin	Median: 7.5 months for clinical follow-up (5 months for radiological follow-up)	Ischemic stroke Major bleeding	Ischemic stroke Major bleeding	1 8	70 70	1 1	40 40	
Daou 2017** ¹¹²	EAD+IAD	47	Anticoagulants	Antiplatelets	Mean : 24 months	mRS 0–2 Ischemic stroke Major bleeding	mRS 0–2 Ischemic stroke Major bleeding	85 2 2	88 88 88	182 6 4	195 195 195	
Dziewas 2003 ⁴⁵	EAD	43.2 ± 11.2	Anticoagulants	Antiplatelets	Max 6 months	Mortality ^a Ischemic stroke ^a Intracranial hemorrhage ^a	Mortality ^a Ischemic stroke ^a Intracranial hemorrhage ^a	1 0 1	71 71 71	0 0 0	7 7 7	
Gensicke 2015 ¹¹⁷	EAD	46 (40–52)*	Anticoagulants	Antiplatelets	30 days	Major bleeding ^a mRS 0–1	Major bleeding ^a mRS 0–1	0 19	71 25	0 30	7 43	
Georgiadis 2009 ¹¹³	EAD	Anticoagulants group: 46 ± 10; Aspirin group: 46 ± 11	Anticoagulants	Aspirin	3 months	Mortality Ischemic stroke Intracranial hemorrhage Major bleeding	Mortality Ischemic stroke Intracranial hemorrhage Major bleeding	0 2 2 2	202 202 202 202	0 0 0 0	96 96 96 96	

(continued)

Table 10. (continued)

Study	Population	Mean age	Intervention		Comparison	Duration of follow-up	Outcome	IPD			
								intervention		IPD control	
			Type	Type	Type			N	N	N	N
Kennedy 2012 ¹¹⁴	EAD	Anticoagulants: 43 Antiplatelets: 45	Anticoagulants	Antiplatelets			Mortality	0	28	0	59
							Ischemic stroke	1	28	1	59
Metso 2009** ¹¹⁵	EAD+IAD	46.6 (range: 15–79)	Anticoagulants	Antiplatelets ^a	4 years (median 3.1, range: 0.1–13.2)		Mortality ^a	1	140	0	4
							Ischemic stroke ^a	3	140	1	4
							Intracranial hemorrhage ^a	2	140	0	4
Ramchand 2018 ¹¹⁶	EAD	47 ± 15	Anticoagulants	Antiplatelets	Median 3.5 months		Ischemic stroke	1	36	1	39
							Intracranial hemorrhage	1	36	0	39
							Major bleeding	1	36	0	39
							Intracerebral bleeding	1	36	0	39

*Median (IQR) ; IQR : interquartile range .

**these studies include a small subset of IAD patients (60 patients in Daou 2017; 18 patients in Arnold 2006; 3 patients in Beletsky 2003; 95 patients in Metso 2009; 27 patients with pure IAD and 23 patients with IAD+EAD in Arauz 2013).

^aFigures from previously published meta-analysis and review.^{4,119}

Figure 28. Meta-analysis of effects on mortality of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

Figure 29. Meta-analysis of effects on good functional outcome (mRS 0–2) of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

Figure 30. Meta-analysis of effects on excellent functional outcome (mRS 0–1) of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

Figure 31. Meta-analysis of effects on ischemic stroke of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

Figure 32. Meta-analysis of effects on intracranial hemorrhage of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

Figure 33. Meta-analysis of effects on major bleedings of anticoagulant versus antiplatelet treatment at the acute phase of EAD in observational studies.

the superiority of aspirin over anticoagulants in the acute phase of ischemic stroke in general are empirical arguments in favor of prescribing aspirin rather than anticoagulants in patients with IAD and cerebral ischemia. In case of recurrent thromboembolic events despite aspirin, dual antiplatelet treatment or anticoagulants could be considered. Before initiation of antithrombotic treatment in patients with IAD and cerebral ischemia, a lumbar puncture can be performed if neuroimaging cannot formally rule out minor SAH.¹³ In IAD patients without SAH and cerebral ischemia, or in rare cases when both SAH and cerebral ischemia are present, close monitoring without antithrombotic treatment has been suggested.¹⁰³

Evidence-based Recommendation

In the acute phase of symptomatic EAD, we recommend that clinicians can prescribe either anticoagulants or antiplatelet therapy.

Quality of evidence: Moderate $\oplus\oplus$

Strength of recommendation: Strong for an intervention $\uparrow\uparrow$

Expert consensus statement

For symptomatic EAD patients treated with anticoagulants in the acute phase, all but three experts felt that DOACs could be used in place of vitamin K antagonists.

For symptomatic EAD patients, all but one expert felt it was reasonable to use dual antiplatelet therapy with aspirin and clopidogrel in the acute phase in patients with TIA or minor stroke and restricted to a few weeks.

For acute symptomatic IAD patients with ischemic stroke or TIA and no SAH, all but one expert felt antiplatelet agents had a better risk/benefit ratio than anticoagulants.

PICO 6: In extracranial artery dissection (EAD) patients and in intracranial artery dissection (IAD) patients without subarachnoid hemorrhage (SAH) does endovascular or surgical treatment of a stenosis or a dissecting aneurysm outside the acute phase versus optimal medical treatment

Figure 34. PRISMA flow chart of study selection for PICO6 “Wrong study design, language, setting, intervention, outcomes, and (patient) population” in COVIDENCE corresponds to studies that do not match criteria for this PICO.

alone reduce the risk of death, ischemic stroke, intracerebral hemorrhage (ICH), and SAH?

Analysis of current evidence

The literature search identified no RCT and no comparative observational studies meeting our sample size criteria (Figure 34).

Importantly, there is overwhelming data on the very low rate of recurrent ischemic events in EAD patients.^{23,125} In

the combined CADISP and French-Swiss EAD cohorts, out of 1,931 EAD patients, overall sixty patients (3.1%) had new-onset cerebral ischemia during follow-up at 3 to 6 months, of whom 33 patients (1.7%) an ischemic stroke, and 32 patients (1.7%) a TIA.²³ In the CADISS trial, comparing anticoagulation to antiplatelet therapy in 250 EAD patients, the rate of ischemic events during follow-up was 2% at 3 months and 2.4% at 1 year.^{28,33} The presence of dissecting aneurysms was not associated with a higher risk of stroke during follow-up: at 12 months, stroke occurred in

2.1% of patients with and 3.2% of patients without dissecting aneurysm (OR, 0.84 [95% CI, 0.10–7.31], $p = 0.88$).²⁸

Additional information

We identified two small observational single-arm studies addressing PICO6 (Table 11) matching our inclusion criteria in terms of sample size.^{126,127} Although the time lapse was not clearly indicated, we considered only subsets of patients who were reported to be in a chronic disease phase. Persistent symptoms despite appropriate medical management (ischemic or thromboembolic events), severe stenosis, or developing/expanding dissecting aneurysm were the most common indications for endovascular or surgical intervention. Müller et al. (48 chronic EAD patients) reported surgical treatment using saphenous vein graft replacement after resection of the diseased internal carotid segment was performed in 80% of the patients.¹²⁶ Moon et al. (51 chronic EAD patients) reported endovascular treatment with stenting and/or coiling.¹²⁷ Antithrombotic therapy after endovascular or surgical treatment differed across studies. In the study by Müller et al., low-dose heparin was administered for 1 week in all patients.¹²⁶ In the study by Moon et al., patients were treated with dual antiplatelet therapy (oral aspirin, 325 mg and clopidogrel, 75 mg) after stenting for at least 6 months.¹²⁷ Follow-up duration ranged between 1 and 190 months (1–190 months in Müller et al.¹²⁶ and 1–146 months in Moon et al.¹²⁷). Regarding outcomes after the intervention, data are scarce. Müller et al. described one ischemic stroke (2%), one intracranial hemorrhage (2%) and three deaths (6%) amongst the 48 chronic EAD patients.¹²⁶ Moon et al. reported no stroke or TIA during follow-up amongst the 51 chronic EAD patients.¹²⁷

In addition to the CADISS trial mentioned above, several observational studies add to the evidence that residual stenosis and dissecting aneurysms have a benign prognosis in EAD patients. In a long-term follow-up study of patients with carotid EAD and persistent (46 patients, 6.2 years) or transient (46 patients, 7.2 years) severe stenosis or occlusion, there was no relation between residual arterial pathology and stroke rate.¹²⁸ A systematic review gathering 166 EAD patients with internal carotid artery dissection (from 9 studies on <50 patients each) followed the evolution of dissecting aneurysms and reported that they rarely become symptomatic and rarely increase in size:²⁷ 3% increased in size, 52% remained unchanged in diameter, 21% diminished in size, 19% completely resolved, 2% thrombosed, and 3% were repaired surgically; 2% underwent surgery 0.5–5.0 years later.

Much less data is available on IAD. In 52 IAD patients with asymptomatic vertebrobasilar dissecting aneurysms discovered incidentally (thus considered here as chronic, out of a total of 113 IAD patients followed up for 2.9 years

[range, 27 days to 8 years]), Kobayashi et al. observed no stroke during follow-up and only one case of clinical deterioration due to mass effect.¹⁰² Aneurysm size remained unchanged in 96% of the 52 patients during follow-up.¹⁰²

Evidence-based Recommendation

In post-acute EAD patients with residual stenosis or dissecting aneurysm, there is uncertainty over the benefits and risks of endovascular or surgical treatment and therefore it is not possible to make a recommendation.

Quality of evidence: Very low ⊕

Strength of recommendation: -

Expert consensus statement

Given the overwhelming evidence of a very low rate of recurrent ischemic events in post-acute EAD patients under medical treatment and the lack of evidence for an impact of residual stenosis or dissecting aneurysms on the rate of these events, all but one expert suggest against routine use of endovascular/surgical treatment in these patients: based on current limited evidence, endovascular/surgical treatment of post-acute EAD patients with residual stenosis or dissecting aneurysms may be carefully considered in exceptional situations, such as recurrent ischemic events despite optimal antithrombotic therapy or expanding dissecting aneurysms causing compression, after assessment by a multidisciplinary team (neurologist, neuroradiologist, neurosurgeon, and neurointerventionalist).

Discussion

We produced a total of four recommendations, of which one was with moderate, one with low, and two with very low quality of evidence. These were based on the results of two phase 2 RCTs,³²⁻³⁴ and 26 comparative observational studies^{16,17,44-48,56-58,67-70,108-117,126,127} that were identified through systematic review and met our inclusion criteria (Supplementary Table 7). For one PICO no study met our inclusion criteria but 5 smaller studies, of which 2 comparative, were analyzed (PICO4). All recommendations are summarized in Table 12.

The three recommendations with low and very low quality of evidence were derived from observational studies: on IVT (PICO1) and EVT (PICO2) at the acute phase of EAD-related ischemic stroke, and on endovascular or surgical interventions in IAD with SAH (PICO3).

Table 11. Summary of observational single-arm study findings relevant for PICO6.

Study	Population	Mean age	Intervention	Duration follow-up	Mean delay between EAD and intervention	Reasons for intervention	Outcome	IPD intervention		
								N event	N total	Frequency (95% CI)
Moon 2017 ¹²⁷	EAD	44.9 (5–76)	Stent placement; coil occlusion of parent artery; stenting with contralateral vessel occlusion	41.6 months (1–146 months) ^a	chronic (no more details)	Failure of medical treatment ^b	Recurrent stroke or TIA	0	51	0%
Müller 2000 ¹²⁶	EAD (carotid)	45.4	Surgical (Saphenous vein graft replacement after resection of the diseased internal carotid segment in 80%)	70 months (1–190 months)	9 months (2 months–5 years)	Aneurysm formation at distal end of the dissection near the skull base; stenosis >80%	Mortality Intracranial hemorrhage Recurrent ischemic stroke	3	48*	0.06 (0.00–0.13) 0.02 (0.00–0.06) 0.02 (0.00–0.06)

* 1 patient with acute dissection was removed from the descriptive analysis.
^afor the overall study (116 EAD patients, of whom only 51 were in the post-acute phase).
^bMedical therapy was considered to have failed if ischemic or thromboembolic events continued during clinical follow-up, but not if pseudoaneurysms enlarged, unless they caused thromboembolic phenomena.

Table 12. Summary table of all evidence-based recommendations.

Evidence-based recommendations	Quality of evidence	Strength of recommendation
In patients with symptomatic EAD with acute ischemic stroke within 4.5 hours of onset, we suggest using intravenous thrombolysis with alteplase, if the standard inclusion / exclusion criteria are met.	Low ⊕⊕	Weak for an intervention ↑?
In patients with symptomatic IAD with acute ischemic stroke within 4.5 hours of onset, there is insufficient data to provide a recommendation.	Very low ⊕	-
In acute ischemic stroke patients with EAD and large vessel occlusion of the anterior circulation we suggest using MT	Very low ⊕	Weak for an intervention ↑?
In acute ischemic stroke patients with IAD there is insufficient data to provide a recommendation regarding the use of EVT.	Very low ⊕	-
In patients suffering IAD with SAH we suggest early surgical or endovascular intervention. There is insufficient data to provide a recommendation on the type of intervention to prioritize and the precise time window.	Very low ⊕	Weak for an intervention ↑?
For symptomatic IAD patients with an intracranial dissecting aneurysm and isolated headache (no TIA, no acute ischemic stroke, no SAH), there is uncertainty over the benefits and risks of endovascular or surgical treatment and therefore it is not possible to make a recommendation.	Very low ⊕	-
In the acute phase of symptomatic EAD, we recommend that clinicians can prescribe either anticoagulants or antiplatelet therapy.	Moderate ⊕⊕⊕	Strong for an intervention ↑↑
In post-acute EAD patients with residual stenosis or dissecting aneurysms, there is uncertainty over the benefits and risks of endovascular or surgical treatment and therefore it is not possible to make a recommendation.	Very low ⊕	-

Considering the nature of the evidence described in these observational studies, the very low incidence of EAD/IAD making it a major challenge to gather higher quality evidence, the strong benefit/risk ratio of some interventions in ischemic stroke overall, and information on the natural history of the disease, we nevertheless proposed recommendations for intervention (graded as weak because of the observational nature of the data), as we felt that running a clinical trial to more definitely answer these questions would be unethical. The recommendation with moderate quality of evidence was based on two phase 2 RCTs comparing anticoagulation and antiplatelet therapy in EAD (PICO5). While we acknowledge that additional trials may be warranted in the future to account for therapeutic innovations and novel antithrombotic regimens, we considered that it would be unrealistic to enroll the required number of patients to complete a phase 3 trial on an uncommon condition such as EAD. As results were consistent across the two published RCTs, we therefore also rated this recommendation as strong. For PICO4 and PICO6, addressing endovascular or surgical intervention in IAD with isolated headache, and in the post-acute phase of EAD, the quality of evidence was so low that no recommendation could be made.

In addition to the 4 recommendations, we proposed 9 expert consensus statements. All expert consensus statements are summarized in [Table 13](#). For all of these there was agreement by at least 8 out of 11 voting members of the MWG.

Many of these pertain to the management of IAD for which the quality of evidence was too low to provide recommendations, specifically for the use of IVT (PICO1) and EVT (PICO2) in the acute phase of IAD-related ischemic stroke, antiplatelet agents over anticoagulation in IAD with TIA or ischemic stroke (PICO5), and endovascular or surgical interventions when the IAD is associated with headache only (without SAH or cerebral ischemia, PICO4). Some expert consensus statements were aimed at sharing expert opinion on the implementation of certain recommendations, such as the use of direct anticoagulants in symptomatic EAD or of dual antiplatelet therapy in the first weeks after EAD-related TIA or minor stroke (PICO5), or the decision to conduct EVT for the treatment of the EAD/IAD lesion (PICO2). Finally, some expert consensus statements recommended in the absence of clear data to use complementary local expertise through multidisciplinary assessment on complex questions such as the choice of the type of intervention in IAD-related SAH (PICO3) or the decision to perform an intervention for post-acute EAD in the uncommon situation of recurrent ischemic events despite optimal antithrombotic therapy or expanding dissecting aneurysms causing compression (PICO6).

To our knowledge, this is the first comprehensive European guideline on the management of EAD and IAD, implementing GRADE methodology.³⁵ Recommendations and expert consensus statements are based on the work of a large group of experts with broad expertise relevant to the PICO questions, spanning across disciplines (neurology,

Table 13. Summary table of all expert consensus statements.

Expert consensus statement based on voting by all WMG members	Votes
In patients with an acute ischemic stroke suspected to be caused by IAD, all but one expert suggested that IVT should be considered, after ruling out standard contra-indications, including subtle signs of subarachnoid bleeding on brain imaging.	10/11
For patients with EAD as a cause of acute ischemic stroke with intracranial large vessel occlusion of the anterior circulation, all but one expert suggest EVT (other than MT) for the treatment of the EAD lesion in case of carotid occlusion without patent circle of Willis or with recurrent embolism.	10/11
For patients with IAD as a cause of acute ischemic stroke with intracranial large vessel occlusion of the anterior circulation, all experts suggest EVT (other than MT) for the treatment of the IAD lesion at the hyperacute phase after assessing the risk/benefit ratio based on the location of the dissection and bleeding risk.	11/11
Different types of surgical and endovascular treatment methods can be used for treating IAD with SAH. In the absence of RCTs and considering the limited data from observational studies with high risk of bias, all experts suggest that the choice of intervention type in acute IAD-related SAH should ideally be the result of a multidisciplinary assessment.	11/11
For symptomatic IAD patients with an intracranial aneurysm and isolated headache, all but one expert suggest against endovascular or surgical treatment unless the aneurysmal size increases significantly on follow-up imaging, or signs of compression occur.	10/11
For symptomatic EAD patients treated with anticoagulants in the acute phase, all but three experts felt that DOACs could be used in place of vitamin K antagonists.	8/11
For symptomatic EAD patients, all but one expert felt it was reasonable to use dual antiplatelet therapy with aspirin and clopidogrel in the acute phase in patients with TIA or minor stroke and restricted to a few weeks.	10/11
For acute symptomatic IAD with ischemic stroke or TIA and no SAH, all but one expert felt antiplatelet agents had a better risk/benefit ratio than anticoagulants.	10/11
Given the overwhelming evidence of a very low rate of recurrent ischemic events in post-acute EAD patients under medical treatment and the lack of evidence for an impact of residual stenosis or dissecting aneurysms on the rate of these events, all but one expert suggest against routine use of endovascular/surgical treatment in these patients: based on current limited evidence, endovascular/surgical treatment of post-acute EAD may be carefully considered in exceptional situations, such as recurrent ischemic events despite optimal antithrombotic therapy or expanding dissecting aneurysms causing compression, after assessment by a multidisciplinary team (neurologist, neuroradiologist, neurosurgeon, and neurointerventionalist).	10/11

neurointervention, neurosurgery, and neuroepidemiology), countries, and continents. The latter was specifically important for this guideline due to important differences in the epidemiology of EAD and IAD between European and East Asian ancestry populations and more extensive experience with IAD management in East Asia. While most of our PICOs relied on observational data only, we have applied very strict inclusion criteria, in contrast to previously published meta-analyses on this topic, most of which also included small case series. Importantly, this guideline also presents the first meta-analysis of the only two RCTs comparing anticoagulants and antiplatelets for the management of EAD.

We acknowledge limitations. The observational nature and small size of most of the studies described in this guideline is a major limitation. For some PICOs and outcomes, there was an important heterogeneity between included studies. Some studies combined EAD and IAD without providing the data that would enable a separate analysis of each. Not all studies used strict diagnostic criteria for dissection, especially older ones and for IAD. Important heterogeneity was noted in follow-up duration. For the two published RCTs, more elaborate individual level data meta-analysis could be warranted, but this was beyond

the scope of the present study. Finally, the random effects meta-analyses included in this manuscript were based on raw data extracted from the publications. Secondary meta-analyses using adjusted ORs provided in the manuscripts or by contacting the authors could be of interest but are beyond the scope of the present work.

While conducting phase 3 RCTs on the management of EAD/IAD seems unrealistic based on power estimates from the CADISS trial, further prospective collection of data in the setting of international registries could provide important “real life” estimates of the impact of different therapeutic approaches on EAD/IAD-related outcomes. Some specific therapeutic questions, for which there is extremely limited data, could not be addressed, such as the management of traumatic dissection (severe trauma) and lifestyle guidance (on management of future pregnancies and limiting practice of physical activity). We equally did not address the management of EAD/IAD in the rare context of inherited connective tissue diseases, such as vascular Ehlers-Danlos syndrome.¹²⁹ We also did not examine the question of optimal vascular risk factor management after EAD, especially pertaining to hypertension and hypercholesterolemia after EAD-related stroke (and systematic prescription of antihypertensive and lipid lowering

therapies). The impact of intervention on imaging-based outcomes (recurrent dissection; new ischemic lesions on DWI sequences; frequency of normalization or stability of vessel patency on imaging) was not covered.

Plain language summary

Extracranial artery dissection (EAD) and intracranial artery dissection (IAD) are characterized by bleeding in the wall of an artery in the neck or in the brain. They represent a leading cause of stroke in young and middle-aged adults, either due to clotting in the artery near the wall hematoma (for EAD and IAD) leading to vessel occlusion and causing an ischemic stroke, or due to rupture of the artery (for IAD) causing intracranial bleeding in the subarachnoid space surrounding the brain tissue (subarachnoid hemorrhage).

When patients with EAD present with an acute ischemic stroke, it is recommended that they receive the same acute phase treatments as other patients with ischemic stroke, which mainly aim at removing the blood clot from the vessel to restore normal blood flow. This can be done by using intravenous thrombolysis (i.e., injection of a product that dissolves the clot) and by retrieving the clot mechanically (mechanical thrombectomy). After these emergency treatments other medications are indicated for several months to avoid new clotting within the dissected artery. For this, it is recommended that physicians can choose between two types of blood thinning medications, antiplatelets (moderate blood thinning), or anticoagulants (intensive blood thinning). Indeed, two recent clinical trials where EAD patients were randomly assigned to receive either antiplatelets or anticoagulants found these two options to be equally efficient. For patients with IAD who suffer from a subarachnoid hemorrhage, an urgent intervention is recommended (either endovascular or surgical) as it is known to substantially reduce the risk of rebleeding and improve survival. Finally, for patients with IAD who present with headache only (without any ischemic stroke or subarachnoid hemorrhage) and in EAD patients with residual vessel narrowing or widening (so-called dissecting aneurysm) after the acute phase, there isn't enough evidence to make recommendations on the use of endovascular treatment or surgery to improve the outcome of patients.

Acknowledgements

We thank the reviewers of the PICO Qs and final text, editorial support of the ESJ and ESO staff Luzia Balmer and Sabrina Mutter for excellent administrative support. We thank Terry Quinn for serving as an external advisor to our MWG.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Funding for the development of these guidelines was provided by the European Stroke Organisation, Basel, Switzerland. SD received funding from Grants from EU H2020, JPND, ANR (French National funding agency, general and 'Investment for the Future' Programme ANR-18-RHUS-002, Claude Pompidou foundation, for research on environmental and genetic determinants of stroke, cerebral small vessel disease, MRI-markers of brain aging, cognitive decline and dementia PI on national investment for the future programme on covert cerebral small vessel disease (RHU SHIVA). JM received funding from NIH/NINDS 1U24NS107228 and 1U24NS107156; NCMRR R01HD105718. JK received funding honorary/travel support compensation from Boehringer Ingelheim and Pfizer, all outside the current topic. MA received Grants from Swiss Heart Foundation and Swiss National Science Foundation; Co-Chief Investigator of TREAT-CAD <https://clinicaltrials.gov/ct2/show/NCT02046460>, honoraria for lectures from AstraZeneca, Bayer, Covidien, and Medtronic, and honoraria for scientific advisory boards of Amgen, Bayer, Bristol Myers, Squibb, Daiichi Sankyo, Medtronic, Novartis, and Sanofi all outside the current topic. SE received grants from Swiss National Science Foundation, Swiss Heart Foundation, Stroke Funds Basel, University Hospital Basel University of Basel, Academic Society Basel [Freiwillige Akademische Gesellschaft Basel, FAG], he is Chief Investigator of TREAT-CAD <https://clinicaltrials.gov/ct2/show/NCT02046460>. He also received funding for an investigator-initiated clinical study from Daiichi-Sankyo outside the current topic, and speaker Honorary/Travel support/Advisory board compensation from Bayer, Boehringer Ingelheim, Daiichi-Sankyo, BMS/Pfizer, and MindMaze; all outside the current topic. MM received grants from the French National Agency of Health, and the French ministry of health. He also received Speaker Honorary/Travel support/Advisory board compensation from Amgen, Boehringer Ingelheim, Air Liquide, Acticor Biotech; all outside the current topic. AP received grants from the University of Brescia; ASST Spedali Civili, Brescia; Nova Science Publishers, Inc, Hauppauge NY. He also received funding from the Associazione per la Lotta alla Trombosi e alle Malattie Cardiovascolari (ALT) for the Italian Project on Stroke at Young Age (IPSYS); from the Regione Lombardia for the STROKOVID Network, and Speaker Honorary/Travel support: compensation from Bayer, Boehringer Ingelheim, MAGI, all outside the topic of the current paper. HSM received Grants from Stroke Association, is chief Investigator of CADISS and received funding from BIBA for conference organization and is an Editor in Chief at the International Journal of Stroke. AB received grants from Minister of Health and is Chief Investigator of GENOMA and SENECA project. MK received speaker honorary from Bayer Yakuhin, BMS/Pfizer, Daiichi-Sankyo, and Nippon Boehringer Ingelheim and research funds from Takeda, Daiichi-Sankyo, Nippon Boehringer Ingelheim, Astellas, and Shionogi.

Contributorship

All listed authors have contributed to the preparation and writing of the manuscript.

ORCID iDs

Stephanie Debette <https://orcid.org/0000-0001-8675-7968>

Janika Kõrv <https://orcid.org/0000-0002-6074-0727>

Supplemental material

Supplemental material for this article is available online.

References

- Putala J, Metso AJ, Metso TM, et al. Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke: the Helsinki young stroke registry. *Stroke* 2009; 40: 1195–1203.
- Leys D, Bandu L, Hénon H, et al. Clinical outcome in 287 consecutive young adults (15 to 45 years) with ischemic stroke. *Neurology* 2002; 59: 26–33.
- Caplan LR. Dissections of brain-supplying arteries. *Nat Clin Pract Neurol* 2008; 4: 34–42.
- Lyrer P and Engelter S. Antithrombotic drugs for carotid artery dissection. *Cochrane Database Syst Rev* 2010; 6(10): CD000255.
- Schievink WI. Spontaneous dissection of the carotid and vertebral arteries. *N Engl J Med* 2001; 344: 898–906.
- Arnold M, Fischer U and Boussier MG. Treatment issues in spontaneous cervicocephalic artery dissections. *Int J Stroke* 2011; 6: 213–218.
- Debette S, Metso T, Pezzini A, et al. Association of vascular risk factors with cervical artery dissection and ischemic stroke in young adults. *Circulation* 2011; 123: 1537–1544.
- Debette S, Goeggel Simonetti B, Schilling S, et al. Familial occurrence and heritable connective tissue disorders in cervical artery dissection. *Neurology* 2014; 83: 2023–2031.
- Debette S and Leys D. Cervical-artery dissections: predisposing factors, diagnosis, and outcome. *Lancet Neurol* 2009; 8: 668–678.
- Walmsley JG and Canham PB. Orientation of nuclei as indicators of smooth muscle cell alignment in the cerebral artery. *Blood Vessels* 1979; 16: 43–51.
- Lee RM. Morphology of cerebral arteries. *Pharmacol Ther* 1995; 66: 149–173.
- Chen M and Caplan L. Intracranial dissections. *Front Neurol Neurosci* 2005; 20: 160–173.
- Debette S, Compter A, Labeyrie MA, et al. Epidemiology, pathophysiology, diagnosis, and management of intracranial artery dissection. *Lancet Neurol* 2015; 14: 640–654.
- Béjot Y, Daubail B, Debette S, et al. Incidence and outcome of cerebrovascular events related to cervical artery dissection: the Dijon stroke registry. *Int J Stroke* 2014; 9: 879–882.
- Lee VH, Brown RD Jr, Mandrekar JN, et al. Incidence and outcome of cervical artery dissection: a population-based study. *Neurology* 2006; 67: 1809–1812.
- Arnold M, Boussier MG, Fahrni G, et al. Vertebral artery dissection: presenting findings and predictors of outcome. *Stroke* 2006; 37: 2499–2503.
- Arauz A, Ruiz A, Pacheco G, et al. Aspirin versus anti-coagulation in intra- and extracranial vertebral artery dissection. *Eur J Neurol* 2013; 20: 167–172.
- Kwak JH, Choi JW, Park HJ, et al. Cerebral artery dissection: spectrum of clinical presentations related to angiographic findings. *Neurointervention* 2011; 6: 78–83.
- Huang YC, Chen YF, Wang YH, et al. Cervicocranial arterial dissection: experience of 73 patients in a single center. *Surg Neurol* 2009; 72(Suppl 2): S20–27. discussion S27.
- Debette S, Kamatani Y, Metso TM, et al. Common variation in PHACTR1 is associated with susceptibility to cervical artery dissection. *Nat Genet* 2015; 47: 78–83.
- Mizutani T. Natural course of intracranial arterial dissections. *J Neurosurg* 2011; 114: 1037–1044.
- Caplan LR, Baquis GD, Pessin MS, et al. Dissection of the intracranial vertebral artery. *Neurology* 1988; 38: 868–877.
- Compter A, Schilling S, Vaineau CJ, et al. Determinants and outcome of multiple and early recurrent cervical artery dissections. *Neurology* 2018; 91: e769–e780.
- Patel RR, Adam R, Maldjian C, et al. Cervical carotid artery dissection: current review of diagnosis and treatment. *Cardiol Rev* 2012; 20: 145–152.
- Benninger DH, Gandjour J, Georgiadis D, et al. Benign long-term outcome of conservatively treated cervical aneurysms due to carotid dissection. *Neurology* 2007; 69: 486–487.
- Pelkonen O, Tikkakoski T, Leinonen S, et al. Extracranial internal carotid and vertebral artery dissections: angiographic spectrum, course and prognosis. *Neuroradiology* 2003; 45: 71–77.
- Paraskevas KI, Batchelder AJ and Naylor AR. Fate of distal false aneurysms complicating internal carotid artery dissection: a systematic review. *Eur J Vasc Endovasc Surg* 2016; 52: 281–286.
- Larsson SC, King A, Madigan J, et al. Prognosis of carotid dissecting aneurysms: results from CADISS and a systematic review. *Neurology* 2017; 88: 646–652.
- Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke: a guideline for healthcare professionals from the American heart Association/American stroke association. *Stroke* 2019; 50: e344–e418.
- Ahmed N, Steiner T, Caso V, et al. Recommendations from the ESO-Karolinska stroke update conference, Stockholm 13–15 November 2016. *Eur Stroke J* 2017; 2: 95–102.
- Berge E, Whiteley W, Audebert H, et al. European stroke organisation (ESO) guidelines on intravenous thrombolysis for acute ischaemic stroke. *Eur Stroke J* 2021; 6: I–LXII.

32. Markus HS, Hayter E, Levi C, et al. Antiplatelet treatment compared with anticoagulation treatment for cervical artery dissection (CADISS): a randomised trial. *Lancet Neurol* 2015; 14: 361–367.
33. Markus HS, Levi C, King A, et al. Antiplatelet therapy vs anticoagulation therapy in cervical artery dissection: the cervical artery dissection in stroke study (CADISS) randomized clinical trial final results. *JAMA Neurol* 2019; 76: 657–664.
34. Engelter ST, Traenka C, Gensicke H, et al. Aspirin versus anticoagulation in cervical artery dissection (TREAT-CAD): an open-label, randomised, non-inferiority trial. *Lancet Neurol* 2021; 20: 341–350.
35. Guyatt GH, Oxman AD, Schünemann HJ, et al. GRADE guidelines: a new series of articles in the journal of clinical epidemiology. *J Clin Epidemiol* 2011; 64: 380–382.
36. Ntaios G, Bornstein NM, Caso V, et al. The European stroke organisation guidelines: a standard operating procedure. *Int J Stroke* 2015; 10(Suppl A100): 128–135.
37. Kaatz S, Ahmad D, Spyropoulos AC, et al. Definition of clinically relevant non-major bleeding in studies of anticoagulants in atrial fibrillation and venous thromboembolic disease in non-surgical patients: communication from the SSC of the ISTH. *J Thromb Haemost* 2015; 13: 2119–2126.
38. Higgins JP, Altman DG, Gøtzsche PC, et al. The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343: d5928.
39. Murad MH, Mustafa RA, Schünemann HJ, et al. Rating the certainty in evidence in the absence of a single estimate of effect. *Evid Based Med* 2017; 22: 85–87.
40. Wardlaw JM, Murray V, Berge E, et al. Thrombolysis for acute ischaemic stroke. *Cochrane Database Syst Rev* 2014; 2014: CD000213.
41. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995; 333: 1581–1587.
42. Fusco MR and Harrigan MR. Cerebrovascular dissections—a review part I: spontaneous dissections. *Neurosurgery* 2011; 68: 242–257. discussion 257.
43. Intracerebral hemorrhage after intravenous t-PA therapy for ischemic stroke. The NINDS t-PA stroke study group. *Stroke* 1997; 28: 2109–2118.
44. Engelter ST, Dallongeville J, Kloss M, et al. Thrombolysis in cervical artery dissection—data from the cervical artery dissection and ischaemic stroke patients (CADISP) database. *Eur J Neurol* 2012; 19: 1199–1206.
45. Dziewas R, Konrad C, Dräger B, et al. Cervical artery dissection—clinical features, risk factors, therapy and outcome in 126 patients. *J Neurol* 2003; 250: 1179–1184.
46. Bernardo F, Nannoni S, Strambo D, et al. Efficacy and safety of endovascular treatment in acute ischemic stroke due to cervical artery dissection: a 15-year consecutive case series. *Int J Stroke* 2019; 14: 381–389.
47. Qureshi AI, Chaudhry SA, Hassan AE, et al. Thrombolytic treatment of patients with acute ischemic stroke related to underlying arterial dissection in the United States. *Arch Neurol* 2011; 68: 1536–1542.
48. Jensen J, Salottolo K, Frei D, et al. Comprehensive analysis of intra-arterial treatment for acute ischemic stroke due to cervical artery dissection. *J Neurointerv Surg* 2017; 9: 654–658.
49. Bernardo F, Nannoni S, Strambo D, et al. Intravenous thrombolysis in acute ischemic stroke due to intracranial artery dissection: a single-center case series and a review of literature. *J Thromb Thrombolysis* 2019; 48: 679–684.
50. Zinkstok SM, Vergouwen MD, Engelter ST, et al. Safety and functional outcome of thrombolysis in dissection-related ischemic stroke: a meta-analysis of individual patient data. *Stroke* 2011; 42: 2515–2520.
51. Engelter ST, Rutgers MP, Hatz F, et al. Intravenous thrombolysis in stroke attributable to cervical artery dissection. *Stroke* 2009; 40: 3772–3776.
52. Tsivgoulis G, Zand R, Katsanos AH, et al. Safety and outcomes of intravenous thrombolysis in dissection-related ischemic stroke: an international multicenter study and comprehensive meta-analysis of reported case series. *J Neurol* 2015; 262: 2135–2143.
53. Doijiri R, Yokota C, Suzuki R, et al. Intravenous recombinant tissue plasminogen activator thrombolysis in acute ischemic stroke due to middle cerebral artery dissection. *J Stroke Cerebrovasc Dis* 2012; 21: e917–e919.
54. Moon Y, Lee JH, Cho HJ, et al. Intravenous thrombolysis in a patient with acute ischemic stroke attributable to intracranial dissection. *Neurologist* 2012; 18: 136–138.
55. Demaerschalk BM, Kleindorfer DO, Adeoye OM, et al. Scientific rationale for the inclusion and exclusion criteria for intravenous Alteplase in acute ischemic stroke: a statement for healthcare professionals from the American heart association/American stroke association. *Stroke* 2016; 47: 581–641.
56. Li S, Zi W, Chen J, et al. Feasibility of thrombectomy in treating acute ischemic stroke because of cervical artery dissection. *Stroke* 2018; 49: 3075–3077.
57. Marnat G, Lapergue B, Sibon I, et al. Safety and outcome of carotid dissection stenting during the treatment of tandem occlusions: a pooled analysis of TITAN and ETIS. *Stroke* 2020; 51: 3713–3718.
58. Traenka C, Jung S, Gralla J, et al. Endovascular therapy versus intravenous thrombolysis in cervical artery dissection ischemic stroke - results from the SWISS registry. *Eur Stroke J* 2018; 3: 47–56.
59. Boisseau W, Fahed R, Lapergue B, et al. Predictors of parenchymal hematoma after mechanical thrombectomy: a multicenter study. *Stroke* 2019; 50: 2364–2370.
60. Papanagiotou P, Haussen DC, Turjman F, et al. Carotid stenting with antithrombotic agents and intracranial thrombectomy leads to the highest recanalization rate in patients with acute stroke with tandem lesions. *JACC Cardiovasc Interv* 2018; 11: 1290–1299.
61. Turc G, Bhogal P, Fischer U, et al. European stroke organisation (ESO) - European society for minimally invasive

- neurological therapy (ESMINT) guidelines on mechanical thrombectomy in acute ischaemic stroke endorsed by stroke alliance for Europe (SAFE). *Eur Stroke J* 2019; 4: 6–12.
62. Marnat G, Mourand I, Eker O, et al. Endovascular management of tandem occlusion stroke related to internal carotid artery dissection using a distal to proximal approach: insight from the RECAST study. *AJNR Am J Neuroradiol* 2016; 37: 1281–1288.
 63. Karam A, Bricout N, Khyeng M, et al. Safety and outcome of mechanical thrombectomy in ischaemic stroke related to carotid artery dissection. *J Neurol* 2021.
 64. Nogueira RG and Tsivgoulis G. Large vessel occlusion strokes after the DIRECT-MT and SKIP trials: is the alteplase syringe half empty or half full? *Stroke* 2020; 51: 3182–3186.
 65. Marnat G, Bühlmann M, Eker OF, et al. Multicentric experience in distal-to-proximal revascularization of tandem occlusion stroke related to internal carotid artery dissection. *AJNR Am J Neuroradiol* 2018; 39: 1093–1099.
 66. Yamada M, Kitahara T, Kurata A, et al. Intracranial vertebral artery dissection with subarachnoid hemorrhage: clinical characteristics and outcomes in conservatively treated patients. *J Neurosurg* 2004; 101: 25–30.
 67. Anxionnat R, de Melo Neto JF, Bracard S, et al. Treatment of hemorrhagic intracranial dissections. *Neurosurgery* 2003; 53: 289–300. discussion 300–281.
 68. Rabinov JD, Hellinger FR, Morris PP, et al. Endovascular management of vertebrobasilar dissecting aneurysms. *AJNR Am J Neuroradiol* 2003; 24: 1421–1428.
 69. Mizutani T, Aruga T, Kirino T, et al. Recurrent subarachnoid hemorrhage from untreated ruptured vertebrobasilar dissecting aneurysms. *Neurosurgery* 1995; 36: 905–911. discussion 912–903.
 70. Zhao WY, Krings T, Alvarez H, et al. Management of spontaneous haemorrhagic intracranial vertebrobasilar dissection: review of 21 consecutive cases. *Acta Neurochir (Wien)* 2007; 149: 585–596. discussion 596.
 71. Albuquerque FC, Fiorella DJ, Han PP, et al. Endovascular management of intracranial vertebral artery dissecting aneurysms. *Neurosurg Focus* 2005; 18: E3.
 72. Endo H, Matsumoto Y, Kondo R, et al. Medullary infarction as a poor prognostic factor after internal coil trapping of a ruptured vertebral artery dissection. *J Neurosurg* 2013; 118: 131–139.
 73. Kashiwazaki D, Ushikoshi S, Asano T, et al. Long-term clinical and radiological results of endovascular internal trapping in vertebral artery dissection. *Neuroradiology* 2013; 55: 201–206.
 74. Kim BM, Shin YS, Kim SH, et al. Incidence and risk factors of recurrence after endovascular treatment of intracranial vertebrobasilar dissecting aneurysms. *Stroke* 2011; 42: 2425–2430.
 75. Fang YB, Zhao KJ, Wu YN, et al. Treatment of ruptured vertebral artery dissecting aneurysms distal to the posterior inferior cerebellar artery: stenting or trapping? *Cardiovasc Intervent Radiol* 2015; 38: 592–599.
 76. Ikeda DS, Marlin ES, Shaw A, et al. Endovascular flow diversion therapy for an actively hemorrhaging aneurysm after intraoperative rupture. *J Clin Neurosci* 2015; 22: 1839–1842.
 77. Lee JM, Kim TS, Joo SP, et al. Endovascular treatment of ruptured dissecting vertebral artery aneurysms—long-term follow-up results, benefits of early embolization, and predictors of outcome. *Acta Neurochir (Wien)* 2010; 152: 1455–1465.
 78. Mohammadian R, Taheraghdam AA, Sharifipour E, et al. Endovascular treatment of intracranial artery dissection: clinical and angiographic follow-up. *Neurol Res Int* 2013; 2013: 968380.
 79. Motoyama Y, Takamura Y, Park HS, et al. Appropriate flow reduction for unilateral ruptured vertebral artery dissection by proximal clipping to prevent rebleeding and medullary infarction. *World Neurosurg* 2019; 130: e627–e633.
 80. Nakamura H, Fujinaka T, Nishida T, et al. Endovascular therapy for ruptured vertebral artery dissecting aneurysms: results from nationwide, retrospective, multi-center registries in Japan (JR-NET3). *Neurol Med Chir (Tokyo)* 2019; 59: 10–18.
 81. Ohkuma H, Suzuki S and Ogane K. Study group of the association of cerebrovascular disease in Tohoku J. Dissecting aneurysms of intracranial carotid circulation. *Stroke* 2002; 33: 941–947.
 82. Oran I, Cinar C, Yağci B, et al. Ruptured dissecting aneurysms arising from non-vertebral arteries of the posterior circulation: endovascular treatment perspective. *Diagn Interv Radiol* 2009; 15: 159–165.
 83. Park W, Kwon DH, Ahn JS, et al. Treatment strategies for dissecting aneurysms of the posterior cerebral artery. *Acta Neurochir (Wien)* 2015; 157: 1633–1643.
 84. Raper DMS, Caldwell J, Brew S, et al. A comparison of endovascular strategies in the treatment of ruptured vertebral artery aneurysms. *J Clin Neurosci* 2020; 75: 168–175.
 85. Satow T, Ishii D, Iihara K, et al. Endovascular treatment for ruptured vertebral artery dissecting aneurysms: results from Japanese registry of neuroendovascular therapy (JR-NET) 1 and 2. *Neurol Med Chir (Tokyo)* 2014; 54(Suppl 2): 98–106.
 86. Shibukawa M, Sakamoto S, Kiura Y, et al. Ruptured vertebral artery dissecting aneurysms treated with endovascular treatment. *Hiroshima J Med Sci* 2009; 58: 55–60.
 87. So TY, Mitchell PJ, Dowling RJ, et al. Efficacy, complications and clinical outcome of endovascular treatment for intracranial intradural arterial dissections. *Clin Neurol Neurosurg* 2014; 117: 6–11.
 88. Zang Y, Wang C, Zhang Y, et al. Long-term follow-up study of 35 cases after endovascular treatment for vertebrobasilar dissecting aneurysms. *Clin Neurol Neurosurg* 2015; 137: 121–131.
 89. Zhang Y, Tian Z, Sui B, et al. Endovascular treatment of spontaneous intracranial fusiform and dissecting aneurysms: outcomes related to imaging classification of 309 cases. *World Neurosurg* 2017; 98: 444–455.

90. Zhao KJ, Zhao R, Huang QH, et al. The interaction between stent(s) implantation, PICA involvement, and immediate occlusion degree affect symptomatic intracranial spontaneous vertebral artery dissection aneurysm (sis-VADA) recurrence after reconstructive treatment with stent(s)-assisted coiling. *Eur Radiol* 2014; 24: 2088–2096.
91. Lee JW, Jung JY, Kim YB, et al. Spontaneous dissecting aneurysm of the intracranial vertebral artery: management strategies. *Yonsei Med J* 2007; 48: 425–432.
92. Fukasawa I, Sasaki H and Nukui H. Surgical treatment for ruptured vertebral artery dissecting aneurysms. *Neurol Med Chir (Tokyo)* 1998; 38(Suppl 1): 104–106.
93. Sikkema T, Uyttenboogaart M, van Dijk JM, et al. Clinical features and prognosis of intracranial artery dissection. *Neurosurgery* 2015; 76: 663–670. discussion 670–661.
94. Steiner T, Juvela S, Unterberg A, et al. European stroke organization guidelines for the management of intracranial aneurysms and subarachnoid haemorrhage. *Cerebrovasc Dis* 2013; 35: 93–112.
95. Bederson JB, Connolly ES Jr, Batjer HH, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a statement for healthcare professionals from a special writing group of the stroke council, American heart association. *Stroke* 2009; 40: 994–1025.
96. Nakamura H, Fujinaka T, Nishida T, et al. Endovascular therapy for ruptured vertebral artery dissecting aneurysms: results from nationwide, retrospective, multi-center registries in Japan (JR-NET3). *Neurol Med Chir (Tokyo)* 2019; 59: 10–18.
97. Ohkuma H, Suzuki S and Ogane K. Dissecting aneurysms of intracranial carotid circulation. *Stroke* 2002; 33: 941–947.
98. Nakazawa T, Takeichi Y, Yokoi T, et al. Treatment of spontaneous intradural vertebral artery dissections. *Neuroradiol J* 2011; 24: 699–711.
99. Naito I, Iwai T and Sasaki T. Management of intracranial vertebral artery dissections initially presenting without subarachnoid hemorrhage. *Neurosurgery* 2002; 51: 930–937. discussion 937–938.
100. Ahn JY, Han IB, Kim TG, et al. Endovascular treatment of intracranial vertebral artery dissections with stent placement or stent-assisted coiling. *AJNR Am J Neuroradiol* 2006; 27: 1514–1520.
101. Nam KH, Ko JK, Cha SH, et al. Endovascular treatment of acute intracranial vertebral artery dissection: long-term follow-up results of internal trapping and reconstructive treatment using coils and stents. *J Neurointerv Surg* 2015; 7: 829–834.
102. Kobayashi N, Murayama Y, Yuki I, et al. Natural course of dissecting vertebrobasilar artery aneurysms without stroke. *AJNR Am J Neuroradiol* 2014; 35: 1371–1375.
103. Kai Y, Nishi T, Watanabe M, et al. Strategy for treating unruptured vertebral artery dissecting aneurysms. *Neurosurgery* 2011; 69: 1085–1091. discussion 1091–1082.
104. Traenka C, Gensicke H, Schaedelin S, et al. Biomarkers and antithrombotic treatment in cervical artery dissection - design of the TREAT-CAD randomised trial. *Eur Stroke J* 2020; 5: 309–319.
105. Fonseca AC, Merwick A, Dennis M, et al. European stroke organisation (ESO) guidelines on management of transient ischaemic attack. *European Stroke Journal* 2021; 1–24.
106. International Stroke Trial Collaborative Group (IST): a randomised trial of aspirin, subcutaneous heparin, both, or neither among 19435 patients with acute ischaemic stroke. *Lancet* 1997; 349: 1569–1581.
107. Menon RK, Markus HS and Norris JW. Results of a UK questionnaire of diagnosis and treatment in cervical artery dissection. *J Neurol Neurosurg Psychiatry* 2008; 79: 612.
108. Arauz A, Hoyos L, Espinoza C, et al. Dissection of cervical arteries: long-term follow-up study of 130 consecutive cases. *Cerebrovasc Dis* 2006; 22: 150–154.
109. Ast G, Woimant F, Georges B, et al. Spontaneous dissection of the internal carotid artery in 68 patients. *Eur J Med* 1993; 2: 466–472.
110. Beletsky V, Nadareishvili Z, Lynch J, et al. Cervical arterial dissection: time for a therapeutic trial? *Stroke* 2003; 34: 2856–2860.
111. Caprio FZ, Bernstein RA, Alberts MJ, et al. Efficacy and safety of novel oral anticoagulants in patients with cervical artery dissections. *Cerebrovasc Dis* 2014; 38: 247–253.
112. Daou B, Hammer C, Mouchtouris N, et al. Anticoagulation vs antiplatelet treatment in patients with carotid and vertebral artery dissection: a study of 370 patients and literature review. *Neurosurgery* 2017; 80: 368–379.
113. Georgiadis D, Arnold M, von Buedingen HC, et al. Aspirin vs anticoagulation in carotid artery dissection: a study of 298 patients. *Neurology* 2009; 72: 1810–1815.
114. Kennedy F, Lanfranconi S, Hicks C, et al. Antiplatelets vs anticoagulation for dissection: CADISS nonrandomized arm and meta-analysis. *Neurology* 2012; 79: 686–689.
115. Metso TM, Metso AJ, Salonen O, et al. Adult cervicocerebral artery dissection: a single-center study of 301 Finnish patients. *Eur J Neurol* 2009; 16: 656–661.
116. Ramchand P, Mullen MT, Bress A, et al. Recanalization after extracranial dissection: effect of antiplatelet compared with anticoagulant therapy. *J Stroke Cerebrovasc Dis* 2018; 27: 438–444.
117. Gensicke H, Ahlhelm F, Jung S, et al. New ischaemic brain lesions in cervical artery dissection stratified to antiplatelets or anticoagulants. *Eur J Neurol* 2015; 22: 859–865.
118. Sarikaya H, da Costa BR, Baumgartner RW, et al. Antiplatelets versus anticoagulants for the treatment of cervical artery dissection: Bayesian meta-analysis. *PLoS One* 2013; 8: e72697.
119. Chowdhury MM, Sabbagh CN, Jackson D, et al. Antithrombotic treatment for acute extracranial carotid artery dissections: a meta-analysis. *Eur J Vasc Endovasc Surg* 2015; 50: 148–156.

120. Menon R, Kerry S, Norris JW, et al. Treatment of cervical artery dissection: a systematic review and meta-analysis. *J Neurol Neurosurg Psychiatry* 2008; 79: 1122–1127.
121. Wang Y, Zhao X, Liu L, et al. Clopidogrel with aspirin in acute minor stroke or transient ischemic attack. *N Engl J Med* 2013; 369: 11–19.
122. Johnston SC, Easton JD, Farrant M, et al. Clopidogrel and aspirin in acute ischemic stroke and high-risk TIA. *N Engl J Med* 2018; 379: 215–225.
123. Hao Q, Tampi M, O'Donnell M, et al. Clopidogrel plus aspirin versus aspirin alone for acute minor ischaemic stroke or high risk transient ischaemic attack: systematic review and meta-analysis. *BMJ* 2018; 363: k5108.
124. Johnston SC, Amarenco P, Denison H, et al. Ticagrelor and aspirin or aspirin alone in acute ischemic stroke or TIA. *N Engl J Med* 2020; 383: 207–217.
125. Schievink WI, Mokri B and O'Fallon WM. Recurrent spontaneous cervical-artery dissection. *N Engl J Med* 1994; 330: 393–397.
126. Müller BT, Luther B, Hort W, et al. Surgical treatment of 50 carotid dissections: indications and results. *J Vasc Surg* 2000; 31: 980–988.
127. Moon K, Albuquerque FC, Cole T, et al. Stroke prevention by endovascular treatment of carotid and vertebral artery dissections. *J Neurointerv Surg* 2017; 9: 952–957.
128. Kremer C, Mosso M, Georgiadis D, et al. Carotid dissection with permanent and transient occlusion or severe stenosis: long-term outcome. *Neurology* 2003; 60: 271–275.
129. Adham S, Billon C, Legrand A, et al. Spontaneous cervical artery dissection in vascular ehlers-danlos syndrome: a cohort study. *Stroke* 2021; 52: 1628–1635.