Published in final edited form as:

Acta Biochim Pol. 2021 March 15; 68(1): 15–22. doi:10.18388/abp.2020_5587.

Crystal structure of timothy grass allergen Phl p 12.0101 reveals an unusual profilin dimer*

Andrea O'Malley¹, A. Brenda Kapingidza¹, Noah Hyduke¹, Coleman Dolamore¹, Krzysztof Kowal^{2,3}, Maksymilian Chruszcz^{1,✉}

¹Department of Chemistry and Biochemistry, University of South Carolina, Columbia, SC 29208, USA

²Department of Allergology and Internal Medicine, Medical University of Białystok, Białystok 15-276, Poland

³Department of Experimental Allergology and Immunology, Medical University of Białystok, Białystok 15-276, Poland

Abstract

Timothy grass pollen is a source of potent allergens. Among them, Phl p 1 and Phl p 5 are thought to be the most important, as a majority of timothy grass-allergic individuals have IgE antibodies directed against these two allergens. The profilin from timothy grass (Phl p 12) has been registered as a minor allergen, with up to 35% of individuals in populations of grass pollen allergic patients showing IgE binding to Phl p 12. Profilins are primarily minor allergens and are known for a high likelihood of co-sensitization as well as cross-reactivity situations caused by their sequence and structure similarity. The crystal structure of Phl p 12.0101 was determined and it revealed that this allergen may form an unusual dimer not previously observed among any profilins. For example, the Phl p 12 dimer has a completely different geometry and interface when compared with the latex profilin (Hev b 8) dimer that has its crystal structure determined. The structure of Phl p 12.0101 is described in the context of allergenic sensitization and allergy diagnostics. Moreover, the structure of the Phl p 12.0101 dimer is discussed, taking into account the production of recombinant allergens and their storage.

Keywords

grass allergy; allergen; profilin; *Phleum pratense*; pollen-food syndrome; oral allergy syndrome

*This manuscript is dedicated as a Birthday tribute to Prof. Wlodek Minor to whom the corresponding author owes (among many other things) interest in molecular allergology.

✉ chruszcz@mailbox.sc.edu .

Conflict of Interest

A.O, A.B.K., N.H, C.D., and M.C, declare no conflict of interest. K.K. received lecture fees from: ALK Abello, Astra Zeneca, Berlin Chemie, Chiesi, Emma, Hal Allergy, Meda Pharma, Orion Pharma, and royalties from UpToDate.

INTRODUCTION

Timothy (*Phleum pratense*) is a perennial grass native to most of Europe and found in most of the United States. Timothy grass grows well in wet, humid areas and is quite winter-hardy (Ogle et al., 2011). It is grown commercially as a common animal fodder, especially for horses, other grazing animals like cattle, and for domesticated pets such as guinea pigs. It can also be used as erosion control in some regions (Ogle et al., 2011). In America, timothy is mainly cultivated in the Pacific Northwest (humid areas around the Puget Sound), the Northeast, and the Midwest (Hitchcock & Chase, 1951). Due to this ubiquitous nature of timothy grass, it is a potent source of allergens for many patients sensitized to the grass (Scaparrotta et al., 2013; Rossi et al., 2008; Sekerkova et al., 2012; Almeida et al., 2019). Sensitization and reaction to pollen from timothy grass is a contributor to allergic rhinitis and other symptoms such as rhinoconjunctivitis. Due to high pollen counts and long pollen seasons, grasses are the most important seasonal allergen source in Europe and the USA (Kowal, 2020; Durham et al., 2012; Maloney et al., 2014). Moreover, high cumulative exposure to grass pollens each season leads to development of a robust IgE response to a wide range of grass proteins in allergic patients (Hatzler et al., 2012).

Currently, ten *P. pratense* allergens have been registered by the World Health Organization and International Union of Immunological Societies (WHO/IUIS) Allergen Nomenclature Sub-committee (www.allergen.org) (Radauer et al., 2014). Among them, Phl p 1 and Phl p 5 are considered the most important, as a majority of IgE in timothy grass-allergic individuals is directed against these molecules (Rossi et al., 2008). A significant fraction of the *P. pratense* allergens were characterized in terms of their molecular and structural properties, with structures determined for Phl p 1, Phl p 2, Phl p 3, Phl p 4, Phl p 5, Phl p 6, and Phl p 7 (Zafred et al., 2013; Padavattan et al., 2009; Henzl et al., 2013; Gobl et al., 2017; Winter et al., 2020; Mitropoulou et al., 2018; Schweimer et al., 2008; De Marino et al., 1999).

The profilin from timothy grass, Phl p 12, has been registered as a minor allergen. In different populations of grass pollen allergic patients, up to 35% show IgE binding to Phl p 12 (Cudowska et al., 2020; Kowal et al., 2020). Although Phl p 12 is regarded as a minor allergen, it displays T cell response prevalence and strength comparable to Phl p 1, a major allergen from *P. pratense* (Lund et al., 2018). Phl p 12 has three registered isoallergens, and only one of them (Phl p 12.0101) was characterized at the molecular level (Cudowska et al., 2020).

Profilins are small, ubiquitous proteins found extensively in plants. Their primary function involves binding actin in plant cytoskeletons and controlling its polymerization and communication (Jimenez-Lopez et al., 2012; Jimenez-Lopez et al., 2013). Profilins are included in the panallergen classification due to their wide range of functions, high likelihood of IgE cross-reactivity, and conserved sequences and structures (McKenna et al., 2016). Profilins are particularly noted for their marked similarity between protein sequences, with 57 registered plant allergens ranging from 65 to 92% sequence identity. As a protein family, profilins are primarily minor allergens and are known for a high likelihood of co-sensitization as well as cross-reactivity situations caused by their sequence and structure similarity (Chruszcz et al., 2018; Offermann et al., 2016).

Here we present a crystal structure of Phl p 12.0101, which reveals that this allergen may form an unusual dimeric structure that was not previously observed among any profilins. The structure of Phl p 12.0101 is discussed in the context of allergenic sensitization, IgE binding, and allergy diagnostics, as well as in production of recombinant allergens.

MATERIALS AND METHODS

Protein purification and crystallization

Recombinant Phl p 12.0101 was produced and purified as previously described (Cudowska et al., 2020). Briefly, the Phl p 12.0101 construct with a cleavable N-terminal polyhistidine tag was produced in *E. coli* BL21(DE3) cells. The protein was purified using a combination of immobilized metal affinity chromatography and size exclusion chromatography. 20 mM β -mercaptoethanol (β -ME) was used during protein purification; however, it was not included in the final size exclusion chromatography step. Cleavage of the purification tag was performed according to the previously described protocol (Booth et al., 2018). Purified protein was stored in buffer containing 50 mM Tris-HCl, 150 mM NaCl, pH 7.4.

Protein was crystallized at 277 K over the course of one year. Using the vapor diffusion method in a 4 μ L drop, the best-diffracting crystals were obtained when 20 mg/mL protein solution was mixed 1:1 with crystallization solution containing 0.5 M sodium citrate at pH 6.5. Prior to data collection, crystals were cryocooled in liquid nitrogen.

Data collection, processing and structure determination

The South East Regional Collaborative Access Team beamline 22ID at the Advanced Photon Source (Argonne National Laboratory, Lemont, IL) was used to perform diffraction experiments. HKL-2000 software package was used for data processing (Otwinowski, 1997), and data collection statistics are reported in Table 1.

Phl p 12.0101 structure was solved by molecular replacement using MOLREP (Vagin & Teplyakov, 1997) integrated with HKL-3000 (Minor et al., 2006). The structure of maize profilin (Zea m 12; PDB code: 5FEF) was used as a search model. Initial model was rebuilt with BUCCANEER (Cowtan, 2006), COOT (Emsley & Cowtan, 2004), and HKL-3000. REFMAC (Murshudov et al., 2011) and COOT were used for structure refinement. Translation-Libration-Screw (TLS) Motion Determination server (Painter & Merritt, 2006) was used to divide the protein model into segments, and TLS parameters were implemented in the final stages of the refinement process. COOT and MOLPROBITY (Davis et al., 2007) were used for validation of the Phl p 12.0101 model. Refinement and validation statistics are summarized in Table 1. The Phl p 12.0101 structure together with its structure factors was deposited to the PDB with accession code 7KYW. Diffraction images were deposited in the Integrated Resource for Reproducibility in Macromolecular Crystallography (<https://proteindiffraction.org>; Grabowski et al., 2019; Grabowski et al., 2016; with <https://doi.org/10.18430/m3.irrmc.5715>).

Various computational approaches

PDBePISA was used to investigate an oligomeric form of Phl p 12.0101 in crystal state (Krissinel & Henrick, 2007). PDBeFOLD (Krissinel & Henrick, 2004) and DALI (Holm & Rosenstrom, 2010) servers were used to find proteins that have similar structures to Phl p 12.0101. Sequence alignment was prepared using Jalview (Waterhouse et al., 2009) and MUSCLE (Edgar, 2004). Figures showing protein structures were prepared with PYMOL (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.). APBS implemented in PyMOL was used for electrostatics calculations (Jurrus et al., 2018).

RESULTS AND DISCUSSION

Phl p 12.0101 – overall structure

Recombinant Phl p 12.0101 crystallized in $P3_121$ space group with one molecule in the asymmetric unit, and the protein model was refined at 2.3 Å resolution. The model contains all residues forming the mature form of the protein (residues 2–131), as well as glycine (residue 1) that corresponds to a fragment of the purification tag. The overall fold of the protein chain is very similar to that observed in other profilins that have their structures determined. The structure reported here superposes very well (over the whole sequence; rmsd values ~1 Å) with other models of plant profilins such as profilins 1–3 from *Arabidopsis thaliana* (Qiao et al., 2019, Thorn et al., 1997), *Hevea brasiliensis* (Hev b 8.0102) (Galicia et al., 2015; Mares-Mejia et al., 2016), muskmelon (Cuc m 2.0101) (Kapingidza et al., 2019), mugwort (Art v 4.0101), and ragweed (Amb a 8.0101) (Offermann et al., 2016).

The core of the molecule includes an antiparallel beta sheet composed of five strands ($\beta 2 \uparrow \beta 1 \downarrow \beta 7 \uparrow \beta 6 \downarrow \beta 5 \uparrow$), which is flanked from one side by two helices (H1 and H3), and from the other side by another helix (H2) and a hairpin motif ($\beta 3 \beta 4$) (Fig. 1). However, helix H3 in Phl p 12.0101 is shorter in comparison to the equivalent helices from plant profilins, such as that observed in mugwort profilin Art v 4.0101 (Fig. 1B). In Phl p 12.0101, an initial fragment of helix H3 is unwound and Cys115, which in other plant profilin structures is pointing toward the protein core, is facing toward the protein surface.

The structural analysis revealed that both Cys13 and Cys115 form intermolecular disulfide bridges (Cys13-Cys115' and Cys115-Cys13'; Fig. 2) that are responsible for Phl p 12.0101 dimer formation (Fig. 3). Symmetry of the dimer coincides in this case with crystal symmetry. The dimer interface has an area of 709 Å², and this molecular assembly, in addition to the two covalent bonds, is stabilized by a mixture of hydrophobic and hydrogen bonding interactions. Each molecule forming the dimer contributes approximately 15 residues to the interface, and most of these residues are highly conserved among profilins that are reported as allergens. However, the presence of dimerization in the crystal structure of Phl p 12.0101 was unexpected, as initial stages of protein purification were performed in reducing conditions, and the protein was determined to be monomeric prior to crystallization (Cudowska et al., 2020). Most likely, oxidation of the protein during the crystallization process, which took several months, resulted in dimer formation.

Phl p 12 has three officially registered isoallergens that have the same length and vary by just a few residues. Phl p 12.0102 can be considered a Phl p 12.0101 mutant in which arginine that is present in the 12.0101 isoallergen is substituted by alanine. Phl p 12.0103 has four differing residues from Phl p 12.0101 (97% sequence identity), including R81A as seen in Phl p 12.0102. Phl p 12.0101 and Phl p 12.0103 also feature differences between leucine and phenylalanine (L26F and F54L) and one change from glycine to alanine (G69A) in comparison to Phl p 12.0101 and Phl p 12.0102. Both cysteines (C13 and C115) found in Phl p 12.0101 are found in Phl p 12.0102 and Phl p 12.0103, as well as other residues forming the dimer interface, indicating that these isoallergens may form dimers identical to the one observed in Phl p 12.0101 crystal.

Comparison of Phl p 12.0101 with other profilins

As already mentioned, the overall fold of Phl p 12.0101 is very similar to that observed for other profilins. However, the observed conformation of Phl p 12.0101 fragment containing Cys115 is very unusual and was not reported previously (Fig. 1). Similarly, the participation of the conserved Cys115 (Fig. 2) in formation of an oligomeric assembly in plant profilins was also not observed. At the same time, profilins have been shown to exist as monomers or higher-order oligomers (Mares-Mejia et al., 2016; Mittermann et al., 1998; Wopfner et al., 2002; Willeroider et al., 2003). It was suggested that this may be accomplished through disulfide bond formation between two or more identical profilin molecules. Phl p 12.0101 as presented here is the second profilin allergen with its structure solved as a homodimer, following Hev b 8.0102 (Mares-Mejia et al., 2016). However, there are significant differences between the dimers formed by Phl p 12.0101 and Hev b 8.0102. For example, while the Phl p 12.0101 dimer is stabilized by two disulfide bridges, only one such covalent link is present in Hev b 8.0102 (Fig. 3). In addition, despite the fact that most of the residues forming the dimer interfaces originate from the N- and C-terminal fragments of the protein chains (including helices H1 and H3), the arrangements of the molecules in the dimers are different (Fig. 3). At the same time, residues that mediate dimer formation in Hev b 8.0102 and Phl p 12.0101 are highly conserved, suggesting that other plant profilins may form such oligomeric assemblies.

Plant profilins that are reported as allergens have at least two cysteine residues. Until now, all structural studies indicated that only residues equivalent to Cys13 of Phl p 12.0101 were exposed to solvent. The solvent-exposed thiol group was shown to be prone to modifications and disulfide bond formation, like in the case of Hev b 8.0102 (Mares-Mejia et al., 2016). Moreover, reducing agents such as 2-mercaptoethanol (β -ME) and dithiothreitol (DTT) were used in purification of these proteins and may have an effect on profilin stability (Soh et al., 2017; Mittermann et al., 1998). Under reducing conditions, profilin exists nearly exclusively in the monomeric form (Cudowska et al., 2020; Kapingidza et al., 2019; Offermann et al., 2016). However, when the profilin is under non-reducing conditions (in buffer without the reducing agent), the profilin has the potential to oligomerize. Bet v 2, the profilin from birch, was shown in SDS-PAGE to form dimers and trimers, with monomers in higher concentration than the dimeric or trimeric form (Mittermann et al., 1998). In non-reducing conditions, Hev b 8 forms primarily monomers and dimers, with the suggestion that it forms transient oligomers during gel filtration (Mares-Mejia et al., 2016). Art v 4 was shown to

form dimers, trimers, and tetramers under non-reducing conditions (Wopfner et al., 2002). Art v 4 has an extra cysteine residue (C95) in addition to the conserved C13 and C115 observed in Phl p 12 (Fig. 1), Hev b 8, and Bet v 2, possibly making oligomerization easier for Art v 4 in comparison to the other profilins. Similar to Hev b 8, varying oligomerization was observed when Art v 4 tetramers were converted back to stable dimers upon the addition of the reducing agent DTT (Wopfner et al., 2002). Interestingly, in contrast to other profilins that seem to exist primarily as monomers, Cap a 2 (*Capsicum annuum*, various peppers) was found almost exclusively in dimeric form with size exclusion chromatography, and the addition of DTT acted only partially to convert it to a monomer (Willerroider et al., 2003).

Profilin oligomerization in the context of biological function

Profilins interact with many proteins, and some of these interactions are mediated by binding to sequences that include proline-rich regions (Witke, 2004). It was shown in the dimeric structure of Hev b 8.0102 that the regions responsible for binding of proline-rich peptides partially overlap with the oligomerization interface (Kapingidza et al., 2019), which is also true for the Phl p 12.0101 dimer. Superposition of the profilin dimer structures with a structure of Amb a 8 in complex with polyproline (PDB code: 5EV0) and with a complex of human profilin-1:poly-Pro:actin (PDB code: 3CHW; Fig. 4) suggests that the dimers may potentially interact with relatively short proline-rich peptides (~ six residues) through the part of the binding site available in monomeric forms of the proteins. However, there is currently no information on the ability of the profilin dimers to interact with proline-rich peptides. If profilins can oligomerize in cells (Babich et al., 1996), it is plausible that the different forms of these proteins may have modified affinity towards binding partners (Kapingidza et al., 2019). Figure 4 also shows that dimeric assemblies of Hev b 8.0102 and Phl p 12.0101 are most likely not able to bind actin. In this case, more potential clashes are observed for the Hev b 8 dimer, due to its bent structure (Fig. 3).

Plant profilins are encoded by multigene families (Kovar et al., 2000). Therefore, several different isoforms of this protein may be present in the same organism, and the isoforms are not functionally alike. Phl p 12.0101 belongs to the profilin-1 class, while Hev b 8.0102 is a member of the profilin-2 class (Kovar et al., 2000). Therefore, one can speculate that the different dimeric assemblies observed for these proteins may be representative of different profilin classes. Despite the structural differences, there are large patches of surface that are negatively charged in both dimers (Fig. 3) which can be potentially involved in interactions with positively charged molecules. Similarly, positively charged surface areas may participate in interactions with negatively charged molecules. For example, it was demonstrated that profilins can interact with phosphatidylinositol 4,5-bisphosphate (PIP2) and inhibit hydrolysis of this molecule by phospholipase C (Goldschmidt-Clermont et al., 1990). According to UniProt (The UniProt, 2017), the Phl p 12.0101 region responsible for PIP2 binding corresponds to the Arg81-Thr97 fragment that contains several positively charged residues. Analysis of the profilin dimers' structures indicates that the putative PIP2 binding site is relatively far from the dimerization interfaces, and therefore the dimeric Hev b 8.0102 and Phl p 12.0101 may interact with the lipid.

Structure of Phl p 12.0101 in relation to allergy

Although plant profilins recognized as allergens are considered to exist mainly as monomeric proteins, it was demonstrated that oligomerization has an impact on their allergenicity. For example, it was shown that Hev b 8.0102 dimers and monomers were cross-reactive when tested with an IgE mAb (Mares-Mejia et al., 2020); however, higher basophil degranulation in a rat basophilic leukemia cell line was observed with the dimeric form of the allergen as compared to the monomer due to increased cross-linking between bound mAb and FcεRI receptors (Mares-Mejia et al., 2020). This may be related to the structure of FcεRI-IgE complexes and the distance between cross-linked IgE molecules required for optimal mast cell activation (Knol, 2006). Moreover, it has recently been demonstrated for polcalcins that multivalency of oligomers can overcome the need for high-affinity interactions necessary for effective mast cell activation (Bucaite et al., 2019). Dimer formation of Hev b 8.0102 also exposed known IgE-binding epitopes on the surface of the homodimer (Mares-Mejia et al., 2016), leaving the dimer able to bind more IgE than the monomer. A similar situation with changed IgE-binding epitopes may be possible with other allergenic profilins that can form dimers, such as Phl p 12, Cuc m 2 and Amb a 8.

While there is experimental evidence pointing to differences between IgE binding of monomeric and dimeric forms of profilins, it is still not clear whether the human body is exposed to the monomeric form of these allergens or a mixture of monomeric and oligomeric forms. Interestingly, reported clinical significance of sensitization to profilins differs in individual patients, ranging from profilin acting as a molecule of no clinical relevance to an active aeroallergen and to a trigger of a systemic anaphylaxis (Rodríguez Del Río et al., 2018). The variable clinical response in patients sensitized to profilins may depend on co-factors which can possibly affect the ability of those molecules to be recognized by IgE on mast cells and basophils (Rodríguez Del Río et al., 2018). Moreover, it must be stressed that the process of profilin purification can introduce a significant bias and result in samples that do not well represent the relative abundance of different oligomeric forms observed in the starting source material. In fact, profilins are often purified from natural sources using protocols that involve denaturing conditions and refolding. Such protocols may lead to formation of oligomeric assemblies that are mediated by intermolecular disulfide bridges. On the other hand, use of affinity columns that contain poly-Pro may lead to selection of profilin forms that have high affinity toward the proline-rich peptides. Additional complications can be related to the presence or absence of various reducing agents that can clearly shift equilibrium between monomeric and oligomeric forms of profilins. Therefore, purification and storage conditions of natural and recombinant profilins that are used in allergy research or diagnostics have to be taken into consideration.

Analysis of 69 plant profilins that are registered as allergens by WHO/IUIS shows that all of these proteins have two conserved Cys residues (Cys13 and Cys115 in Phl p 12.0101). In 15 of these proteins, there are three Cys residues (Ama r 2.0101, Amb a 8.0101, Amb a 8.0102, Art v 4.0101, Art v 4.0201, Hel a 2.0101, Hor v 12.0101, Koc s 2.0101, Pro j 2.0101, Sal k 4.0101, Sal k 4.0201, Tri a 12.0101, Tri a 12.0102, Tri a 12.0103, and Tri a 12.0104). Only two of these proteins have four Cys residues (Aca f 2.0101 and Mus a 1.0101; Fig. 5).

Crystal structures of Amb a 8.0101 and Art v 4.0101 revealed that the additional cysteine (Cys95 in Art v 4.0101; Fig. 5) forms an intramolecular disulfide bridge with Cys115 (Fig. 1B). Taking this into account, one may assume the presence of such an intramolecular disulfide bridge in Aca f 2.0101, Ama r 2.0101, Amb a 8.0102, Art v 4.0201, Art v 4.0201, Hel a 2.0101, Koc s 2.0101, Pro j 2.0101, Sal k 4.0101 and Sal k 4.0201, as they all have an equivalent of Cys95. Most likely this group of profilins does not form the dimeric assembly that was revealed for Phl p 12.0101, because formation of an intermolecular bridge would require breaking the intramolecular disulfide bond. In the case of Hor v 12.0101 and the Tri a 12 isoallergens, the additional cysteine residue precedes Cys13, and there is no experimental data on its involvement in formation of disulfide bridges. However, its localization suggest that the thiol group of this residue should be solvent-exposed and prone to modification.

Aca f 2.0101 has four cysteine residues, which follow the general pattern observed in Amb a 8.0101 and Art v 4.0101 with additional N-terminal Cys. Again, in this case it is not known whether the N-terminal cysteine participates in Aca f 2.0101 oligomerization. Similarly, it is not known whether the four cysteine residues present in the profilin from banana (Mus a 1.0101) participate in formation of intramolecular or intermolecular disulfide bridges.

In summary, the oxidation state of cysteine residues in profilins that were identified as allergens requires more study, as it may be important from the perspective of recombinant protein production and purification of profilins from natural sources, as well as for our understanding of allergy. Moreover, as recombinant proteins are used extensively in allergy diagnostics, the conditions of their storage should prevent unwanted modifications. In addition, characterization of allergenic profilins originating from natural sources is needed to provide information on posttranslational modifications and oligomeric states that are present in this group of proteins and that may presumably have impact on their interactions with the human immune system.

Acknowledgements of financial support:

The study was supported by N/ST/ZB/17/004/1143 grant from the Medical University of Białystok. A.O., A.B.K., and M.C. were partially supported by R01AI077653 grant from National Institute of Allergy and Infectious Diseases. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Structural results shown in this report are derived from data collected at Southeast Regional Collaborative Access Team (SER-CAT; 22 ID) beamline at the Advanced Photon Source, Argonne National Laboratory. Supporting institutions may be found at www.ser-cat.org/members.html. Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract Nos. DE-AC02-06CH11357 and W-31-109-Eng-38. This work was partially supported by an ASPIRE III grant from the Office of the Vice President of Research at the University of South Carolina.

Abbreviations:

β-ME	β-mercaptoethanol
IgE	immunoglobulin E
PDB	Protein Data Bank

REFERENCES

- Almeida E, Caeiro E, Todo-Bom A, Duarte A, Gazarini L (2019) Sensitization to grass allergens: Phl p1, Phl p5 and Phl p7 Phl p12 in adult and children patients in Beja (Southern Portugal), Allergol Immunopathol (Madr) 47: 579–584. 10.1016/j.aller.2019.04.006 [PubMed: 31477404]
- Babich M, Foti LR, Sykaluk LL, Clark CR (1996) Profilin forms tetramers that bind to G-actin. Biochem Biophys Res Commun 218: 125–131. 10.1006/bbrc.1996.0022 [PubMed: 8573117]
- Booth WT, Schlachter CR, Pote S, Ussin N, Mank NJ, Klapper V, Offermann LR, Tang C, Hurlburt KT, Chruszcz M (2018) Impact of an N-terminal polyhistidine tag on protein thermal stability. ACS Omega 3: 760–768. 10.1021/acsomega.7b01598 [PubMed: 29399652]
- Bucaite G, Kang-Pettinger T, Moreira J, Gould HJ, James LK, Sutton BJ, McDonnell JM (2019) Interplay between affinity and valency in effector cell degranulation: a model system with polcalcin allergens and human patient-derived IgE antibodies. J Immunol 203: 1693–1700. 10.4049/jimmunol.1900509 [PubMed: 31462504]
- Chruszcz M, Kapingidza AB, Dolamore C, Kowal K (2018) A robust method for the estimation and visualization of IgE cross-reactivity likelihood between allergens belonging to the same protein family. PLoS One 13: e0208276. 10.1371/journal.pone.0208276 [PubMed: 30496313]
- Cowtan K (2006) The Buccaneer software for automated model building. 1. Tracing protein chains. Acta Crystallogr D Biol Crystallogr 62: 1002–1011. 10.1107/S0907444906022116 [PubMed: 16929101]
- Cudowska B, Kapingidza AB, Pawłowicz M, Pampuch A, Hydeuk N, Pote S, Schlachter CR, Lebensztejn DM, Chruszcz M, Kowal K (2020) Production and use of recombinant profilins Amb a 8, Art v 4, Bet v 2, and Phl p 12 for allergenic sensitization studies. Molecules 25. 10.3390/molecules25020369
- Davis IW, Leaver-Fay A, Chen VB, Block JN, Kapral GJ, Wang X, Murray LW, Arendall WB 3rd, Snoeyink J, Richardson JS, Richardson DC (2007) MolProbity: all-atom contacts and structure validation for proteins and nucleic acids. Nucleic Acids Res 35: W375–W383. 10.1093/nar/gkm216 [PubMed: 17452350]
- De Marino S, Morelli MA, Fraternali F, Tamborini E, Musco G, Vrtala S, Dolecek C, Arosio P, Valenta R, Pastore A (1999) An immunoglobulin-like fold in a major plant allergen: the solution structure of Phl p 2 from timothy grass pollen. Structure 7: 943–952. 10.1016/s0969-2126(99)80121-x [PubMed: 10467147]
- Durham SR, Emminger W, Kapp A, de Monchy JG, Rak S, Scadding GK, Wurtzen PA, Andersen JS, Tholstrup B, Riis B, Dahl R (2012) SQ-standardized sublingual grass immunotherapy: confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. J Allergy Clin Immunol 129: 717–725 e715. 10.1016/j.jaci.2011.12.973 [PubMed: 22285278]
- Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32: 1792–1797. 10.1093/nar/gkh340 [PubMed: 15034147]
- Emsley P, Cowtan K (2004) Coot: model-building tools for molecular graphics. Acta Crystallogr D Biol Crystallogr 60: 2126–2132. 10.1107/S0907444904019158 [PubMed: 15572765]
- Galicía C, Mendoza-Hernández G, Rodríguez-Romero A (2015) Impact of the vulcanization process on the structural characteristics and IgE recognition of two allergens, Hev b 2 and Hev b 6.02, extracted from latex surgical gloves. Mol Immunol 65: 250–258. 10.1016/j.molimm.2015.01.018 [PubMed: 25700348]
- Göbl C, Focke-Tejkl M, Najafi N, Schrank E, Madl T, Kosol S, Madritsch C, Dorofeeva Y, Flicker S, Thalhamer J, Valenta R, Zangger K, Tjandra N (2017) Flexible IgE epitope-containing domains of Phl p 5 cause high allergenic activity. J Allergy Clin Immunol 140: 1187–1191. 10.1016/j.jaci.2017.05.005 [PubMed: 28532654]
- Goldschmidt-Clermont PJ, Machesky LM, Baldassare JJ, Pollard TD (1990) The actin-binding protein profilin binds to PIP2 and inhibits its hydrolysis by phospholipase C. Science 247: 1575–1578. 10.1126/science.2157283 [PubMed: 2157283]
- Grabowski M, Cymborowski M, Porebski PJ, Osinski T, Shabalin IG, Cooper DR, Minor W (2019) The integrated resource for reproducibility in macromolecular crystallography: experiences of the first four years. Struct Dyn 6: 064301. 10.1063/1.5128672 [PubMed: 31768399]

- 10.1107/S2059798316014716 (2016) A public database of macromolecular diffraction experiments. *Acta Crystallogr D Struct Biol* 72: 1181–1193. 10.1107/S2059798316014716 [PubMed: 27841751]
- Hatzler L, Panetta V, Lau S, Wagner P, Bergmann RL, Illi S, Bergmann KE, Keil T, Hofmaier S, Rohrbach A, Bauer CP, Hoffman U, Forster J, Zepp F, Schuster A, Wahn U, Matricardi PM (2012) Molecular spreading and predictive value of preclinical IgE response to *Phleum pratense* in children with hay fever. *J Allergy Clin Immunol* 130: 894–901 e895. 10.1016/j.jaci.2012.05.053 [PubMed: 22841010]
- Henzl MT, Sirianni AG, Wycoff WG, Tan A, Tanner JJ (2013) Solution structures of polyclonal Phl p 7 in three ligation states: Apo-, hemi-Mg²⁺-bound, and fully Ca²⁺-bound. *Proteins* 81: 300–315. 10.1002/prot.24186 [PubMed: 23011803]
- Hitchcock AS, Chase A (1951) *Manual of the grasses of the United States*. US Department of Agriculture
- Holm L, Rosenstrom P (2010) Dali server: conservation mapping in 3D. *Nucleic Acids Res* 38: W545–W549. 10.1093/nar/gkq366 [PubMed: 20457744]
- Jimenez-Lopez JC, Morales S, Castro AJ, Volkmann D, Rodríguez-García MI, Alché JD (2012) Characterization of profilin polymorphism in pollen with a focus on multifunctionality. *PLoS One* 7: e30878. 10.1371/journal.pone.0030878 [PubMed: 22348028]
- Jimenez-Lopez JC, Rodríguez-García MI, Alché JD (2013) Analysis of the effects of polymorphism on pollen profilin structural functionality and the generation of conformational, T- and B-cell epitopes. *PLoS One* 8: e76066. 10.1371/journal.pone.0076066 [PubMed: 24146818]
- Juruss E, Engel D, Star K, Monson K, Brandi J, Felberg LE, Brookes DH, Wilson L, Chen J, Liles K, Chun M, Li P, Gohara DW, Dolinsky T, Konecny R, Koes DR, Nielsen JE, Head-Gordon T, Geng W, Krasny R, Wei GW, Holst MJ, McCammon JA, Baker NA (2018) Improvements to the APBS biomolecular solvation software suite. *Protein Sci* 27: 112–128. 10.1002/pro.3280 [PubMed: 28836357]
- Kapingidza AB, Pye SE, Hyduke N, Dolamore C, Pote S, Schlachter CR, Commings SP, Kowal K, Chruszcz M (2019) Comparative structural and thermal stability studies of Cuc m 2.0101, Art v 4.0101 and other allergenic profilins. *Mol Immunol* 114: 19–29. 10.1016/j.molimm.2019.07.004 [PubMed: 31326654]
- Knol EF (2006) Requirements for effective IgE cross-linking on mast cells and basophils. *Mol Nutr Food Res* 50: 620–624. 10.1002/mnfr.200500272 [PubMed: 16764017]
- Kovar DR, Drøbak BIT, Staiger CJ (2000) Maize profilin isoforms are functionally distinct. *Plant Cell* 12: 583–598. 10.1105/tpc.12.4.583 [PubMed: 10760246]
- Kowal K, Pampuch A, Sacharzewska E, Swiebocka E, Siergiejko Z, Siergiejko G, Chruszcz M, DuBuske L (2020) Serum immunoglobulin E reactivity to cross-reacting panallergen components in northeastern Poland patients pollen sensitized. *Allergy Asthma Proc* 41: 183–191. 10.2500/aap.2020.41.200002 [PubMed: 32375962]
- Krissinel E, Henrick K (2004) Secondary-structure matching (SSM), a new tool for fast protein structure alignment in three dimensions. *Acta Crystallogr D Biol Crystallogr* 60: 2256–2268. 10.1107/S0907444904026460 [PubMed: 15572779]
- Krissinel E, Henrick K (2007) Inference of macromolecular assemblies from crystalline state. *J Mol Biol* 372: 774–797. 10.1016/j.jmb.2007.05.022 [PubMed: 17681537]
- Lund G, Brand S, Ramos T, Jimeno L, Boissy P, Vega F, Arina M, Christensen LH, Hoof I, Meno KH, Barber D, Blanco C, Würtzen PA, Andersen PS (2018) Strong and frequent T-cell responses to the minor allergen Phl p 12 in Spanish patients IgE-sensitized to Profilins. *Allergy* 73: 1013–1021. 10.1111/all.13351 [PubMed: 29121407]
- Maloney J, Bernstein DI, Nelson H, Creticos P, Hébert J, Noonan M, Skoner D, Zhou Y, Kaur A, Nolte H (2014) Efficacy and safety of grass sublingual immunotherapy tablet, MK-7243: a large randomized controlled trial. *Ann Allergy Asthma Immunol* 112: 146–153 e142. 10.1016/j.anai.2013.11.018 [PubMed: 24468255]
- Mares-Mejía I, García-Ramírez B, Torres-Larios A, Rodríguez-Hernández A, Osormo-Hernández AI, Terán-Olvera G, Ortega E, Rodríguez-Romero A (2020) Novel murine mAbs define specific and

- cross-reactive epitopes on the latex profilin panallergen Hev b 8. *Mol Immunol* 128: 10–21. 10.1016/j.molimm.2020.09.017 [PubMed: 33045539]
- Mares-Mejía I, Martínez-Caballero S, Garay-Canales C, Cano-Sánchez P, Torres-Larios A, Lara-González S, Ortega E, Rodríguez-Romero A (2016) Structural insights into the IgE mediated responses induced by the allergens Hev b 8 and Zea m 12 in their dimeric forms. *Sci Rep* 6: 32552. 10.1038/srep32552 [PubMed: 27586352]
- McKenna OE, Asam C, Araujo GR, Roulias A, Goulart LR, Ferreira F (2016) How relevant is panallergen sensitization in the development of allergies? *Pediatr Allergy Immunol* 27: 560–568. 10.1111/pai.12589 [PubMed: 27129102]
- Minor W, Cymborowski M, Otwinowski Z, Chruszcz M (2006) HKL-3000: the integration of data reduction and structure solution – from diffraction images to an initial model in minutes. *Acta Crystallogr D Biol Crystallogr* 62: 859–866. 10.1107/S0907444906019949/ [PubMed: 16855301]
- Mitropoulou AN, Bowen H, Dodev TS, Davies AM, Bax HJ, Bevil RL, Bevil AJ, Gould HJ, James LK, Sutton BJ (2018) Structure of a patient-derived antibody in complex with allergen reveals simultaneous conventional and superantigen-like recognition. *Proc Natl Acad Sci U S A* 115: E8707–E8716. 10.1073/pnas.1806840115 [PubMed: 30150373]
- Mittermann I, Fetrow JS, Schaak DL, Almo SC, Kraft D, Heberle-Bors E, Valenta R (1998) Oligomerization of profilins from birch, man and yeast. Profilin, a ligand for itself? *Sex Plant Rep* 11: 183–191. 10.1007/s004970050140
- Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, Winn MD, Long F, Vagin AA (2011) REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr* 67: 355–367. 10.1107/S0907444911001314 [PubMed: 21460454]
- Offermann LR, Schlachter CR, Perdue ML, Majorek KA, He JZ, Booth WT, Garrett J, Kowal K, Chruszcz M (2016) Structural, functional, and immunological characterization of Profilin panallergens Amb a 8, Art v 4, and Bet v 2. *J Biol Chem* 291: 15447–15459. 10.1074/jbc.M116.733659 [PubMed: 27231348]
- Ogle DG, John LS, Tilley DJ (2011) Plant guide for timothy (*Phleum pratense*), USDA-Natural Resources Conservation Service, Idaho State Office. Boise, ID.
- Otwinowski Z, Minor W (1997) Processing of X-ray diffraction data collected in oscillation mode. *Methods in Enzymology: Macromolecular Crystallography*, vol 276, part A, pp 307–326. Academic Press, New York
- Padavattan S, Flicker S, Schirmer T, Madritsch C, Randow S, Reese G, Vieths S, Lupinek C, Ebner C, Valenta R, Markovic-Housley Z (2009) High-affinity IgE recognition of a conformational epitope of the major respiratory allergen Phl p 2 as revealed by X-ray crystallography. *J Immunol* 182: 2141–2151. 10.4049/jimmunol.0803018 [PubMed: 19201867]
- Painter J, Merritt EA (2006) TLSMD web server for the generation of multi-group TLS models. *J Appl Cryst* 39: 109–111. 10.1107/S0021889805038987
- The PyMOL Molecular Graphics System, Version 2.0. Schrödinger, LLC;
- Qiao Z, Sun H, Ng JTY, Ma Q, Koh SH, Mu Y, Miao Y, Gao YG (2019) Structural and computational examination of the Arabidopsis profilin-Poly-P complex reveals mechanistic details in profilin-regulated actin assembly. *J Biol Chem* 294: 18650–18661. 10.1074/jbc.RA119.011307 [PubMed: 31653702]
- Radauer C, Nandy A, Ferreira F, Goodman RE, Larsen JN, Lidholm J, Pomés A, Raulf-Heimsoth M, Rozynek P, Thomas WR, Breiteneder H (2014) Update of the WHO/IUIS Allergen Nomenclature Database based on analysis of allergen sequences. *Allergy* 69: 413–419. 10.1111/all.12348 [PubMed: 24738154]
- Rodríguez Del Río P, Díaz-Perales A, Sánchez-García S, Escudero C, Ibáñez MD, Méndez-Brea P, Barber D (2018) Profilin, a change in the paradigm. *J Investig Allergol Clin Immunol* 28: 1–12. 10.18176/jiaci.0193
- Rossi RE, Monasterolo G, Prina P, Coco G, Operti D, Rossi L (2008) IgE profiles of Bermuda grass pollen sensitised patients evaluated by Phleum pratense allergens Phl P 1, 2, 4, 5, 6, 7, 11, 12. *Allergol Int* 57: 157–164. 10.2332/allergolint.O-07-503 [PubMed: 18349590]

- Scaparrotta A, Verini M, Consilvio NP, Cingolani A, Rapino D, Attanasi M, Cerasa M, Di Pillo S, Chiarelli F (2013) Sensitization to timothy grass pollen allergenic molecules in children. *Multidiscip Respir Med* 8: 17. 10.1186/2049-6958-8-17 [PubMed: 23448255]
- Schweimer K, Petersen A, Suck R, Becker WM, Rösch P, Matecko I (2008) Solution structure of Phl p 3, a major allergen from timothy grass pollen. *Biol Chem* 389: 919–923. 10.1515/BC.2008.102 [PubMed: 18627309]
- Sekerkova A, Polackova M, Striz I (2012) Detection of Phl p 1, Phl p 5, Phl p 7 and Phl p 12 specific IgE antibodies in the sera of children and adult patients allergic to Phleum pollen. *Allergol Int* 61: 339–346. 10.2332/allergolint.11-OA-0372 [PubMed: 22526205]
- Soh WT, Briza P, Dall E, Asam C, Schubert M, Huber S, Aglas L, Bohle B, Ferreira F, Brandstetter H (2017) Two distinct conformations in Bet v 2 determine its proteolytic resistance to cathepsin S. *Int J Mol Sci* 18. 10.3390/ijms18102156
- The UniProt C (2017) UniProt: the universal protein knowledgebase. *Nucleic Acids Res* 45: D158–D169. 10.1093/nar/gkw1099 [PubMed: 27899622]
- Thorn KS, Christensen HE, Shigeta R, Huddler D, Shalaby L, Lindberg U, Chua NH, Schutt CE (1997) The crystal structure of a major allergen from plants. *Structure* 5: 19–32. 10.1016/s0969-2126(97)00163-9 [PubMed: 9016723]
- Vagin A, Teplyakov A (1997) MOLREP: an automated program for molecular replacement. *J Appl Cryst* 30: 1022–1025. 10.1107/S0021889897006766.
- Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ (2009) Jalview Version 2 – a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25: 1189–1191. 10.1093/bioinformatics/btp033. [PubMed: 19151095]
- Willeroider M, Fuchs H, Ballmer-Weber BK, Focke M, Susani M, Thalhamer J, Ferreira F, Wüthrich B, Scheiner O, Breiteneder H, Hoffmann-Sommergruber K (2003) Cloning and molecular and immunological characterisation of two new food allergens, Cap a 2 and Lyc e 1, profilins from bell pepper (*Capsicum annuum*) and Tomato (*Lycopersicon esculentum*). *Int Arch Allergy Immunol* 131: 245–255. 10.1159/000072136 [PubMed: 12915767]
- Winter P, Stubenvoll S, Scheiblhofer S, Joubert IA, Strasser L, Briganser C, Soh WT, Hofer F, Kamenik AS, Dietrich V, Michelini S, Laimer J, Lackner P, Horejs-Hoeck J, Tollinger M, Liedl KR, Brandstetter J, Huber CG, Weiss R (2020) *In silico* Design of Phl p 6 variants with altered fold-stability significantly impacts antigen processing, immunogenicity and immune polarization. *Front Immunol* 11: 1824. 10.3389/fimmu.2020.01824 [PubMed: 33013833]
- Witke W (2004) The role of profilin complexes in cell motility and other cellular processes. *Trends Cell Biol* 14: 461–469 [PubMed: 15308213]
- Wopfner N, Willeroidee M, Hebenstreit D, van Ree R, Aalbers M, Briza P, Thalhamer J, Ebner C, Richter K, Ferreira F (2002) Molecular and immunological characterization of profilin from mugwort pollen. *Biol Chem* 383: 1779–1789. 10.1515/BC.2002.199 [PubMed: 12530543]
- Zafred D, Nandy A, Pump L, Kahlert H, Keller W (2013) Crystal structure and immunologic characterization of the major grass pollen allergen Phl p 4. *J Allergy Clin Immunol* 132: 696–703 e610. 10.1016/j.jaci.2013.03.021 [PubMed: 23683465]

Figure 1. Comparison of overall structures of Phl p 12.0101 (A) and Art v 4.0101 (PDB code: 5EV0) (B).

Both proteins are shown in cartoon representation. Cysteine residues are marked in magenta. Cys13 in the presented Art v 4.0101 structure is covalently modified by β -ME (Offermann et al., 2016).

Figure 2. Intermolecular disulfide bridge formed by two Phl p 12.0101 protein chains.
The electron density observed for this region of dimer interface is shown as a blue mesh (2Fo-Fc map, rmsd 1.5).

Figure 3. Comparison of dimeric assemblies formed by Phl p 12.0101 (A) and Hev b 8.0102 (B) (PDB code: 5FEG) (Mares-Mejia et al., 2016).

The models of proteins are shown in cartoon and surface representations. The models are shown in three different orientations, which correspond to rotations (0° – top, 90° – middle, and 180° – bottom) around a vertically oriented axis. Distributions of charges is mapped on molecular surfaces with positive charges marked in blue and negative charges marked in red. Cysteine residues are marked in magenta.

Figure 4. Superposition of dimers formed by Phl p 12.0101 (teal) (A) and Hev b 8.0102 (cyan) (B) on structure of tertiary complex between human profilin (magenta), a proline-rich peptide, and actin (blue) (PDB code: 3CHW).

The figure shows that in the case of dimeric profilin assemblies, binding of both actin and proline-rich peptide would most likely be obscured.

Figure 5. Sequence alignment of selected plant profilins that were identified as allergens and that have different content and/or distribution of cysteine residues.

Cysteine residues that are conserved in all officially registered allergens from the profilin family are marked with black arrows. Grey arrows mark positions of other cysteine residues that are not universally conserved in profilins. The sources of the profilins are as follows: Aca f – *Acacia farnesiana* (needle bush), Amb a – *Ambrosia artemisiifolia* (ragweed), Art v – *Artemisia vulgaris* (mugwort), Hev b – *Hevea brasiliensis* (para rubber tree), Hor v – *Hordeum vulgare* (barley), Tri a – *Triticum aestivum* (wheat), Phl p – *Phleum pratense* (timothy grass), and Mus a – *Musa acuminata* (banana).

Table 1.

Data collection and refinement statistics. Values in parentheses are for the highest resolution shell.

Protein	Phl p 12.0101
PDB accession code	7KYW
Data collection	
Diffraction source	APS, 22ID
Wavelength (Å)	1.0000
Space group	P3121
a, c (Å)	56.3, 94.4
Resolution range (Å)	40.00-2.30 (2.34-2.30)
No. of unique reflections	8119 (405)
Completeness (%)	99.8 (100.0)
Redundancy	10.4 (10.1)
$\langle I/\sigma(I) \rangle$	50.3 (2.2)
R _{meas}	0.066 (0.766)
R _{p.i.m}	0.021 (0.825)
CC1/2	0.942 (0.825)
Refinement	
Resolution range (Å)	33.94-2-30 (2.36-2.30)
Completeness (%)	99.7 (99.0)
No. of reflections, working set	7700 (556)
No. of reflections, test set	395 (29)
Final R _{cryst}	0.197 (0.302)
Final R _{free}	0.242 (0.400)
Rmsd bonds (Å)	0.014
Rmsd angles (°)	1.5
Ramachandran Plot	
Allowed regions (%)	98.0
Favored regions (%)	100.0